



[Course](#) > [Week 10](#) > [Practic...](#) > Q3: X-V...

Q3: X-Values

Problem 3: X Values


Part 1



0.0/6.0 points (ungraded)


Instead of the Bellman update equation, consider an alternative update equation, which learns the X value function. The update equation, assuming a discount factor $\gamma = 1$, is shown below:

$$X_{k+1}(s) \leftarrow \max_a \sum_{s'} T(s, a, s') [R(s, a, s') + \max_{a'} \sum_{s''} T(s', a', s'') [R(s', a', s'') + X_k(s'')]]$$

Assuming we have an MDP with two states, S_1, S_2 and two actions, a_1, a_2 , select the expectimax tree rooted at S_1 that corresponds to the alternative update equation







Submit

You have used 0 of 1 attempt

i Answers are displayed within the problem

Part 2

0.0/6.0 points (ungraded)

Select the mathematical relationship between the X_k -values learned using the alternative update equation and the V_k -values learned using a Bellman update equation.

- ☐ $X_k(s) = V_{k+1}(s)$
- ☐ $X_{k+1}(s) = V_k(s)$
- ☐ $X_k(s) = V_{k+2}(s)$
- ☐ $X_{k+2}(s) = V_k(s)$
- ☐ $X_k(s) = V_k(s)$
- ☐ $X_k(s) = V_k(V_{k+1}(s))$
- ☒ $X_k(s) = V_{2k}(s)$ ✓
- ☐ $X_{2k}(s) = V_k(s)$
- ☐ $X_k(s) = V_k(s) + V_{k+1}(s)$

☐ $X_k(s) = V_k(s) + V_k(s')$

Submit

You have used 0 of 1 attempt

i Answers are displayed within the problem