IRRATIONALITY MEASURES FOR $L(2,\chi_{-3})$ AND OTHER DIFFERENCES OF TRIGAMMA VALUES

SEAN LITSEY

ABSTRACT. We prove irrationality measures for several differences of values of the trigamma function ψ' at rational points. Among them are

$$\mu\big(L(2,\chi_{-3})\big) \le 13.7046, \qquad \mu\Big(L(2,\chi_{-3}) \pm \frac{\pi^2}{9}\Big) \le 9.4998,$$

$$\mu\Big(G \pm \frac{5\pi^2}{48}\Big) \le 31.2876,$$

and $\mu(\psi'(\frac{k}{5}) - \pi^2/6) \le 23.8272$ and $\mu(\psi'(\frac{k}{7}) - \pi^2/6) \le 2207.8697$ for k prime to the denominator. For $L(2,\chi_{-3})$ and $L(2,\chi_{-3}) \pm \pi^2/9$ the best bound known before was 24781, due to Calegari, Dimitrov and Tang. For $G \pm 5\pi^2/48$, $\psi'(\frac{k}{5}) - \pi^2/6$ and $\psi'(\frac{k}{7}) - \pi^2/6$ we know of no earlier proof of irrationality.

The approximations are determinants of rectangular Löwner matrices built from the digamma function at integer nodes. We bound their denominators prime by prime, using the corank of the matrix modulo p and cancellations at second and third order in its Smith normal form. Nonvanishing at rational points follows from Hermite interpolation in characteristic zero. The size of the determinants is bounded by Andréief's identity and an energy estimate. The proof is formalized in Lean 4.

CONTENTS

1. Introduction 2
2. Overview of the method 7
3. The construction 10
4. Arithmetic: the denominators 12
5. Nonvanishing 26
6. Analytic decay 27
7. The pair of shifts $(\frac{1}{3}, \frac{2}{3})$ 31
8. Proof of Theorem 1.1 34
9. Limits of the method 35
10. Formalization 36

Date: November 1, 2026.

2020 *Mathematics Subject Classification.* Primary 11J82; Secondary 11J72, 15A15, 31A15, 68V20.

Key words and phrases. Irrationality exponent, trigamma function, $L(2,\chi_{-3})$, Löwner matrices, p-adic valuations, Hermite interpolation, Andréief's identity, logarithmic energy, formal verification, Lean.

Disclosure of AI use 37
References 38

1. Introduction

1.1. **The result.** The values $\psi'(s) = \sum_{n\geq 0}(n+s)^{-2}$ of the trigamma function at rational points $s = k/q$ are among the simplest constants of analysis. Each is a rational linear combination of π^2 and the values $L(2,\chi)$ for the Dirichlet characters χ modulo q. For $q \leq 2$ they are rational multiples of π^2. Already for $q = 3$ and $q = 4$ they involve $L(2,\chi_{-3})$ and Catalan's constant $G = L(2,\chi_{-4})$, whose irrationality was proved only in 2024 and 2026 [6, 31]. Throughout the paper we work with the differences

$$\Psi(s,t) = \psi'(s) - \psi'(t) = \sum_{n=0}^{\infty}\Big(\frac{1}{(n+s)^2} - \frac{1}{(n+t)^2}\Big) \qquad (0 < s < t \leq 1).$$

We prove irrationality measures for $\Psi(s,t)$ for the pairs of Table 1, by a single determinant construction.

The *irrationality exponent* $\mu(\xi)$ of a real number ξ is the infimum of the real numbers μ such that $|\xi - a/b| > b^{-\mu}$ for all integers a and all sufficiently large integers b. It does not change under $\xi \mapsto r\xi + r'$ with $r, r' \in \mathbb{Q}$ and $r \neq 0$. A finite bound for $\mu(\xi)$ is an irrationality measure for ξ, and it shows that ξ is irrational.

(s,t)	k	classical form	μ
$(\frac{k}{2},1)$	1	π^2	6.5253
$(\frac{k}{3},1)$	1	$L(2,\chi_{-3}) + \frac{\pi^2}{9}$	9.4998
	2	$L(2,\chi_{-3}) - \frac{\pi^2}{9}$	9.4998
$(\frac{1}{3},\frac{2}{3})$	–	$L(2,\chi_{-3})$	13.7046
$(\frac{k}{4},1)$	1	$G + \frac{5\pi^2}{48}$	31.2876
	3	$G - \frac{5\pi^2}{48}$	31.2876
$(\frac{k}{5},1)$	$1,2,3,4$	–	23.8272
$(\frac{k}{7},1)$	$1,\ldots,6$	–	2207.8697

Table 1. The pairs of Theorem 1.1, grouped by the denominator of s. The classical form is a number in common notation that differs from $\Psi(s,t)$ by a rational affine map. A dash means that there is none. The last column is an upper bound for $\mu(\Psi(s,t))$.

The classical forms come from the values

$$\psi'(\tfrac14)=\pi^2+8G,\qquad \psi'(\tfrac34)=\pi^2-8G,$$
$$\psi'(\tfrac13)=\frac{2\pi^2}{3}+\frac92 L(2,\chi_{-3}),\qquad \psi'(\tfrac23)=\frac{2\pi^2}{3}-\frac92 L(2,\chi_{-3}),$$

together with $\psi'(\frac12)=\frac{\pi^2}{2}$ and $\psi'(1)=\frac{\pi^2}{6}$. Here

$$G=L(2,\chi_{-4})=\sum_{n\ge0}\Big(\frac{1}{(4n+1)^2}-\frac{1}{(4n+3)^2}\Big)=0.91596\,55941\ldots,$$
$$L(2,\chi_{-3})=\sum_{n\ge0}\Big(\frac{1}{(3n+1)^2}-\frac{1}{(3n+2)^2}\Big)=0.78130\,24128\ldots$$

are the values at 2 of the Dirichlet L-functions of the nontrivial characters modulo 4 and modulo 3. So Catalan's constant and $L(2,\chi_{-3})$ are the first two cases of the same family. In the notation of this paper

$$\Psi(\tfrac14,1)=8\Big(G+\frac{5\pi^2}{48}\Big),\qquad \Psi(\tfrac34,1)=-8\Big(G-\frac{5\pi^2}{48}\Big),\qquad \Psi(\tfrac14,\tfrac34)=16G,$$
$$\Psi(\tfrac13,1)=\frac92\Big(L(2,\chi_{-3})+\frac{\pi^2}{9}\Big),\qquad \Psi(\tfrac23,1)=-\frac92\Big(L(2,\chi_{-3})-\frac{\pi^2}{9}\Big),$$
$$\Psi(\tfrac13,\tfrac23)=9L(2,\chi_{-3}),\qquad \Psi(\tfrac12,1)=\frac{\pi^2}{3}.$$

For $q=5$ and $q=7$ there is no single classical constant. For $\gcd(k,q)=1$ the number $\Psi(\frac kq,1)$ is a rational linear combination of π^2 and the values $L(2,\chi)$ for the Dirichlet characters χ modulo q, since

$$\psi'(\tfrac kq)=q^2\sum_{n\equiv k\ (q)}\frac{1}{n^2}=\frac{q^2}{\varphi(q)}\sum_{\chi \bmod q}\bar\chi(k)L(2,\chi).$$

Theorem 1.1. *Let (s,t) be a pair in Table 1, and let $\mu_{s,t}$ be the entry in the last column of its row. For all integers a and all sufficiently large integers b,*

$$\Big|\Psi(s,t)-\frac ab\Big|>b^{-\mu_{s,t}}.$$

In particular $\Psi(s,t)$ is irrational and $\mu(\Psi(s,t))\le\mu_{s,t}$. The same bound holds for the classical form in the same row.

For $q=4$, 5 and 7 we know of no earlier proof that $\Psi(\frac kq,1)$ is irrational. For $q=4$ this does not follow from the irrationality of $G=\frac{1}{16}\Psi(\frac14,\frac34)$, since it is not known whether 1, G and π^2 are linearly independent over $\mathbb{Q}$. For $q=3$ irrationality was known [6], and the theorem improves the known measures. The bound for $q=2$ is weaker than the known bounds for π^2 and is included for completeness.

One family of determinants, with the same three estimates, covers every row of Table 1.

Theorem 1.1 has been verified in the Lean 4 proof assistant, together with every theorem, proposition and lemma used in its proof. The formal

proof has no unproved steps and uses only the standard axioms. Section 10 describes it. The remarks and Section 9 discuss the method and are not part of the formal proof. The formalization is available at https://github.com/slitsey/catalan-irrationality-measure. The use of AI in this work is disclosed at the end of the paper.

1.2. **Background.** *Catalan's constant.* In October 2026 OpenAI [31] proved that G is irrational and formalized the proof in Lean. Sun's preprint [38] had claimed a proof of the irrationality of G. Wachs [41] reports that its main estimate is not justified. Finite irrationality exponents for G were obtained after OpenAI's proof by Jha [17], Carreira [9], Bojnal [4], Xu [42] and the author [24]. Carreira [8] also records attempts to carry the determinant method to $\beta(4)$. For the arithmetic of G and related values before these results see [36, 45, 44, 20, 21] and the survey [48]. Eskandari, Murty and Nemoto [12] construct a 2-dimensional mixed motive over $\mathbb{Q}$ with period G. They use it to obtain linear forms in 1 and G. Eskandari [11] improves the approximation exponent of these forms of Rivoal–Zudilin type from 0.52 to 0.62. These are explicit approximations, not a lower bound for all rationals. Our searches found no proof that $G - 5\pi^2/48$ or $G + 5\pi^2/48$ is irrational.

The value $L(2, \chi_{-3})$. Calegari, Dimitrov and Tang [6] proved that 1, π^2 and $L(2, \chi_{-3})$ are linearly independent over $\mathbb{Q}$. Their proof applies a new arithmetic holonomy bound to a construction of Zagier. In [7] they made holonomy bounds effective. Theorem 5.1 there gives $\mu(L(2, \chi_{-3})) \leq 24781$. The remark that follows that theorem states that the same theorem and proof hold for any $\mathbb{Q}$-linear combination of $L(2, \chi_{-3})$ and π^2. So 24781 is also the known bound for $L(2, \chi_{-3}) \pm \pi^2/9$. Theorem 1.1 improves it to 13.7046 for $L(2, \chi_{-3})$ and to 9.4998 for $L(2, \chi_{-3}) \pm \pi^2/9$. Leibtag et al. [23] find numerically that Apéry-like approximations from conservative matrix fields have positive approximation exponents for $L(2, \chi_{-3}) \pm \pi^2/9$. A supplement to their paper uses one such field to give an Apéry-style proof of the theorem of Calegari, Dimitrov and Tang. The exponent it proves for each of these two numbers is only greater than 10^{-4}.

The numbers $\psi'(\frac{k}{5}) - \pi^2/6$ and $\psi'(\frac{k}{7}) - \pi^2/6$. Our searches found no irrationality result for them. Prellberg [33] proved that $\mathrm{Li}_2(1/5)$ is irrational, but that is a different number.

Recent results. Since September 2026 a series of new results has appeared in this area. Several of them were obtained with AI systems. They join classical results such as the bound $\mu(\log 2) < 3.5746$ of Marcovecchio [28], the bounds of Rhin and Viola for π^2 and $\zeta(3)$ [34, 35], which followed Apéry and Beukers [1, 3], and Zudilin's bound for π^2 [46]. OpenAI proved $\mu(\pi) = 2$, with a Lean formalization [32]. This improves the bound $\mu(\pi) < 7.1033$ of Zeilberger and Zudilin [43]. Liu, Jiang and Zhang proved $\mu(\log r) = 2$ for every rational $r > 0$ with $r \neq 1$ [25]. Their result also covers nonzero rational combinations of such logarithms. Giraudin lowered the bound for $\mu(\pi^2)$

to 5.0193784 with two hypergeometric constructions of Zudilin [46] at new parameters [15]. Giraudin also lowered the bound for $\mu(\zeta(3))$ to 5.5138800 by applying the Rhin–Viola method at a new integral direction [16]. Leibtag, Barkan, Kalisch, Weinbaum, Shalyt and Kaminer [23] showed that the classical constructions are one-dimensional trajectories in conservative matrix fields. By searching over other trajectories they obtained $\mu(\pi^2) < 4.62675$ and $\mu(\zeta(3)) < 5.4197$. Prellberg [33] proved that $\mathrm{Li}_2(-1/4)$, $\mathrm{Li}_2(1/5)$ and $\mathrm{Li}_2(-1/3)$ are irrational. These are three cases left open in [6]. The proof uses the integrals of Rhin–Viola and Viola–Zudilin. Fauzan [13] announced a proof that $\zeta(5)$ is irrational, using rationally normalized Hankel determinants. Lai [22] combined that method with simultaneous approximations of Zudilin and of Dauguet and Zudilin. Lai showed that 1, $\zeta(3)$ and $\zeta(4)$ are linearly independent over $\mathbb{Q}$.

Determinant methods. Irrationality proofs built on determinants of large size have a precedent in Zudilin's determinantal criterion with Hankel determinants [47]. The preprint of Liu, Jiang and Zhang on logarithms [25] uses a separated-weight interpolation determinant. The arguments of Fauzan and Lai [13, 22] use Hankel determinants. Brown [5] proposes a multivariate transfinite-diameter test for Mellin-type integrals and applies it to π^2. The framework we use is OpenAI's [31]. In it the decay of the determinants is controlled by Cauchy determinants and Andréief's identity, and the denominators are controlled prime by prime. Our determinants are Löwner matrices, that is, matrices of divided differences, of a digamma difference. They are a matrix-valued analogue of Padé and Hermite–Padé approximation [30]. Formal proofs of irrationality results in Lean also include the irrationality of $\zeta(3)$ by Liu, Zhang and Zhi [26].

1.3. **The companion paper.** Catalan's constant is also a difference of trigamma values, $G = \frac{1}{16}\Psi(\frac{1}{4}, \frac{3}{4})$, and the construction applies to this pair. For it, however, our bound for the denominators exceeds our bound for the decay, so we obtain no measure for G. The companion paper [24] refines OpenAI's determinant instead and proves $\mu(G) \le 410$. Our computations for $(\frac{1}{4}, \frac{3}{4})$ are not part of the formal proof.

The row for π^2 is included only to show the reach of the construction. Since $\zeta(2) = \pi^2/6$, much better bounds are known, the best being $\mu(\pi^2) < 4.62675$ of Leibtag et al. [23].

1.4. **What a measure requires.** Let $P(x)$ be a polynomial with rational coefficients for which $P(\xi)$ is small, and suppose that $|\xi - a/b|$ is very small. Three estimates turn this into a lower bound for $|\xi - a/b|$.

(i) *Denominators.* The coefficients of P have a common denominator of size $e^{(\Lambda+o(1))r^2}$.
(ii) *Smallness on a neighbourhood.* $|P(x)| \le e^{-(D-o(1))r^2}$ for every x with $|x - \xi| \le e^{-\rho r}$, not only at ξ.

(iii) *Nonvanishing at the point.* $P(a/b) \neq 0$ for some member of the family whose size can be chosen in terms of b.

In OpenAI's argument for G the size of the determinant is tied to a prime p, and nonvanishing is proved modulo p. Here nonvanishing is proved in characteristic zero, so the size r is free. The matrices are also simpler: their entries are divided differences of a single digamma combination. We prove the three estimates once, uniformly in the shift.

1.5. **New ideas.** The proof rests on the following ideas. Section 2 explains each of them in more detail.

(1) *Rectangular Löwner matrices.* The rows and columns share a long block of nodes $\{0, \dots, r-1\}$, which carries the variable. A further $m \approx cr/2$ row nodes and m column nodes alternate beyond it. The degree of the determinant in the variable is still r, while its decay is that of a larger node set. The border nodes never meet the diagonal, so the denominators grow less than the decay.
(2) *Denominators from local bounds.* For each prime we prove a list of upper bounds for the valuation of the common denominator. Each is an explicit integer expression in r, m, p and the first pole k_p. Integrated against $\log p$, their minimum gives the denominator exponent. Checking this minimum reduces to finitely many comparisons of rational numbers.
(3) *Primes smaller than the border.* When $p \leq 2m$ some residue classes contain two border nodes. Eliminating these pairs by Schur complements leaves a smaller matrix of the same shape, to which the separation argument for large primes applies at third order in the Smith normal form.
(4) *Nonvanishing in characteristic zero.* A kernel vector of the matrix at a rational point corresponds to a solution of a Hermite interpolation problem. A Wronskian argument bounds the corank independently of r, so a principal minor of bounded codimension does not vanish.
(5) *Decay through a trial measure.* Andréief's identity turns the determinant into an integral whose logarithm is a discrete logarithmic energy. Comparison with a piecewise polynomial trial measure bounds this energy, leaving a lower bound for one explicit function on finitely many segments.

The constants of the proof for each row of Table 1 are listed in Table 2.

1.6. **Organization.** Section 2 explains the construction and the main ideas without technicalities, and states the general deduction. Section 3 gives the construction. Sections 4, 5 and 6 prove the three estimates, for the denominators, nonvanishing and decay, uniformly in the shift. Section 7 treats the function F_χ of the pair $(\frac{1}{3}, \frac{2}{3})$. Section 8 deduces Theorem 1.1 and comments on the individual rows. Section 9 discusses the limits of the method. Section 10 describes the formalization. The disclosure of AI use follows it.

1.7. **Notation and conventions.** We write ψ for the digamma function, the logarithmic derivative of Euler's gamma function, and v_p for the p-adic valuation. For $P \in \mathbb{Q}[x]$, $\operatorname{den} P$ is the least positive integer whose product with P lies in $\mathbb{Z}[x]$. We write $[0,r) = \{0,1,\dots,r-1\}$. For finite sets X_0, Y_0 of integers and a matrix A indexed by integers, $A[X_0,Y_0]$ is the submatrix with rows X_0 and columns Y_0.

2. Overview of the method

2.1. **Löwner matrices of digamma differences.** Let F be a function analytic near a finite set of distinct points. The *Löwner matrix* [27] of F is the matrix of divided differences indexed by the points:

$$\mathsf{L}_F[t,t'] = \frac{F(t') - F(t)}{t' - t} \ (t \neq t'), \qquad \mathsf{L}_F[t,t] = F'(t).$$

It is a Cauchy-like structured matrix. Hermite's formula gives $\mathsf{L}_F = \frac{1}{2\pi i} \oint F(z)\, v(z) v(z)^\top dz$ with $v_t(z) = 1/(z-t)$. A representation of F as a Stieltjes or contour integral therefore exhibits L_F as a Gram matrix of Cauchy kernels.

For a rational shift $0 < s < 1$ with denominator q we take the digamma difference F_s. For the pair $(\frac{1}{3}, \frac{2}{3})$ we take a combination F_χ of three digamma functions. The nodes are integers. We put

$$\begin{gathered} F_s(z) = \psi(z+1) - \psi(z+s), \qquad \omega = \tfrac{1}{q^2}, \quad \xi_s = -\tfrac{1}{q^2}\Psi(s,1), \\ F_\chi(z) = -\tfrac{1}{3}\psi(z+\tfrac{1}{3}) + \tfrac{1}{2}\psi(z+\tfrac{1}{2}) - \tfrac{1}{6}\psi(z+1), \\ \omega = -\tfrac{1}{6}, \quad \xi_\chi = \tfrac{1}{36}\Psi(\tfrac{1}{3},\tfrac{2}{3}). \end{gathered} \tag{2.1}$$

In each case $\xi = \omega\, F'(0)$. The differences $F(t') - F(t)$ and $F'(t) - F'(0)$ are rational for integers $t, t' \geq 0$. Let R and Q be finite sets of nonnegative integer nodes, which index the rows and the columns. Then

$$\omega\, \mathsf{L}_F[R,Q] = M[R,Q] + \xi \operatorname{Id}_{R,Q}, \qquad \operatorname{Id}_{R,Q}[t,t'] = \mathbf{1}[t = t'], \tag{2.2}$$

with an explicit symmetric function M on $\mathbb{Z}^2_{\geq 0}$ with rational values, given in §3. Replacing ξ by a variable x gives the matrix polynomial $\Phi(x) = M[R,Q] + x\operatorname{Id}_{R,Q}$. Its determinant $P(x)$ has rational coefficients and degree at most $|R \cap Q|$, and $P(\xi) = \det(\omega\, \mathsf{L}_F[R,Q])$.

Suppose that $|\xi - a/b|$ is very small. Then $P(a/b)$ is very close to $P(\xi)$, which is tiny. On the other hand $b^{|R\cap Q|} \operatorname{den}(P)\, P(a/b)$ is an integer. If it is not zero, we get a contradiction as soon as

$$\underbrace{\log \operatorname{den} P}_{\text{denominators}} + |R \cap Q| \log b \ < \ \underbrace{-\log |P(a/b)|}_{\text{decay}}.$$

Three estimates are therefore needed: an upper bound for $\log \operatorname{den} P$, the *arithmetic estimate*; an upper bound for $|P(x)|$ near ξ, the *analytic estimate* or decay; and the *nonvanishing* of $P(a/b)$ for some member of a family of determinants whose size can be chosen in terms of b. Each estimate is proved once, for every shift, and the families differ only in a few parameters.

2.2. **Rectangular Löwner matrices with an alternating border.** We take a long common block $C = [0, r)$, which carries the variable x, and add m row nodes and m column nodes that alternate beyond it:

$$X' = \{r, r+2, \dots, r+2m-2\}, \qquad Y' = \{r+1, r+3, \dots, r+2m-1\}.$$

The rows are $R = C \sqcup X'$ and the columns are $Q = C \sqcup Y'$ (Figure 1). The determinant $P(x) = \det \Phi(x)$ has degree at most r, since the variable occurs only on the diagonal of the common block. So the factor $b^{\deg P}$ in the contradiction is that of r nodes, while the matrix has size $r+m$ and its decay is that of about $(1+c/2)r$ nodes, where $c = 2m/r$ is the length of the border relative to r. The denominators grow too, but less, because border rows and border columns never share a node. To guarantee nonvanishing we also use the principal minors that delete a bounded number of common nodes. A larger c raises D and Λ together, and we chose c numerically to minimise $\rho/(D-\Lambda)$. Its value for each constant is given in Table 2.




FIGURE 1. The geometry for $r = 8$ and $m = 3$. Rows are indexed by $C \sqcup X'$ and columns by $C \sqcup Y'$. The variable x occurs only on the diagonal of the common block. The border rows $X' = \{r, r+2, r+4\}$ and border columns $Y' = \{r+1, r+3, r+5\}$ alternate and never meet the diagonal.

The interleaving of X' and Y' is essential. It keeps the two Cauchy determinants in Andréief's formula comparable (§6). It also gives the reduction of the matrix modulo a prime an alternating structure. Several of the arithmetic bounds of §4 rest on this structure.

2.3. **Denominators from local bounds.** As in the work of Rhin and Viola [34, 35], we bound the p-adic valuation of the common denominator prime by prime and sum the bounds with the prime number theorem in arithmetic progressions. For $p \asymp r$ the valuation depends on r, m, p and the first pole k_p, the least index $k_p \geq 1$ at which the second digamma term has a pole modulo p. One has $k_p = \kappa p + O(1)$ with $\kappa = j/q$, where the

residue $j \equiv qs\,p^{-1} \pmod q$ is the *root class* of p. The function F_χ has two first poles, and on each class of primes modulo 6 both are affine functions of one index of the same kind (Section 7).

The local bounds come from three sources: the valuations of the entries, the corank of the reduction modulo p, and cancellations at second and third order in the Smith normal form. Each is an explicit piecewise linear expression in (r, m, p, k_p), valid in a range of p/r described by linear inequalities. Replacing (r, m, p, k_p) by $(1, \alpha, \theta, \kappa\theta)$, with $\alpha = c/2$, turns it into a piecewise affine function of $\theta = p/r$, with an explicitly bounded error. The *denominator exponent* is

$$\Lambda = \frac{1}{\varphi(q)} \sum_j \int_0^\infty \mathcal{P}_j(\theta)\, d\theta,$$

where $\mathcal{P}_j$ is the minimum of these functions for the root class j. A table of affine pieces above $\mathcal{P}_j$ certifies Λ by finitely many comparisons of rational numbers (Theorem 4.14). The bounds depend on the shift only through the root class, so one table serves every shift with denominator q, and the numbers $\Psi(s, 1)$ for these shifts receive the same exponent.

2.4. **Primes smaller than the border.** For $p \le 2m$ some residue classes modulo p contain two border nodes, a row t and a column $t + p$. The bounds above are not sharp for these primes. Each such pair gives a pivot that is a p-adic unit, because the interval $(t, t + p]$ contains one pole of each digamma term and their contributions cannot cancel. Eliminating these pivots leaves a matrix of the same shape with a border of $d = \|m\|_p$ rows and columns. To this smaller matrix we apply the separation argument used for large primes, at third order. Two identities for bordered inverses show that the elimination does not disturb the second-order terms (Lemma 4.9). A rational interpolation problem of type (n, n) shows that the remaining border nodes stay separated (Lemma 4.11).

2.5. **Nonvanishing.** Let x_0 be a rational point. The kernel of $\Phi(x_0)$ is identified with a space of pairs of polynomials that satisfy Hermite interpolation conditions at the nodes. Two such pairs have vanishing cross product, by a count of degrees and zeros. A further pair in the kernel would give a rational antidifference of the increment of F. No such antidifference exists, because the pole at -1 is isolated in its class modulo $\mathbb{Z}$. This bounds the corank of $\Phi(x_0)$ by a number that depends only on F and on the alternation of the border. A principal restriction that deletes at most that many common nodes is then nonsingular. This is Theorem 5.1. No prime enters, so the size r can be chosen freely in terms of b.

2.6. **Decay.** Moving Hermite's contour to a polygonal path Γ in the left half-plane writes $\omega \mathbin{\mathsf{L}}_F [R, Q]$ as a Gram matrix $\int v_t v_{t'}\, d\Omega$. Here Ω is a complex measure. It has signed atoms at poles of F and an exponentially small

density on Γ. Andréief's identity with two families of functions, one for the rows and one for the columns, gives

$$|\det \mathsf{L}_F[R,Q]| \le \frac{1}{n_*!}\int |\mathsf{C}_R(z)|\,|\mathsf{C}_Q(z)| \prod_{i=1}^{n_*} d|\Omega|(z_i),$$

where n_* is the size of the matrix and C_R, C_Q are Cauchy determinants. Taking logarithms turns the integrand into a discrete logarithmic energy. The free points z_i interact with the nodes in an external field.

We bound this energy from below by comparing it with a *trial measure* σ, a density that is polynomial on each of finitely many segments. The comparison uses the conditional positive definiteness of the logarithmic kernel. What is left is a lower bound for one explicit function Υ on the contour, on the segment carrying the atoms and on the interval of the nodes. We prove it on a finite cover of these sets by cells, each checked in exact rational arithmetic. One theorem, Theorem 6.2, gives the decay for every geometry parameter $c \le 1$ and every function F whose poles are separated from the contour. Each family then supplies only σ, the polygon and the cover.

2.7. **The deduction.** Write $\eta = D - \Lambda$, where D is the *decay rate*, and let ρ be the *neighbourhood exponent* of the decay estimate: the decay holds at every point within $e^{-\rho r}$ of ξ. Section 8 combines the three estimates into

$$\mu(\xi) \le \frac{\rho}{\eta} = \frac{\rho}{D-\Lambda}.$$

This is the bound $d\rho/\eta$ of the companion paper [24] with degree $d = 1$ per unit of scale.

2.8. **Formal verification.** Every step is formalized in Lean 4 with Mathlib [10, 39]: the local arithmetic bounds and the conditions on the tables for Λ, the interpolation argument for nonvanishing, the contour representation, the cell checks of the decay estimate, and the prime number theorem in arithmetic progressions. The formal proofs share one core, which contains rectangular determinants, Gauss valuations, the local bounds, the tables for Λ, the selection in characteristic zero, the decay theorem and the deduction. A family supplies its function F, its geometry parameter and its numerical tables. The written proofs below are at the level of a research article. For long computations we point to the formal proof, which is the basis for the correctness of Theorem 1.1.

3. The construction

3.1. **The entries of M.** Let $0 < s < 1$ be rational with denominator q. Put $u = q(1-s)$, so that $k-1+s = (qk-u)/q$. For the function F_s of (2.1), the rational matrix M of (2.2) is as follows. Here $0 \le t < t'$ are integers,

and $M[t',t] = M[t,t']$.

$$M[t,t'] = \frac{1}{q^2(t'-t)} \sum_{k=t+1}^{t'} \Big(\frac{1}{k} - \frac{q}{qk-u}\Big),$$

$$M[t,t] = -\frac{1}{q^2} \sum_{k=1}^{t} \Big(\frac{1}{k^2} - \frac{q^2}{(qk-u)^2}\Big).$$

For $s = \frac{3}{4}$ this is $M[t,t'] = -\frac{1}{t'-t} \sum_{k=t+1}^{t'} \frac{1}{16k(4k-1)}$. For the function F_χ,

$$M[t,t'] = \frac{1}{6(t'-t)} \sum_{k=t+1}^{t'} g(k), \qquad M[t,t] = \frac{1}{6} \sum_{k=1}^{t} g'(k),$$

with $g(k) = \frac{1}{3k-2} - \frac{1}{2k-1} + \frac{1}{6k}$. Then $-\frac{1}{6}g'(k) = \frac{1}{2(3k-2)^2} - \frac{1}{3(2k-1)^2} + \frac{1}{36k^2}$.

We write each function as $F = \sum_{\varsigma\in\mathcal{T}} w_\varsigma \psi(z+\varsigma)$, where $\mathcal{T}$ is the set of shifts of F and $\sum_\varsigma w_\varsigma = 0$. For F_s, $\mathcal{T} = \{1, s\}$ with $w_1 = 1$ and $w_s = -1$. For F_χ, $\mathcal{T} = \{\frac{1}{3}, \frac{1}{2}, 1\}$. The number $|\mathcal{T}|$ of digamma terms is 2 for F_s and 3 for F_χ.

Lemma 3.1. *For each function of (2.1), $\omega \mathsf{L}_F[R,Q] = M[R,Q] + \xi \mathrm{Id}_{R,Q}$ for all finite sets $R, Q \subset \mathbb{Z}_{\geq 0}$, with ω and ξ as in (2.1).*

Proof. From $\psi(w+1) - \psi(w) = 1/w$ one gets $F_s(k) - F_s(k-1) = \frac{1}{k} - \frac{1}{k-1+s} = \frac{1}{k} - \frac{q}{qk-u}$. Telescoping gives the off-diagonal entries. In the same way $\psi'(w+1) - \psi'(w) = -1/w^2$ gives $F_s'(t) - F_s'(0)$. Finally $\omega F_s'(0) = q^{-2}(\psi'(1) - \psi'(s)) = -q^{-2}\Psi(s,1) = \xi_s$. For F_χ the entries are computed in the same way. For the value at 0, the multiplication formula $\sum_{i<n} \psi'(z + \frac{i}{n}) = n^2\psi'(nz)$ gives $\psi'(\frac{1}{2}) = 3\psi'(1)$ and $\psi'(\frac{1}{3}) + \psi'(\frac{2}{3}) = 8\psi'(1)$. Hence $F_\chi'(0) = -\frac{1}{3}\psi'(\frac{1}{3}) + \frac{1}{2}\psi'(\frac{1}{2}) - \frac{1}{6}\psi'(1) = -\frac{1}{6}\Psi(\frac{1}{3}, \frac{2}{3})$, and $-\frac{1}{6}F_\chi'(0) = \frac{1}{36}\Psi(\frac{1}{3}, \frac{2}{3}) = \xi_\chi$. □

Since μ is invariant under rational affine maps, Theorem 1.1 is a statement about the numbers ξ of (2.1).

3.2. **Geometries.**

Definition 3.2. Let $r, m \geq 0$. The *geometry* $\mathcal{G}(r,m)$ consists of the common nodes $C = [0,r)$, the row nodes $X' = \{r + 2i : 0 \leq i < m\}$ and the column nodes $Y' = \{r + 2i + 1 : 0 \leq i < m\}$. Rows are indexed by $R = C \sqcup X'$ and columns by $Q = C \sqcup Y'$. Let $I \subseteq C$ be a set of deleted common nodes. The matrix

$$\Phi_{\mathcal{G},I}(x) = \big(M[t,t'] + x\,\mathbf{1}[t = t' \in C \setminus I]\big)_{t\in R\setminus I,\ t'\in Q\setminus I}$$

is square of size $|C \setminus I| + m$. Put $P_{\mathcal{G},I}(x) = \det \Phi_{\mathcal{G},I}(x) \in \mathbb{Q}[x]$. We write $\Phi_\mathcal{G}$ and $P_\mathcal{G}$ when $I = \emptyset$, and we write $N = r + 2m$ for the number of nodes.

By Lemma 3.1, $\Phi_\mathcal{G}(\xi + h) = \omega \mathsf{L}_F[R,Q] + h\mathrm{Id}_{R,Q}$ for real h. The geometry with $m = 0$ is the square Löwner matrix of size r.

Definition 3.3. Let $0 < c \le 1$ and $\alpha = c/2$. The *alternating geometry* of parameter c at an even scale r is $\mathcal{G}(r, m(r))$ with $m(r) = \lceil \alpha r - \frac{1}{2} \rceil$. Then $|m(r) - \alpha r| \le \frac{1}{2}$, and the border $[r, r + 2m)$ has length $cr + O(1)$. For a depth $K \ge 0$, its *members* are the matrices $\Phi_{r,I} = \Phi_{\mathcal{G},I}$ with $|I| \le K$. We put $P_{r,I} = \det \Phi_{r,I}$.

For the functions F_s we use $c = \frac{3}{10}, \frac{1}{5}, \frac{3}{10}, \frac{2}{9}, \frac{1}{6}$ for $q = 2, 4, 3, 5, 7$. For F_χ we use $c = \frac{1}{2}$, as explained in Section 7. All $o(1)$ terms below are uniform over the members as $r \to \infty$.

Lemma 3.4 (integrality)**.** *$P_{\mathcal{G},I}$ has degree at most $|C \setminus I|$. If a and $b \ge 1$ are integers with $P_{\mathcal{G},I}(a/b) \ne 0$, then $|\operatorname{den}(P_{\mathcal{G},I})\, P_{\mathcal{G},I}(a/b)| \ge b^{-|C \setminus I|}$.*

Proof. The variable occurs only in the $|C \setminus I|$ diagonal entries indexed by $C \setminus I$. So each Leibniz term has degree at most $|C \setminus I|$. Then $b^{|C \setminus I|} \operatorname{den}(P_{\mathcal{G},I})\, P_{\mathcal{G},I}(a/b)$ is a nonzero integer. □

Example. Take $s = \frac{3}{4}$, $r = 2$ and $m = 1$, so that $C = \{0, 1\}$, $X' = \{2\}$ and $Y' = \{3\}$. Then

$$\Phi_{\mathcal{G}}(x) = \begin{pmatrix} x & -\frac{1}{48} & -\frac{67}{7392} \\ -\frac{1}{48} & x + \frac{7}{144} & -\frac{47}{14784} \\ -\frac{17}{1344} & -\frac{1}{224} & -\frac{1}{528} \end{pmatrix}$$

and

$$P_{\mathcal{G}}(x) = -\frac{x^2}{528} - \frac{823\,x}{3725568} - \frac{239}{37158912}.$$

The variable occurs only at the common nodes 0 and 1, so the degree is $|C| = 2$ although the matrix has size 3. When $M[X', Y']$ is invertible, a Schur complement shows that $P_{\mathcal{G}}$ is, up to a constant factor, the characteristic polynomial of a rational matrix of size $|C|$, as in the square case. We do not use this factorisation.

4. Arithmetic: the denominators

In this section we bound the common denominator of the coefficients of the members $P_{r,I}$ for the functions F_s. The method is that of Rhin and Viola [34, 35]: the p-adic valuation of the denominator is bounded prime by prime, and the bounds are summed against $\log p$ with the prime number theorem in arithmetic progressions. What is new is the form of the local bounds. Each is an explicit integer expression in four integers attached to the prime, and their minimum, integrated, gives the denominator exponent.

The local bounds come from four mechanisms, of increasing depth.

(a) *Valuations of the entries.* The assignment bounds (T1)–(T5) of §4.4.1 use only the valuations of single entries and LP duality.

(b) *Corank modulo p.* The bounds (LP1), (LP2) and (R) of §§4.4.2 and 4.4.4 compute the corank of the reduction of a rescaled matrix.

(c) *Second order.* The bounds (W) and (S) of §§4.4.3 and 4.4.5 use the pairing of kernel vectors modulo p^2 (§4.3).
(d) *Third order.* The bounds (U_n) of §4.4.6 treat the primes smaller than the border.

Comparing the local bounds with a table of affine pieces is a finite computation with rational numbers, and Theorem 4.14 turns an admissible table into the arithmetic estimate. Section 7 treats the function F_χ.

4.1. **The root class of a prime.** Fix the function F_s, with s of denominator q, and the alternating geometry of parameter $c = 2\alpha$. Let $p > q$ be a prime. The indices $k \geq 1$ at which the entries of M acquire a denominator divisible by p are the multiples of p and the solutions of $qk \equiv u \pmod p$. Let k_p be the least of the latter. Then

$$qk_p = jp + u, \qquad 0 < j < q, \qquad j \equiv qs\,p^{-1} \pmod q.$$

We call j the *root class* of p and put $\kappa = \kappa_j = j/q$. Then $k_p - \kappa p = u/q \in [\frac{1}{q}, 1-\frac{1}{q}]$. So the first pole is at distance less than one from κp, uniformly in s. Throughout we assume $q(r+2m) < p^2$. Then every pole below $\bar{N}$ is simple, and only the first power of p matters. The primes with $p^2 \leq q(r+2m)$ and the prime powers contribute $O(r^{3/2}\log r)$ in total.

4.2. **Gauss valuations, the assignment bound and the rank of the reduction.** Fix a prime p. Let $\mathcal{O} \subset \mathbb{Q}(x)$ be the localisation of $\mathbb{Z}_{(p)}[x]$ at the prime ideal (p). It is a discrete valuation ring with uniformiser p and residue field $\mathbb{F}_p(x)$. Its valuation v, restricted to $\mathbb{Q}[x]$, is the *Gauss valuation* $v(f) = \min_k v_p(f_k)$ for $f = \sum f_k x^k$. By Gauss's lemma $v(fg) = v(f) + v(g)$, and $v_p(\operatorname{den} f) = \max(0, -v(f))$. For a member we write $e_p(r, I) = v_p(\operatorname{den} P_{r,I})$.

Lemma 4.1 (assignment bound and LP duality)**.** *Let A be a square matrix over $\mathbb{Q}[x]$ with rows X_0 and columns Y_0. Let $W_{tt'} = \max(0, -v(A_{tt'}))$, and let $\tau(W)$ be the largest weight $\sum W_{tt'}$ of a perfect matching between X_0 and Y_0. Then $v(\det A) \geq -\tau(W)$. If real numbers u_t, $u'_{t'}$ satisfy $W_{tt'} \leq u_t + u'_{t'}$ for all $t \in X_0$ and $t' \in Y_0$, then $\tau(W) \leq \sum_t u_t + \sum_{t'} u'_{t'}$.*

Proof. Each Leibniz term is a product of entries along a perfect matching. So its valuation is at least minus the weight of the matching. The valuation of a sum is at least the minimum of the valuations of its terms. The second statement is weak LP duality for the assignment problem. The weight of a perfect matching is at most $\sum_t u_t + \sum_{t'} u'_{t'}$. □

Lemma 4.2 (rank of the reduction)**.** *Let (u, u') be integer potentials with $W_{tt'} \leq u_t + u'_{t'}$ for the matrix $\Phi_{r,I}$ at p, of value $\tau' = \sum_t u_t + \sum_{t'} u'_{t'}$. Then $\widetilde{\Phi} = \operatorname{diag}(p^u)\,\Phi_{r,I}\,\operatorname{diag}(p^{u'})$ has entries in $\mathcal{O}$, and*

$$e_p(r, I) \leq \tau' - \operatorname{corank}_{\mathbb{F}_p(x)}(\widetilde{\Phi} \bmod p).$$

Proof. We have $v(\det \Phi_{r,I}) = -\tau' + v(\det \widetilde{\Phi})$. Over the discrete valuation ring $\mathcal{O}$ the Smith normal form shows $v(\det \widetilde{\Phi}) \geq \operatorname{corank}(\widetilde{\Phi} \bmod p)$. □

4.3. **Second-order information.** The rank bounds below use a refinement of Lemma 4.2 at second order. Suppose that, after a unimodular change of rows and columns over $\mathcal{O}$, the first n_1 rows and the first n_2 columns of $\widetilde{\Phi}$ are divisible by p. Suppose also that the $n_1 \times n_2$ block where they meet is divisible by p^2. Dividing the first n_1 rows by p leaves the first n_2 columns divisible by p. So

$$v(\det \widetilde{\Phi}) \geq n_1 + n_2. \tag{4.1}$$

The vectors that produce such rows and columns are kernel vectors of the reduction of $\widetilde{\Phi}$ on single residue classes. The condition on the corner is a statement about their pairing at first order, which we now describe. Let $\widetilde{\Phi} = p^2\Phi_{r,I}$, so that all potentials equal 1. Since $q(r+2m) < p^2$, two distinct nodes $t \neq t'$ have $v_p(t'-t) \geq 1$ only if $t \equiv t' \pmod p$, and then $v_p(t'-t) = 1$. Hence $\widetilde{\Phi}$ has entries in $\mathcal{O}$. Its reduction is block diagonal over the residue classes modulo p, which we call the *fibres*. For nodes t, t' in distinct fibres,

$$\widetilde{\Phi}_{tt'} \equiv \frac{p\,(V(t') - V(t))}{t' - t} \pmod{p^2}. \tag{4.2}$$

Here $V(t) \in \mathbb{F}_p$ is the residue of p times the partial sum $\sum_{k \leq t}(\frac{1}{k} - \frac{q}{qk-u})$, up to the normalising factor q^{-2}. The function V is a step function of t which jumps only at the poles $t \equiv 0$ and $t \equiv k_p$. Since $t' - t$ is prime to p, only its residue matters in (4.2).

For a vector $\mathbf{y}$ indexed by nodes, its *moment pair* is

$$\hat{\mathbf{y}} = \Big(\sum_t y_t,\ \sum_t y_t V(t)\Big).$$

Lemma 4.3 (pairing)**.** *Let $\mathbf{y}$ be supported on the rows of one fibre and $\mathbf{y}'$ on the columns of a different fibre. Let t and t' be nodes of the first and of the second fibre. Then*

$$\mathbf{y}^\top \widetilde{\Phi}\, \mathbf{y}' \equiv \frac{p}{t' - t} \det \begin{pmatrix} \hat{\mathbf{y}} \\ \hat{\mathbf{y}}' \end{pmatrix} \pmod{p^2}.$$

In particular the pairing vanishes modulo p^2 whenever the moment pairs of $\mathbf{y}$ and $\mathbf{y}'$ are proportional.

Proof. Expand $\mathbf{y}^\top \widetilde{\Phi}\, \mathbf{y}'$ with (4.2). □

The moment pairs are never both zero. This follows from the displacement structure of Löwner matrices.

Lemma 4.4 (moment lemma)**.** *Let A be a matrix with rows indexed by a set X_0 of distinct integers and columns by a set Y_0, with $|X_0| = |Y_0| + 1$. Suppose that its off-diagonal entries are $A(t,t') = (\mathcal{L}(t') - \mathcal{L}(t))/(t' - t)$ for some function $\mathcal{L}$, and that its diagonal entries, on $X_0 \cap Y_0$, are arbitrary. Then A has a nonzero left kernel vector $\mathbf{y}$ with $(\sum_t y_t, \sum_t y_t \mathcal{L}(t)) \neq (0,0)$.*

Proof. Write $\operatorname{diag}(X_0)$ and $\operatorname{diag}(Y_0)$ for the diagonal matrices of the indices. Then $\operatorname{diag}(X_0)A - A\operatorname{diag}(Y_0) = \mathcal{L}_{X_0}\mathbf{1}^\top - \mathbf{1}\mathcal{L}_{Y_0}^\top$, because the diagonal entries cancel on common indices. The left kernel $\mathcal{V}$ of A is nonzero. Suppose that every $\mathbf{y} \in \mathcal{V}$ had zero moment pair. Multiplying on the left by $\mathbf{y}^\top$ would give $\mathbf{y}^\top \operatorname{diag}(X_0)A = 0$. So $\operatorname{diag}(X_0)$ would preserve $\mathcal{V}$. As this matrix is diagonal with distinct entries, $\mathcal{V}$ would contain a standard basis vector. That vector is supported on one index t, and its moment pair is $(1, \mathcal{L}(t)) \neq (0,0)$. This is a contradiction. □

4.4. **The local bounds.** We now list the local bounds. Each is a statement of the following form. Let $P_{r,I}$ be a member with $|I| \le K$, and let $p > p_0$ be a prime with $q(r+2m) < p^2$ that satisfies the stated conditions. Then

$$e_p(r, I) \le \tau(r, m, p, k_p) + \epsilon_\tau\,|I|. \tag{4.3}$$

Here τ is the expression in the table and $\epsilon_\tau \in \{0,1\}$ is given in its last column. The constant p_0 depends only on F. We write $(\cdot)_+ = \max(\cdot, 0)$, $\|m\|_p$ for the distance of m to $p\mathbb{Z}$, and $\bar{u} = (k_p + p - r)_+$.

Theorem 4.5 (local bounds)**.** *The following bounds of the form* (4.3) *hold.*

	bound τ	*conditions*	ϵ_τ
(T1)	$2(r+m)$	*none*	0
(T1′)	$2r + 2m - \|m\|_p$	*none*	0
(T2)	$2(r-\bar{u})_+ + 2m + \min(\bar{u}, 2m)$	$2m < p \le r$	0
(T3)	$2r + m$	$2m < p \le r$	0
(T4)	$2(r-k_p)_+ + 2m + (2m - (p-r))_+$	$r < p$	0
(T5)	$(2m - (k_p - r)_+)_+$	$\bar{N} \le p$, $r \le k_p$	0
(LP1)	$\tau_{\mathrm{T2}} - (\min(\bar{u}, 2m) - m)_+$	$p \le r < p + k_p$, $\bar{N} \le 2p$	1
(LP2)	$2(r-k_p) + 3m$	$r < p < r + m$, $k_p \le r$, $\bar{N} \le 2p$	1
(W)	$2r + 2m - (t_2 - t_1)$ *for each window* $[t_1, t_2)$ *of Theorem 4.7*	*as there*	0
(R)	Π_R *of* (4.4)	$2m < p$	0
(S)	Π_S *of* (4.5)	$2m \le p$	1
(U_n)	$\Pi_{U,n}$ *of* (4.6), $n \ge 0$	$np \le m < (n+1)p$, *and* $p \le 2m$ *if* $n = 0$	0

The rest of this subsection proves Theorem 4.5, family by family.

4.4.1. *The assignment bounds (T1)–(T5).* These follow from Lemma 4.1 with explicit potentials. Consider an off-diagonal entry $M[t, t']$. By the ultrametric inequality, $-v_p(M[t,t'])$ is at most the number of the following two events that occur: the interval $(t, t']$ contains a pole, and p divides $t' - t$.

A diagonal entry $M[t,t]$ has $-v_p \le 2$ if $t \ge k_p$ and $-v_p \le 0$ if $t < k_p$. The bound (T1) is the trivial bound, with potentials 1 on every row and column.

For (T1′) the potentials depend on the residue class modulo p. On the rows of a class they are $\frac{1}{2}$, $\frac{3}{2}$ or 1, according as X' has more, fewer or as many nodes in the class as Y'. On the columns they are $\frac{3}{2}$, $\frac{1}{2}$ or 1 in the same three cases. The border alternates between X' and Y'. So the imbalance is at most one per class, and the number of unbalanced classes is $2\|m\|_p$.

For (T2)–(T5) one starts from the potential $\mathbf{1}[t \ge k_p]$ on the common nodes t, which accounts for the diagonal. One adds 1, or $1+\mathbf{1}[t \ge p]$, on the border nodes t. Then one raises the potentials of the common nodes that have a congruent partner in the border. In each range one checks $W \le u+u'$ entrywise. Two facts are used. Two distinct nodes not congruent modulo p have weight at most 1. Two border nodes are never congruent once $2m < p$. For (T5) we have $\bar{N} \le p$ and $r \le k_p$. Then no common node accounts for the diagonal, and only the border nodes beyond k_p carry potential 1. So (T5) vanishes once $k_p \ge \bar{N}$, which happens for $p/r > \max(1+2\alpha, 1/\kappa)+o(1)$. All cases are checked in the formal proof.

4.4.2. *Primes near r: the bounds (LP1) and (LP2).* For p close to r the reduction of the rescaled matrix has a large block of rank one per residue class. Its corank can be computed. We give the proof of (LP2). The bound (LP1) applies to the primes with $p \le r < p + k_p$ and $\bar{N} \le 2p$. It is proved by the same method. With the potentials of (T2), eliminating an invertible block of common nodes brings the reduction to the form $\left(\begin{smallmatrix} \tilde{x}\mathrm{Id} & \mathcal{J}_Y \\ \mathcal{J}_X & 0 \end{smallmatrix}\right)$, where $\tilde{x}$ is the variable x plus a constant. In this form the border nodes below $p+k_p$ contribute rows and columns proportional to unit vectors. The Frobenius–König theorem bounds its rank. The saving over (T2) is $(\min(2m, p+k_p-r)-m)_+$, where $\min(2m, p+k_p-r)$ is the number of border nodes below $p+k_p$. We refer to the formal proof for the details.

Theorem 4.6. *Assume $r < p < r+m$, $k = k_p \le r$ and $\bar{N} = r+2m \le 2p$, and that every pole below $\bar{N}$ is simple. Then $v(\det \Phi_{r,\emptyset}) \ge -(2(r-k)+3m)$. The same holds for every member, with $\epsilon_\tau = 1$.*

Proof. Use the potentials $u(t) = \mathbf{1}[k \le t]$ for common nodes $t < r$ and $u(t) = 1 + \mathbf{1}[p \le t]$ for border nodes t. Their value is $\tau' = 2(r-k) + 2m + (\bar{N} - p)$. Since $\bar{N} \le 2p$, the poles below $\bar{N}$ are k, p, and $p+k$ if $p+k < \bar{N}$. Each is simple. The residues of p times the increment $\frac{1}{t} - \frac{q}{qt-u}$ at these poles are $-q/j$, 1 and $-q/(q+j)$.

Rows and columns beyond $p+k$ vanish. Let $t \in [p+k, \bar{N})$ be a border node, so that $u(t) = 2$. Let t' be a node of the other kind. If $t' < k$, then $u(t') = 0$. The partial sum over $(t', t]$ then has valuation at least -1, and $t - t' \in (p, 2p)$ is prime to p. Otherwise $u(t') \ge 1$ and $v_p(M[t', t]) \ge -2$. In both cases the rescaled entry is divisible by p.

The remaining entries. For nodes below $p+k$ the only poles are k and p. Put $C_0 = [0, k)$ and $C_1 = [k, r)$. Recall that $\omega = q^{-2}$ is the normalisation of

(2.1). The reduction has the following blocks. On $C_0 \times C_0$ it is the regular part plus x on the diagonal. On $C_0 \times C_1$ the entries are $-\omega q/(j(t-t'))$ with $t' \in C_0$ and $t \in C_1$. On $C_1 \times C_1$ it is the scalar matrix $\omega(q/j)^2\mathrm{Id}$. A border node $t \in [r,p)$ against $t' \in C_0$ gives $\omega q/(j(t-t'))$. A border node $t \in [p,p+k)$ against $t' \in C_0$ gives $\omega(1-q/j)$ if $t' = t-p$, and 0 otherwise. All other border entries vanish.

Rank. Delete the zero rows and columns found above. This does not change the rank. Eliminate the scalar block on C_1. The Schur complement on C_0 is $\tilde{x}\mathrm{Id}$ plus a constant multiple of the matrix with entries $\sum_{i=1}^{p-r} \frac{1}{(t+i)(t'+i)}$ for $t,t' \in C_0$. Here $\tilde{x}$ is x plus a constant, and the computation uses the reflection $H_{p-1-n} \equiv H_n \pmod p$ of harmonic numbers. The border nodes $t \in [r,p)$ are indexed by $i = p-t$. They are all present because $\bar{N} > p$. Even offsets are rows and odd offsets are columns, and together they cover $1 \le i \le p-r$. So row and column operations remove that multiple and leave $\left(\begin{smallmatrix} \tilde{x}\mathrm{Id} & \mathcal{J}_Y \\ \mathcal{J}_X & 0 \end{smallmatrix}\right)$. Here it is essential that x is an indeterminate, so that $\tilde{x}$ is a unit. A Schur complement on $\tilde{x}\mathrm{Id}$ then shows that the rank is $k + \operatorname{rank}(\mathcal{J}_X\mathcal{J}_Y)$. In $\mathcal{J}_X\mathcal{J}_Y$ the block between the high rows and the high columns is zero, since their offsets $t-p$ have opposite parities. So $\operatorname{rank}(\mathcal{J}_X\mathcal{J}_Y) \le p-r$. In total the rank of the reduction is at most $(r-k)+k+(p-r) = p$, and the corank is at least $r+m-p$. Lemma 4.2 gives $v(\det) \ge (r+m-p) - \tau' = -(2(r-k)+3m)$. Deleting a common node changes the potentials by at most one. □

Under the hypothesis $\bar{N} \le p+k$ the third pole does not occur, and the proof is shorter. The border nodes beyond $p+k$ only contribute rows and columns that vanish modulo p. This observation allows the whole range $\bar{N} \le 2p$, which is automatic on the interval $r < p < r+m$.

4.4.3. *Pole-free windows: the bound (W).* Like the rank bounds, the bound (W) does not come from the valuations of the entries alone. Between two consecutive poles the step function V of (4.2) is constant. The reductions of the fibres of nearby residue classes are then translates of one another. Kernel vectors can be transported from one fibre to the next. Their pairings vanish modulo p^2 by Lemma 4.3. The corner bound (4.1) turns this into a saving at second order.

A *cut* is an integer t with $t \equiv 0$, $t \equiv k_p$ or $t \equiv r \pmod p$. A window $[t_1,t_2)$ of integers is *cut-free* if no cut lies in the open interval (t_1,t_2).

Theorem 4.7. *Assume that every pole below $\bar{N}$ is simple and $\bar{N} < p^2$. Let $n \ge 0$ with $np \le m$, and put $\bar{N}_n = r + 2(m-np)$. Let $[t_1,t_2)$ be a cut-free window with $\max(r, \bar{N}_n - p) \le t_1 \le t_2 \le \min(\bar{N}_n, r+p)$. Then every member satisfies*

$$e_p(r,I) \le 2r + 2m - (t_2 - t_1).$$

For $n = 0$ the windows lie in the border above $\max(r, \bar{N}-p)$, and the bound applies to primes $p > m$. For $n = \lfloor m/p \rfloor \ge 1$ it applies to the small

primes $p \le m$. For these primes only (T1) and (T1′) are available among the assignment bounds. Read modulo p, the window range is then the arc of residues that receive an odd number of border nodes. The saving over (T1′) is the amount, if any, by which the length of the longest cut-free run of this arc exceeds $\|m\|_p$. For other n the window range is empty. The proof uses F only through its poles. For a function with several digamma terms the same statement holds when the cuts are the residues of all the poles and of r.

Proof. Let $\widetilde{\Phi} = p^2\Phi$, with independent diagonal variables on the common nodes. Every member is a coefficient of this determinant, so a bound for it bounds every member. Let $i < i'$ be two positions of the window. They lie in different residue classes. The translation $t \mapsto t + (i' - i)$ maps the nodes of the fibre of i bijectively onto those of the fibre of i'. It maps common nodes to common nodes, since no $t \equiv r$ lies in $(i, i']$. It maps border nodes to border nodes, since both classes lie in the window range. It maps X' to Y' when $i' - i$ is odd. There is no pole in $(i, i']$. So the step function V and the diagonal residues are unchanged, and so are the differences of nodes. Hence the reduced block of the fibre of i' equals that of i when $i' - i$ is even, and its transpose when $i' - i$ is odd.

In the window the fibres alternate between one more row than column and one more column than row. Take a nonzero left kernel vector $\mathbf{y}$ of one block and transport it to every fibre of the window. It is a left kernel vector on the fibres with a surplus row and a right kernel vector on the others. Its moment pair is the same everywhere. By Lemma 4.3 all pairings between these vectors vanish modulo p^2. There are n_1 such vectors on rows and n_2 on columns, with $n_1 + n_2 = t_2 - t_1$. Extending them to unimodular matrices, (4.1) gives $v(\det \widetilde{\Phi}) \ge t_2 - t_1$. Since $\det \widetilde{\Phi} = p^{2(r+m)} \det \Phi$, the theorem follows. □

4.4.4. *The rank bound (R).* Let $2m < p$. Write $r = k_0 p + r_p$ with $0 \le r_p < p$. Note that $k_p = \lceil jp/q \rceil$, since $0 < u < q$. Put $\mathrm{arc}_0 = [0, k_p - 1]$ and $\mathrm{arc}_1 = [k_p, p - 1]$. For $k \ge 0$ and $\iota \in \{0, 1\}$ let

$$n_{k,\iota} = \big|[r, r + 2m) \cap [kp + \min \mathrm{arc}_\iota,\ kp + \max \mathrm{arc}_\iota]\big|,$$
$$n_{\mathrm{sq}} = \big|[\max(0, r + 2m - p), \min(r, k_p))\big|,$$

and

$$\Pi_R = 2(r + m) - \max\big(m + n_{\mathrm{sq}},\ n_{0,0} + n_{1,0} + 2n_{\mathrm{sq}},\ n_{k_0,0},\ n_{k_0,1},\ n_{k_0+1,0},\ n_{k_0+1,1}\big). \tag{4.4}$$

Proof of the bound (R). Every node is $t = i + pk$ with residue $i \in [0, p)$ and digit k. Write $k(i)$ for the number of common nodes in the fibre of i. The border has length $2m < p$. So this fibre contains the common digits $0, \dots, k(i) - 1$ and at most one border digit, namely $k(i) \in \{k_0, k_0 + 1\}$. This border node is a row or a column according to the parity of $i + pk(i) - r$.

The step function V is constant on the fibre up to the bit $\iota(i) = \mathbf{1}[i \geq k_p]$. Thus the reduced block of a fibre depends only on $(k(i), \iota(i))$ and on the role of its border digit. Its entries have the Löwner form of Lemma 4.4.

A fibre with a border row has a left kernel vector with nonzero moment pair. A fibre with a border column has such a right kernel vector. For the digit 0 below the pole the step function vanishes. So these fibres have moment pairs proportional to $(1,0)$. A fibre of type $(1,0)$ without border digit has the zero block and contributes one row and one column. Choosing one such vector per fibre annihilates $m + n_{\mathrm{sq}}$ rows and as many columns. This gives the saving $m + n_{\mathrm{sq}}$.

Within a class of fibres of the same type (k, ι) all moment pairs are equal. So by Lemma 4.3 the pairings between the annihilated rows and columns of one class vanish modulo p^2. Then (4.1) gives a saving equal to the number of annihilated coordinates of the class. This number is $n_{k,\iota}$ for $k \geq 1$ and $(k,\iota) \neq (1,0)$. It is $n_{0,0} + n_{1,0} + 2n_{\mathrm{sq}}$ for the class with moment pair $(1,0)$. Taking the best class gives (4.4). A deleted common node changes at most one fibre. Give up that fibre, and one further fibre to restore the balance of rows and columns. The trivial certificate then costs at most two annihilated coordinates of each kind. The smaller matrix compensates for them, so $\epsilon_\tau = 0$. □

4.4.5. *Separated border nodes: the bound (S).* The bound (S) uses a different source of cancellation. Choose a window $[t_0, t_v]$ of border nodes whose residues lie in one arc between consecutive poles. Consider a border node outside it. Suppose the interval between the node and t_0 contains a singular index. Then the residues modulo p of p times the partial sums at the two nodes differ, because two singular contributions in an interval of length less than p cannot cancel. The leading coefficient in x of the bordered determinant of a fibre is this residue. So the cross product of the two bordered fibre determinants has a nonzero leading coefficient. The border node is then eliminated against t_0 with no further information. Count the residues of the window together with the residues of the eliminated border nodes. This gives an explicit bound, a minimum over the four window types $(k,\iota) \in \{k_0, k_0+1\} \times \{0,1\}$, which we now state.

Let $2m \leq p$, and let k_0, r_p, arc_0, arc_1 and $n_{k,\iota}$ be as in §4.4.4. For a type (k,ι) let $\mathcal{K}_{k,\iota}$ be the interval of the integers $kp + i$ with $i \in \mathrm{arc}_\iota$, and with $i < r_p$ if $k = k_0 + 1$. The border nodes in $\mathcal{K}_{k,\iota}$ form the window, and there are $n_{k,\iota}$ of them. Let n_K be the number of border nodes in $\mathcal{K}_{k,\iota}$ and in its two translates $\mathcal{K}_{k,\iota} \pm p$. Put

$$n_\emptyset = p - |\mathcal{K}_{k,\iota}| - (2m - n_K), \qquad n_+ = (n_{k,\iota} - m)_+.$$

Since $2m \leq p$, each residue class receives at most one border node. So n_K is the number of border nodes whose residues are those of $\mathcal{K}_{k,\iota}$. And $n_\emptyset$ is the number of the other residue classes that receive no border node. Then

$$\Pi_S = \min_{(k,\iota)} (2r + m - n_+ - (n_+ - n_\emptyset)_+), \tag{4.5}$$

where a type whose window is empty contributes $2r + m$. The bound (S) is $e_p(r, I) \le \Pi_S + |I|$. Its proof carries out the elimination above for every border node outside the window. We refer to the formal proof for it.

4.4.6. *Primes smaller than the border: the bound (U_n).* When $p \le 2m$ the border meets some residue classes more than once, and the bound (S) is not available. The bound (U_n) extends it to these primes. It works at third order in the Smith normal form. We show two things. After the elimination of a block of unit pivots, the rescaled matrix has the Smith invariants of a smaller geometry, to which the separation argument of (S) applies. The third-order information of that argument survives the elimination.

Write $m = np + \bar{m}$ with $0 \le \bar{m} < p$, so that $n = \lfloor m/p \rfloor$. Put

$$d = \|m\|_p = \min(\bar{m}, p - \bar{m}), \qquad \bar{r} = r + m - d.$$

For $i \in \{0, 1, 2\}$ and $\iota \in \{0, 1\}$ let $Z_i = p\lfloor r/p \rfloor + (n + i)p$, and put

$$w_0 = Z_i + \min \operatorname{arc}_\iota, \quad t_0 = \max(\bar{r}, w_0), \quad t_v = \min(Z_i + \max \operatorname{arc}_\iota, \bar{r} + p - 1).$$

We call the geometry with common block $[0, \bar{r})$ and border $[\bar{r}, \bar{r} + 2d)$ the *auxiliary geometry* $(\bar{r}, d)$. In it, let $n_+ = \big(\min(t_v + 1, \bar{r} + 2d) - t_0 - d\big)_+$. Let n_K be the number of border nodes in $[w_0, t_v]$ and in its two translates by $\pm p$, and let $n_\emptyset = p - (t_v + 1 - w_0) - (2d - n_K)$. Then

$$\Pi_{U,n} = \min_{i,\iota} \Big(2(r + m) - d - n_+ - (n_+ - n_\emptyset)_+ \Big), \tag{4.6}$$

all subtractions being truncated. This is the expression (4.5) for the auxiliary geometry $(\bar{r}, d)$, with the windows $[w_0, t_v]$ in place of $\mathcal{K}_{k,\iota}$. Note that $2(r + m) - d = 2\bar{r} + d$.

Theorem 4.8. *Let $p > p_0$ be a prime with $q(r + 2m) < p^2$, and let $n \ge 0$ with $np \le m < (n + 1)p$. Assume $p \le 2m$ if $n = 0$. Then every member satisfies $e_p(r, I) \le \Pi_{U,n}$.*

We first describe the proof for $n = 0$, that is $m < p \le 2m$. We then give the changes for $n \ge 1$. As in the proof of Theorem 4.7 we work with $\widetilde{\Phi} = p^2\Phi$. We put an independent indeterminate x_t on each common diagonal entry. We work over the discrete valuation ring $\mathcal{O}$ of Gauss valuations in these indeterminates. Every member is a coefficient of $\det \widetilde{\Phi}$, up to the deletion of common rows and columns, which we treat at the end.

Pivots from pairs of border nodes. For $n = 0$ there are $2m - p$ residue classes that carry two border nodes $t < t + p$, one a row and the other a column. There are $2d$ classes that carry one border node, and they lie in an interval of length $2d < p$. Eliminate the common block. Its fibres are invertible because the x_t are indeterminates. Consider a class with two border nodes. Let A be its common fibre, let $\mathbf{a}$ and $\mathbf{a}'$ be the couplings of its border column and border row to A, and let $\varkappa$ be its border entry. After the elimination the class has the pivot $\varpi = \varkappa - \mathbf{a}'^{\top} A^{-1} \mathbf{a}$. As the x_t tend to infinity, ϖ tends to a nonzero multiple of the residue of p times the

increment of F over $(t, t+p]$. This interval contains exactly one pole of each digamma term. Let $k'p$ be the multiple of p in it, and t' the pole of the second term. If the two contributions cancelled, then $q(t'-k'p) \equiv u$ with $|q(t'-k'p)-u| < q$. Hence $q(t'-k'p) = u$, which is impossible for $0 < u < q$. So these $2m-p$ pivots are units.

After them, the matrix of size $r+m = \bar{r}+d$ is equivalent over $\mathcal{O}$ to $\mathrm{diag}(\mathrm{Id}_{\bar{r}}, p\,\Xi)$, with Ξ of size d. Its Schur complement on the border has the expansion

$$S = p\,S_1 + p^2(S_2 - \mathcal{C}_{UB}\mathcal{B}^{-1}\mathcal{C}_{BU}) + O(p^3). \tag{4.7}$$

Here $\mathcal{B}$ is the diagonal of these pivots. The matrices $\mathcal{C}_{UB}$ and $\mathcal{C}_{BU}$ are the first-order couplings between these pivots and the single border nodes. The correction term in (4.7) is not negligible. The following identities account for it.

Lemma 4.9 (bordered inverses)**.** *Let A be an invertible square matrix over a commutative ring, and let $\mathbf{u}$ be a vector.*

(a) Let A' be obtained from A by replacing one column by a vector $\mathbf{w}$, and suppose that A' is invertible. Let $\mathbf{u}'$ be obtained from $\mathbf{u}$ by replacing the corresponding entry by $\mathbf{u}^\top A^{-1}\mathbf{w}$. Then $\mathbf{u}'^\top A'^{-1}\mathbf{u} = \mathbf{u}^\top A^{-1}\mathbf{u}$.

(b) Suppose that the matrix

$$A' = \begin{pmatrix} A & \mathbf{a} \\ \mathbf{a}'^\top & \varkappa \end{pmatrix}$$

is invertible. Then $(\mathbf{u}, y)^\top A'^{-1}(\mathbf{u}, \mathbf{a}'^\top A^{-1}\mathbf{u}) = \mathbf{u}^\top A^{-1}\mathbf{u}$ for every scalar y.

No symmetry of A or A' is assumed.

Proof. (a) Comparing columns gives $\mathbf{u}'^\top = \mathbf{u}^\top A^{-1}A'$. (b) We have $A'\,(A^{-1}\mathbf{u}, 0)^\top = (\mathbf{u}, \mathbf{a}'^\top A^{-1}\mathbf{u})^\top$. □

Moments. In the separation argument of (S), a border node is counted when its moment relative to a reference node vanishes. That moment is the bilinear form $\mathbf{u}^\top A^{-1}\mathbf{u}$ in the inverse of the common fibre A of its residue class. After the elimination, the common fibre of a class with two border nodes is replaced by a mixed block A' as in Lemma 4.9. Its lower node is a row and its upper node a column, or conversely. The product added to A'^{-1} is exactly the product subtracted in (4.7). So, by Lemma 4.9, the moments of the counted classes are unchanged. The single border nodes outside the window are separated from it as in (S), since two singular contributions in an interval of length $2d < p$ cannot cancel. Applied to the auxiliary geometry $(\bar{r}, d)$, the count of (S) then gives (4.6). It uses the third-order analogue of (4.1). Call the *coupling* of a residue class the residue modulo p of its border entry minus its moment against the reference node. Suppose that n_+ kernel vectors on each side survive, and that at most n_0 residue classes have a nonzero coupling. Then $\sum_i \min(v_i, 3) \geq d + n_+ + (n_+ - n_0)_+$, where

$v_1, v_2, \dots$ are the valuations of the Smith invariants. The first term d is the factor p of $p\,\Xi$.

Members. Deleting ℓ common nodes lowers the trivial bound $2(r+m)$ by 2ℓ. Consider a class with one deleted common node whose bordered system keeps its value at infinity ϑ (Lemma 4.10 below). Then the class keeps its moment. Indeed, let A be the common fibre and delete its last coordinate. The residual of the smaller solution is supported on that coordinate. Since the two values at infinity are equal, its pairing with the data vanishes. A class whose value at infinity changes gives a new separation instead. Classes with two or more deleted nodes, and classes outside the window, cost at most one each. Let k_1, k_2 and k' be the numbers of classes of these three kinds. Then $k_1 + 2k_2 + k' \le \ell$. An elementary inequality for the count $n_+ + (n_+ - n_0)_+$ shows that the loss is at most 2ℓ. Hence the bound holds for every member with $\epsilon_\tau = 0$.

Larger n. Let $np \le m < (n+1)p$. Each class of the remaining border absorbs n pairs of border nodes. Its block after the elimination is a mixed block. Its rows and columns are the nodes of $2n+1$ consecutive quotients $\beta, \beta+1, \dots, \beta+2n$. Here β is the number of common nodes in the class, and the quotient of a node t in the class of a residue i is $(t-i)/p$. The window counts are those of the first case at the scale $r+np$, translated by np. The new difficulty is to show that a remaining border node separated from the window by a pole has a nonzero coupling. The values at infinity are now those of $(n+1)\times(n+1)$ bordered systems, which we describe first.

Lemma 4.10 (bordered systems and rational interpolation)**.** *Let $\mathbb{K}$ be a field and $\mathcal{Z} = \{0, 1, \dots, 2n\}$, with $(2n)! \neq 0$ in $\mathbb{K}$. Let $\mathcal{L} : \mathcal{Z} \to \mathbb{K}$, and let $\mathcal{Z} = X \sqcup Y$ with $|X| = n+1$ and $|Y| = n$. Consider the bordered system*

$$\sum_{y \in Y} \frac{\mathcal{L}(y) - \mathcal{L}(x)}{y - x}\, z_y + \vartheta = \mathcal{L}(x) \qquad (x \in X)$$

in the unknowns z_y and ϑ, with data $\mathcal{L}$. Its solutions correspond bijectively to the pairs $(\mathcal{N}, \mathcal{D})$ of polynomials with $\mathcal{D}$ monic of degree n, $\deg \mathcal{N} \le n$ and $\mathcal{N}(t) = \mathcal{L}(t)\mathcal{D}(t)$ for $t \in \mathcal{Z}$. Under this correspondence ϑ is the coefficient of t^n in $\mathcal{N}$. In particular the system is nonsingular exactly when the interpolation problem of type (n, n) has a unique solution. Then $\vartheta = \lim_{t\to\infty} \mathcal{N}(t)/\mathcal{D}(t)$ depends only on $\mathcal{L}$ and not on the split. We call ϑ the value at infinity *of the system.*

Proof. Let $\mathcal{W}(t) = \prod_{y \in Y}(t - y)$, and let $\mathcal{W}_y = \mathcal{W}/((t-y)\mathcal{W}'(y))$ be the Lagrange polynomials of Y. Given a solution, put $\mathcal{D} = \mathcal{W} - \sum_y z_y \mathcal{W}'(y)\, \mathcal{W}_y$ and $\mathcal{N} = \vartheta\mathcal{W} - \sum_y z_y \mathcal{L}(y)\mathcal{W}'(y)\, \mathcal{W}_y$. Then $\mathcal{N}(y) = \mathcal{L}(y)\mathcal{D}(y)$ for $y \in Y$. Dividing $\mathcal{N}(x) - \mathcal{L}(x)\mathcal{D}(x)$ by $\mathcal{W}(x) \neq 0$ gives the equation of the row x. Conversely, Lagrange interpolation on Y recovers $z_y = -\mathcal{D}(y)/\mathcal{W}'(y)$ and ϑ from $(\mathcal{N}, \mathcal{D})$. □

Lemma 4.11 (harmonic data)**.** *Let $\mathcal{A} \in \mathbb{K}$ with $\mathcal{A} + i \neq 0$ for $0 \le i < 2n$, and let $\mathcal{L}_\mathcal{A}(t) = \sum_{0 \le i < t} 1/(\mathcal{A} + i)$. Let the characteristic of $\mathbb{K}$ be 0 or larger*

than $2n$. For every split, the bordered system of Lemma 4.10 with $\mathcal{L} = \mathcal{L}_{\mathcal{A}}$ is nonsingular. Its value at infinity is $\vartheta = \mathcal{L}_{\mathcal{A}}(n) + \mathsf{c}_n$, with $\mathsf{c}_n \in \mathbb{K}$ depending only on n. Consequently, for the harmonic data $\mathcal{L}(t) = H_t$ on the quotients $\beta, \dots, \beta + 2n$ the value at infinity is $H_{\beta+n} + \mathsf{c}_n$. The values at infinity of two classes with β and $\beta + 1$ common nodes differ by $1/(\beta + n + 1)$.

Proof. Write Δ for the forward difference. For $l \geq 1$ one has $\Delta^l \mathcal{L}_{\mathcal{A}}(k) = (-1)^{l-1}(l-1)!/(\mathcal{A}+k)_l$, where $(y)_l$ is the rising factorial. One also has $\Delta^l[\binom{t}{k} f(t)](0) = \binom{l}{k} \Delta^{l-k} f(k)$. By Lemma 4.10 a solution is a pair $(\mathcal{N}, \mathcal{D})$ with $\mathcal{N} = \mathcal{L}_{\mathcal{A}}\mathcal{D}$ on $\mathcal{Z}$. Since $\mathcal{N}$ has degree at most n, this means $\Delta^l(\mathcal{L}_{\mathcal{A}}\mathcal{D})(0) = 0$ for $n < l \leq 2n$. Write $\mathcal{D} = \sum_k \mathsf{d}_k \binom{t}{k}$ and insert the two formulas. After removing nonzero factors of the rows and columns, these n equations become a system for the coefficients d_k with $k < n$. Its matrix is the Cauchy matrix $\left(1/(l-k)\right)_{n<l\leq 2n,\, 0\leq k<n}$, and its right-hand side comes from $\mathsf{d}_n = 1$. The Cauchy matrix is nonsingular. So $\mathcal{D}$ is unique, which is the nonsingularity.

The same computation shows that $\mathsf{d}_k/(\mathcal{A}+k)_{n-k}$ does not depend on $\mathcal{A}$. For $k < n$, $\Delta^{n-k}\mathcal{L}_{\mathcal{A}}(k)$ is a constant multiple of $1/(\mathcal{A}+k)_{n-k}$. Hence $\vartheta = \Delta^n(\mathcal{L}_{\mathcal{A}}\mathcal{D})(0)/n! = \mathcal{L}_{\mathcal{A}}(n) + \mathsf{c}_n$ with c_n independent of $\mathcal{A}$. For the harmonic data on the quotients $\beta + t$ we have $H_{\beta+t} = H_\beta + \mathcal{L}_{\beta+1}(t)$. A constant added to $\mathcal{L}$ is added to ϑ. So the value at infinity is $H_\beta + \mathcal{L}_{\beta+1}(n) + \mathsf{c}_n = H_{\beta+n} + \mathsf{c}_n$. □

Remark 4.12. We note without proof that the interpolation problem of Lemma 4.11 can be solved in closed form. Its denominator is the polynomial $\mathcal{D}_n(t) = \sum_{k=0}^{n} \binom{n}{k}\binom{2n-k}{n-k}(\mathcal{A}+k)_{n-k}\, t(t-1)\cdots(t-k+1)$, of Hahn type [19]. The constant is $\mathsf{c}_n = 2H_n$. The determinant of the bordered system is $\mathrm{sf}(n)^2\,\mathrm{sf}(n-1)^2/(\prod_{x\in X, y\in Y}|x-y| \cdot \prod_{i<n}(\mathcal{A}+i)_{n+1})$, with $\mathrm{sf}(n) = \prod_{i\leq n} i!$. All its prime factors are at most $2n$ or divide some $\mathcal{A}+i$. These closed forms are not used in the proof and are not part of the formalization.

Separation for $n \geq 1$. Between a remaining border node and the reference node of the window lies a cut. Suppose first that the cut is a pole of the second digamma term. Deform the shift to $s + pY$, with an indeterminate Y. Evaluate at the value of Y at which that pole becomes central. After multiplication by the vanishing factor, the bordered systems of the two classes become block systems built from Cauchy matrices on the quotients. The square blocks are nonsingular by Cauchy's determinant formula. A rectangular Cauchy matrix with $k+1$ rows and k columns has a left kernel vector with nonzero coordinate sum. Otherwise the rational function $\sum_i y_i/(T_i - T)$, formed from a kernel vector (y_i) and the row nodes T_i, would vanish at the k column nodes while its numerator has degree less than k. This gives the two values at infinity the values 0 and the weight of the term, which differ.

Suppose next that the cut is a multiple of p. Take instead the leading coefficient in Y. Every fractional pole term tends to 0, and the data become the harmonic numbers. The two classes have β and $\beta + 1$ common nodes.

Lemma 4.11 gives values at infinity that differ by $1/(\beta+n+1)$. This is a unit because $\beta+2n+1<p$, by $q(r+2m)<p^2$.

In both cases the comparison is made on the cleared numerators of the original determinants, not on inverses. The result returns to $Y=0$ by Gauss's lemma. The cleared denominators are products of the elementary denominators of the entries, which are units at $Y=0$. So a lower bound for the Gauss valuation of the deformed determinant gives the same lower bound for every coefficient of the original one. With these separations, the count and the treatment of the members are those of the case $n=0$.

4.5. **Profiles.** Let $\mathcal{E}$ be the set of integer expressions in r,p,m,k built from integer constants with the following operations: addition, multiplication by integer constants, truncated subtraction $(X-Y)_+$, min, max, and the distance $\|X\|_p$ of an expression X to $p\mathbb{Z}$. The quantities $p\lfloor r/p\rfloor$ and $r \bmod p$ are also allowed. Every bound of Theorem 4.5 lies in $\mathcal{E}$. For (R) and (S) this is (4.4) and (4.5). The ends of the intervals $\mathcal{K}_{k,\iota}$ are $kp=p\lfloor r/p\rfloor+(k-k_0)p$ plus 0, k_p, k_p-1, $p-1$ or r_p-1. For (U_n) this is (4.6) with $d=\|m\|_p$ and $Z_i=p\lfloor r/p\rfloor+(n+i)p$. For (W) the window ends are the cuts nearest to the ends of the window range, which are expressions of the same kind.

The *homogenisation* $\hat\tau(\theta)$ of $\tau\in\mathcal{E}$ replaces r,p,m,k by $1,\theta,\alpha,\kappa\theta$. It replaces every constant by 0, $\|X\|_p$ by $\theta\|\hat X/\theta\|$ and $p\lfloor r/p\rfloor$ by $\theta\lfloor 1/\theta\rfloor$. It is piecewise affine with rational breakpoints, finitely many in every interval bounded away from 0.

Lemma 4.13 (homogenisation)**.** *For each $\tau\in\mathcal{E}$ there is an explicit constant $\mathrm{err}(\tau)$ such that $|\tau(r,p,m(r),k_p)-r\hat\tau(p/r)|\le\mathrm{err}(\tau)$ for all $r,p\ge1$. It is computed recursively from $\mathrm{err}(m)=\frac12$, $\mathrm{err}(k)=1-\frac1q$ and the rules $\mathrm{err}(\tau+\tau')=\mathrm{err}(\tau)+\mathrm{err}(\tau')$ and $\mathrm{err}(\min(\tau,\tau'))\le\max(\mathrm{err}(\tau),\mathrm{err}(\tau'))$.*

Proof. The real version of τ, with the constants dropped, is positively homogeneous of degree one. Every operation is 1-Lipschitz in the maximum norm. The inputs differ from their homogeneous values by $|m-\alpha r|\le\frac12$ and $|k_p-\kappa p|\le1-\frac1q$. Induct on τ. □

The conditions of Theorem 4.5 are linear inequalities in r,p,m,k, strict or not. They homogenise to intervals of θ. At an endpoint the error of such an inequality is a combination of $m-\alpha r$ and $k_p-\kappa p$. With the rounding $m=\lceil\alpha r-\frac12\rceil$ two endpoints are *exact*. These are $\theta=2\alpha$, from $2m<p$, and $\theta=(1+2\alpha)/(1+\kappa)$, from $\bar N\le p+k$. At an exact endpoint the condition holds for every prime on the correct side. The other endpoints are exact, or hold outside a strip of primes of bounded width around the endpoint. On that strip another bound must be used.

4.6. **Tables for Λ and the arithmetic estimate.** For the root class j let $\mathcal{P}_j(\theta)$ be the minimum of the homogenised bounds $\hat\tau(\theta)$ over the bounds τ of Theorem 4.5 valid at θ. A *table for* Λ, for the function F_s and the geometry

of parameter c, consists of the parameters (q, α, K) and the following data for every root class j:

- breakpoints $0 < \theta_0 < \theta_1 < \cdots < \theta_{\mathrm{end}}$;
- on each interval $(\theta_{i-1}, \theta_i]$, an affine function ℓ_i, a constant $\mathrm{err}_i \geq 0$ and a local bound τ_i of Theorem 4.5, with a second bound at each inexact endpoint of the range of τ_i;
- a constant ℓ_0 for the initial interval $(0, \theta_0]$;

together with a rational number Λ_{fin}. We call the table *admissible* if it satisfies the following conditions.

(C1) For each shift of the table, every unit class modulo the modulus q^* of the prime number theorem used is assigned a root class with a table.

(C2) On each piece ℓ_i is nonnegative, and the piece lies in the validity interval of its bound. Moreover $\hat{\tau}$ lies below ℓ_i at both ends and at both one-sided limits of each breakpoint inside the piece.

(C3) At each inexact endpoint the second bound is valid on both sides and lies below ℓ_i there. The constant err_i exceeds the error constants of Lemma 4.13, the numbers $\epsilon_\tau K$, the strip widths times the slopes, and the trivial bound at the scales below an explicit threshold.

(C4) $\ell_0 \geq 2 + 2\alpha$ and $0 < \theta_0 \leq 1$. Also $\theta_{\mathrm{end}} \geq \max(q(1+2\alpha), p_0/2)$ and $\kappa_j \theta_{\mathrm{end}} \geq 1 + 2\alpha$.

(C5) $\frac{1}{\varphi(q)} \sum_j \big(\ell_0\theta_0 + \sum_i \int_{\theta_{i-1}}^{\theta_i} \ell_i(\theta)\, d\theta\big) = \Lambda_{\mathrm{fin}}$.

Each condition is a finite set of comparisons of rational numbers.

Theorem 4.14 (denominators)**.** *Let $0 < c \leq 1$, and let F_s be the function with shift s of denominator q. Suppose that an admissible table for Λ of depth K has s among its shifts. Then for every even r and every member with $|I| \leq K$,*

$$\log \operatorname{den} P_{r,I} \leq (\Lambda_{\mathrm{fin}} + o(1))\, r^2.$$

Proof. By definition $\operatorname{den} P_{r,I} = \prod_p p^{e_p(r,I)}$. Fix a piece $(\theta_{i-1}, \theta_i]$ of the root class j. Let p be a prime in a unit class assigned to j, with $\theta_{i-1} r < p \leq \theta_i r$. Let r be beyond the threshold of (C3). Then the conditions of the bound τ_i hold, either exactly or outside a strip where the second bound applies. Lemma 4.13 with (C2) and (C3) gives $e_p(r, I) \leq r\, \ell_i(p/r) + \mathrm{err}_i$. Below the threshold the trivial bound is at most err_i.

For $p \leq \theta_0 r$ the bound (T1) gives $e_p \leq \ell_0 r + O(1)$. The primes with $p^2 \leq q(r + 2m)$, the prime powers and the finitely many primes below p_0 contribute $O(r^{3/2} \log r)$. For $p > \theta_{\mathrm{end}} r$, (C4) gives $\bar{N} \leq p$ and $k_p \geq \bar{N}$, and (T5) gives $e_p = 0$.

Fix a unit class modulo q^*. By the prime number theorem in arithmetic progressions,

$$\sum_{\substack{\theta_{i-1} r < p \leq \theta_i r \\ p \text{ in the class}}} r\, \ell_i(p/r) \log p = \frac{r^2}{\varphi(q^*)} \int_{\theta_{i-1}}^{\theta_i} \ell_i(\theta)\, d\theta + o(r^2).$$

The class j receives $\varphi(q^*)/\varphi(q)$ unit classes modulo q^*, because the map that sends a unit to qs times its inverse permutes the units modulo q. Summing and using (C5) gives the theorem. The constants err_i contribute $O(r)$. $\square$

The tables of this paper are generated from the functions $\mathcal{P}_j$ themselves. The pieces are the maximal intervals on which $\mathcal{P}_j$ is affine and attained by one bound. So $\Lambda_{\mathrm{fin}} = \frac{1}{\varphi(q)} \sum_j \int \mathcal{P}_j$ up to the initial interval. Every bound of Theorem 4.5 is available to the tables: (T1)–(T5), (LP1), (LP2), (R), (S), the windows (W) and the bounds (U_n). The windows with $n = 0$ are used for $m < p$, between consecutive cuts.

On $(0, \alpha]$, that is for $p \le m$, the only other bounds are (T1), (T1′) and (U_n). With (T1′) alone the profile there is the sawtooth $2 + 2\alpha - \theta\|\alpha/\theta\|$. Its integral is

$$\int_0^\alpha \Big(2 + 2\alpha - \theta\Big\|\frac{\alpha}{\theta}\Big\|\Big)\,d\theta = 2\alpha + \frac{\alpha^2}{2} + 2\alpha^2 \log 2,$$

since $\int_1^\infty \|y\| y^{-3}\,dy = \frac{3}{2} - 2\log 2$. The windows with $n = \lfloor m/p \rfloor \ge 1$ lower this profile on the strips $\alpha/(n+1) < \theta \le \alpha/n$. The tables use them for $n \le 8$, again with each window running between two consecutive cuts. For $q = 3, 4, 5, 7$ the tables also use (U_0) on $(\alpha, 2\alpha]$ and (U_n) on the strips with $1 \le n \le 24$. The table for $q = 2$ uses only the other bounds. Beyond the strips used, a table resolves the sawtooth on each interval $(\alpha/(n+1), \alpha/n]$ with $n < n_0$. It ends with a constant piece on $(0, \alpha/n_0]$, which costs $\alpha^2 T_{n_0}$ with $0 < T_n \le 1/(4n^2)$. The number Λ_{fin} is then rational, and (C5) is an identity between rational numbers. The bounds depend on the shift s only through the root class, and their profiles only through (q, j, α). So one table serves all shifts with denominator q.

5. Nonvanishing

The deduction needs, for every rational x_0 and every large scale, one member of the family that does not vanish at x_0. For the functions of (2.1) this holds in characteristic zero. The number of deleted common nodes is bounded independently of r. This is the statement used in Section 8.

5.1. **Selection in characteristic zero.** Normalise F so that the divided differences of F at consecutive nodes have partial sums f_t, with $f_{t+1} - f_t = \Delta_F(t)$. The diagonal partial sums $\tilde{f}_t$ satisfy $\tilde{f}_{t+1} - \tilde{f}_t = -\Delta_F'(t)$. For F_s,

$$\Delta_F(z) = \omega\Big(\frac{1}{z+1} - \frac{1}{z+s}\Big) = \frac{\omega(s-1)}{(z+1)(z+s)}.$$

For F_χ, with $\omega = -\frac{1}{6}$,

$$\Delta_F(z) = \frac{1 - z}{216\,(z+1)(z+\frac{1}{3})(z+\frac{1}{2})}.$$

Theorem 5.1 (selection)**.** *Let F be a function of (2.1), and consider the alternating geometry of parameter c. There is K, depending only on $|\mathcal{T}|$, with the following property. For every large even r and every $x_0 \in \mathbb{Q}$ some member $P_{r,I}$ with $|I| \le K$ satisfies $P_{r,I}(x_0) \neq 0$.*

Sketch of proof. By barycentric interpolation, a vector in the right kernel of $\Phi_{r,\emptyset}(x_0)$ corresponds to a pair of polynomials (u, w) of degree less than $r+m$. The pair satisfies the *value conditions* $\mathsf{w}(t) = f_t \mathsf{u}(t)$ at the $\bar{N}$ nodes. It satisfies the *jet conditions* $\mathsf{w}'(t) = f_t \mathsf{u}'(t) + (x_0 - \tilde{f}_t)\mathsf{u}(t)$ at the common nodes t. The left kernel is described in the same way. Two such pairs have zero cross product $\mathsf{w}_1\mathsf{u}_2 - \mathsf{w}_2\mathsf{u}_1$. Indeed its degree is at most $2(r+m)-2$, while the value and jet conditions give $\bar{N}+r = 2(r+m)$ zeros, counted with multiplicity.

Consider a pair whose degrees are small enough. Write $\Delta_F = n_\Delta/d_\Delta$ in lowest terms. The discrete Wronskian

$$\mathrm{Wr}(z) = d_\Delta(z)\big[\mathsf{w}(z+1)\mathsf{u}(z) - \mathsf{w}(z)\mathsf{u}(z+1)\big] - n_\Delta(z)\mathsf{u}(z)\mathsf{u}(z+1)$$

has more prescribed zeros at the nodes than its degree allows. So it vanishes identically. Then w/u would be a rational antidifference of Δ_F. This is impossible, because the simple pole of Δ_F at -1 is isolated in its class modulo $\mathbb{Z}$. The other poles of Δ_F are $-s$ for F_s, and $-\frac{1}{3}$ and $-\frac{1}{2}$ for F_χ. None of them is congruent to -1 modulo $\mathbb{Z}$. Clearing denominators in $\mathsf{w}(z+1)/\mathsf{u}(z+1) - \mathsf{w}(z)/\mathsf{u}(z) = \Delta_F(z)$ would propagate a root of the denominator indefinitely in one direction. Counting the zeros along chains of nodes of step 2 bounds the dimension of the kernel. So the corank of $\Phi_{r,\emptyset}(x_0)$ is at most a number K that depends only on $|\mathcal{T}|$ and on the number of such chains. That number is bounded independently of r.

Finally, the left and right kernels map to the same interpolation space. At a common node t their coordinates are the values $\mathsf{u}(t)$ times nonzero factors. No nonzero kernel pair has u vanishing at every common node. Otherwise, dividing by the node polynomial of the common nodes would give a pair of degree less than m that satisfies the value conditions at all $\bar{N}$ nodes, and the Wronskian argument applies again. Hence there is a set I of $|I| =$ corank common nodes such that no nonzero left or right kernel vector vanishes on I. The rows outside I then span the row space, and the columns outside I span the column space. So the principal minor that deletes I is nonzero. □

The theorem applies to every function of (2.1). For F_χ the pole of Δ_F at -1 is again isolated in its class modulo $\mathbb{Z}$, since the other poles are $-\frac{1}{3}$ and $-\frac{1}{2}$. In the formal proof the depth is $K \le 64$ for all the functions of this paper.

6. Analytic decay

In this section we bound $|P_{r,I}(x)|$ for x near ξ. The argument is that of OpenAI [31] for square determinants, with three changes. The determinant has two families of nodes, one for the rows and one for the columns. The

comparison measure is a density on finitely many segments rather than an explicit equilibrium measure. The inequalities that remain are verified on a finite cover by cells. The result is one theorem for every geometry parameter $c \le 1$ and every function of (2.1).

6.1. **A contour representation.** Hermite's formula gives $\mathsf{L}_F[t,t'] = \frac{1}{2\pi i}\oint F(z)\upsilon_t(z)\upsilon_{t'}(z)\,dz$ with $\upsilon_t(z) = 1/(z-t)$. The contour encloses the nodes and no pole of F. Write $F = \sum_{\varsigma\in\mathcal{T}} w_\varsigma \psi(z+\varsigma)$ as in §3. The reflection formula $\psi(w) = \psi(1-w) - \pi\cot\pi w$ splits $F = \tilde{F} + E$. Here $\tilde{F}$ is analytic in the left half-plane and $O(1/|z|)$ there, and

$$E(z) = -\pi \sum_{\varsigma\in\mathcal{T}} w_\varsigma \cot\pi(z+\varsigma).$$

The function E is 1-periodic. Since the weights sum to zero, $E(z) = O(e^{-2\pi|\operatorname{Im} z|})$ for $|\operatorname{Im} z| \ge 1$. For F_s one has $E(z) = \pi[\cot\pi(z+s) - \cot\pi z]$.

Let Γ be a polygonal path, symmetric under complex conjugation. It crosses the real axis at a point $x^* < 0$ that is not a pole of F, runs through the left half-plane and ends in two rays. The poles of F are the points $-(i+\varsigma)$ with $i \ge 0$ and $\varsigma \in \mathcal{T}$. Moving the contour to Γ picks up the residues of F at the poles in $(x^*, 0)$. The part of the integrand with $\tilde{F}$ integrates to 0 over Γ. Hence, for all integers $t, t' \ge 0$,

$$\mathsf{L}_F[t,t'] = \int \upsilon_t \upsilon_{t'}\, d\Omega. \tag{6.1}$$

Here Ω is the complex measure with the mass $-\operatorname{res}_y(F)$ at each pole $y \in (x^*, 0)$ of F and the density $(2\pi i)^{-1}E(z)\,dz$ on Γ. This is a Gram representation with respect to a measure with finitely many atoms and an exponentially small density on Γ. After scaling by r, the atoms lie on a segment $[-r_0, 0]$ with density $|\mathcal{T}|$. The separation condition on x^* is a finite check on the fractional parts of $\varsigma - x^*$ for $\varsigma \in \mathcal{T}$.

6.2. **Two-family Andréief bound.** For a set $C'' \subseteq C$ of common nodes put $\mathcal{M}_{C''} = \omega\,\mathsf{L}_F[C'' \sqcup X', C'' \sqcup Y']$. Consider a member with common nodes $C' = C \setminus I$. Put $R' = C' \sqcup X'$, $Q' = C' \sqcup Y'$ and $n_* = |C'| + m$. By multilinearity in the diagonal,

$$P_{r,I}(\xi + h) = \sum_{C''\subseteq C'} h^{|C'|-|C''|} \det \mathcal{M}_{C''}. \tag{6.2}$$

Andréief's identity [14] for the two families $(\upsilon_t)_{t\in R'}$ and $(\upsilon_{t'})_{t'\in Q'}$ gives

$$\det \mathsf{L}_F[R', Q'] = \frac{1}{n_*!}\int \det[\upsilon_t(z_i)]\,\det[\upsilon_{t'}(z_i)] \prod_{i=1}^{n_*} d\Omega(z_i).$$

Cauchy's determinant formula evaluates both factors. We take absolute values. The logarithmic potential theory behind what follows is in [37]. We

obtain

$$|\det \mathsf{L}_F[R',Q']| \le \frac{1}{n_*!}\int \prod_{i<i'}|z_i-z_{i'}|^2 \prod_{\mathcal{F}\in\{R',Q'\}} \frac{\prod_{t<t'\in\mathcal{F}}|t'-t|}{\prod_{i,\,t\in\mathcal{F}}|z_i-t|}\ \prod_i d|\Omega|(z_i).$$

The integrand is the exponential of minus a discrete logarithmic energy. The free points z_i repel each other with weight 1. Each node family repels itself with weight $\frac{1}{2}$ and attracts the free points with weight 1. The density $|E| \approx e^{-2\pi|\operatorname{Im} z|}$ acts as an external field. The rows and columns differ only in the interleaved border. So the two node families have the same macroscopic density. After scaling by r, both are

$$\nu_c = \mathrm{Leb}|_{[0,1]} + \tfrac{1}{2}\mathrm{Leb}|_{[1,1+c]},$$

of mass $|\nu_c| = 1 + c/2$.

6.3. **The energy certificate.** For a regularisation radius $\epsilon > 0$ let $k_\epsilon(z,w) = -\frac{1}{2}\log(|z-w|^2+\epsilon^2)$. Then $k_\epsilon(z,w) \le -\log|z-w|$.

Lemma 6.1 (conditional positive definiteness)**.** *Let $z_i \in \mathbb{C}$ be finitely many points and y_i real numbers with $\sum_i y_i = 0$. Then $\sum_{i,j} y_i y_j k_\epsilon(z_i,z_j) \ge 0$.*

Proof. We use Frullani's integral $\log T = \int_0^\infty (e^{-u} - e^{-Tu})\,du/u$ for $T > 0$. Together with $\sum_i y_i = 0$ it gives $\sum_{i,j} y_i y_j(-\log(|z_i-z_j|^2+\epsilon^2)) = \int_0^\infty \frac{e^{-u\epsilon^2}}{u}\sum_{i,j} y_i y_j e^{-u|z_i-z_j|^2}du$. Gaussian kernels are positive definite. Indeed, expanding $e^{2u\langle z,z'\rangle}$ in powers and using the binomial theorem writes the inner sum as a convergent sum of squares. □

A *trial measure* is a probability measure σ, symmetric under conjugation, carried by finitely many segments $\mathcal{S}_i \subset \Gamma \cup [-r_0,0] \cup [0,1]$ with rational endpoints. On each segment it has a density $\sigma_i(y)\,dy$ in the parameter $y \in [0,1]$ of the segment. Here σ_i is a polynomial with rational coefficients, and $0 \le \sigma_i \le \bar\sigma_i$ for a rational $\bar\sigma_i$. We write $\Theta_\sigma = \sum_i \bar\sigma_i/|\mathcal{S}_i|$ for the sum of the bounds of its densities with respect to arclength.

We compare the free points and the deleted nodes with $|\nu_c|\sigma$ through Lemma 6.1. This bounds the free–free energy from below by a sum of one-point terms and the constant $|\nu_c|^2\mathcal{I}(\sigma)$. Riemann-sum comparisons replace the node sums by the potential U^{ν_c} and the node self-energy by $\mathcal{I}(\nu_c)$, up to $O(r\log r)$. With the external field one arrives at the function

$$\Upsilon(z) = |\nu_c|\,U^\sigma(z) - U^{\nu_c}(z) + (1-\varepsilon_0)\pi|\operatorname{Im} z|$$

for a small fixed $\varepsilon_0 > 0$. Here $U^\sigma(z) = \int \log\frac{1}{|z-w|}\,d\sigma(w)$ is the logarithmic potential of σ, and U^{ν_c} is that of ν_c. A *certificate* consists of σ and rational numbers Υ_Γ, Υ_0, J, $\mathcal{I}_\sigma$, $\mathcal{I}_\nu$ with the following properties.

(a) $\Upsilon \ge \Upsilon_\Gamma$ on Γ.

(b) $\Upsilon \geq \Upsilon_0$ on $[0,1]$, the scaled position of the deleted common nodes.
(c) $J \leq \int_{-r_0}^{0} \min(\Upsilon(x) - \Upsilon_\Gamma, 0)\, dx$.
(d) $\mathcal{I}(\sigma) \leq \mathcal{I}_\sigma$ and $\mathcal{I}(\nu_c) \geq \mathcal{I}_\nu$.

Theorem 6.2 (decay on a neighbourhood)**.** *Let $0 < c \leq 1$. Let F be a function of (2.1) whose poles are separated from x^*. Let $(\sigma, \Upsilon_\Gamma, \Upsilon_0, J, \mathcal{I}_\sigma, \mathcal{I}_\nu)$ be a certificate for Γ and ν_c. Let $\epsilon > 0$, $\delta_\epsilon \geq \pi\epsilon\,\Theta_\sigma$, $\delta_{\mathrm{m}} > 0$ and $\delta_J > 0$ be rational, and put*

$$\delta = 3|\nu_c|^2\delta_\epsilon + 4|\nu_c|^2\delta_{\mathrm{m}} + 2|\mathcal{T}|\,\delta_J.$$

Let D and ρ be rational numbers with

$$D \leq 2(|\nu_c|\Upsilon_\Gamma + |\mathcal{T}|J) - |\nu_c|^2\mathcal{I}_\sigma + \mathcal{I}_\nu - \delta, \qquad \rho \geq 2(\Upsilon_\Gamma - \Upsilon_0).$$

Then for every depth K, every member of the alternating geometry of parameter c and every real h with $|h| \leq e^{-\rho r}$,

$$|P_{r,I}(\xi + h)| \leq \exp\bigl(-(D - o(1))r^2\bigr).$$

The loss δ comes from replacing σ by finitely many atoms in the comparison with the free points. The regularisation radius ϵ of Lemma 6.1 and the mesh of the atoms lower Υ_Γ by $|\nu_c|(\delta_\epsilon + \delta_{\mathrm{m}})$. They raise $\mathcal{I}_\sigma$ by $\delta_\epsilon + 2\delta_{\mathrm{m}}$. Darboux sums for the integral in (c) lower J by δ_J. Inserted in the expression for D, these changes give δ. The tolerances δ_{m} and δ_J may be any positive rational numbers, since the atoms can be taken as fine as needed. In the certificates of this paper $\epsilon = 10^{-12}$ and $\delta < 10^{-9}$.

The node energy $\mathcal{I}(\nu_c)$ is a finite rational combination of the values $\mathcal{H}(y_k)$, where $\mathcal{H}(y) = \frac{1}{2}y^2 \log|y| - \frac{3}{4}y^2$ and the y_k are the rational differences of the endpoints of the two pieces of ν_c. The lower bound $\mathcal{I}_\nu$ is checked with rational enclosures of the logarithms. These come from the series of artanh and a reduction to $[1,2)$. The energy $\mathcal{I}(\sigma)$ is a finite sum of pair energies of polynomial densities on segments. Each is a closed form of the type polynomial times logarithm plus arctangent, and is bounded in the same way. The conditions (a)–(c) are statements about Υ on sets of positive measure. They are proved on a cover by cells.

6.4. **The cover by cells.** Each of the conditions (a)–(c) is checked on a finite cover by cells with exact rational arithmetic. On a cell, the potential U^σ is split into the contribution of the segments of σ near the cell and the rest. A near segment is bounded by a Taylor polynomial in the cell variable with an explicit remainder. For a cell that meets the line of a segment beyond its end, the same bound is used for the segment with its orientation reversed. At a corner of Γ the two incident segments are treated together. Far segments are bounded by a multipole expansion with an explicit tail. The potential U^{ν_c} is an explicit elementary function, and it is bounded above by monotonicity on the cell.

For (a) the cells are intervals of the arclength on each edge of Γ. The bound is proved on a horizontal neighbourhood of the edge of fixed width. Beyond the last vertex the ray is handled by an analytic tail bound. For (b)

the cells are intervals of $[0,1]$. For (c) the integrand is bounded above on each cell of $[-r_0,0]$ by a constant or by a chord. Then J is the sum of the integrals of these majorants.

Proposition 6.3 (certified constants)**.** *For each row of Table 2 there is a trial measure σ on a polygon Γ, with a cover by cells satisfying the checks above, for which Theorem 6.2 gives the constants D and ρ of the table.*

All inequalities are verified by the Lean kernel, as described in Section 10. The certificate data are the polygon, the density σ and the cells. They were computed with floating-point optimisation and interval arithmetic in Arb [18]. Lean checks them with rational arithmetic only. The data play no role in the proof beyond satisfying these checks.

Remark 6.4 (how good a certificate can be). For an exponent μ_0 the certificate meets μ_0 exactly when $\mu_0(D-\Lambda)-\rho\ge 0$. This quantity is a concave function of σ. Its maximum over all probability measures on $\Gamma\cup[-r_0,0]\cup[0,1]$ is attained at a measure σ^*. For this measure $|\nu_c|\sigma^*$ is the weighted equilibrium measure of the constrained energy problem on this set, with the external field $(1-\varepsilon_0)\pi|\operatorname{Im} z|-U^{\nu_c}$ and the upper constraint $|\mathcal{T}|\,dx$ on the segment [37]. Weak duality expresses the deficit of any certificate as a sum of nonnegative terms. The first is the energy of $2\mu_0|\nu_c|(\sigma-\sigma^*)$. This is the difference between $2\mu_0|\nu_c|\sigma$ and the optimal dual measure $2\mu_0|\nu_c|\sigma^*$. So a certificate close to the best possible one must be close to σ^* in energy. Then $\Upsilon-\Upsilon_\Gamma$ is a small ripple on the support of σ^*. This is why a pointwise verification on a cover is needed. This remark is not used in the proof of Theorem 1.1 and is not part of the formalization.

6.5. **Node removal and the neighbourhood.** The argument applies to every subset $C''\subseteq C'$ of the common nodes. Removing common nodes turns them into points of the configuration that are not integrated over. Only common nodes can be removed. A node removed from one family only would leave an uncancelled term of weight $\frac12$. By condition (b) one obtains

$$|\det \mathcal{M}_{C''}|\le \exp\bigl(-(D-o(1))r^2+\rho r\,|C'\setminus C''|\bigr),$$

uniformly in C''. Insert this into (6.2) with $|h|e^{\rho r}\le 1$. The sum has at most $2^{|C'|}=e^{O(r)}$ terms, each at most $e^{-(D-o(1))r^2}$. This gives Theorem 6.2. The deletion of at most K common nodes in a member changes the energy by $O(r)$.

7. The pair of shifts $(\frac13,\frac23)$

For $\Psi(\frac13,\frac23)$ we use the function F_χ of (2.1), which combines three digamma terms, with the alternating geometry of parameter $c=\frac12$ at all even scales r. Then $\alpha=\frac14$ and $m=\lceil r/4-\frac12\rceil$. The decay and nonvanishing estimates of Theorems 6.2 and 5.1 apply unchanged. The denominators need more care. At a prime p the increments $g(k)$ of §3 have poles in three

residue classes instead of two, so the local bounds depend on two first poles. This section describes the two poles, the bounds we use, and the resulting constant.

7.1. **The first poles.** Let $p > 3$ be a prime. The increment $g(k) = \frac{1}{3k-2} - \frac{1}{2k-1} + \frac{1}{6k}$ has a pole modulo p exactly when p divides k, $3k-2$ or $2k-1$. Let k_A and k_B be the least $k \geq 1$ with $p \mid 3k-2$ and with $p \mid 2k-1$. Then

$$k_B = \frac{p+1}{2}, \qquad k_A = \begin{cases} \frac{p+2}{3}, & p \equiv 1 \pmod 6, \\ \frac{2p+2}{3}, & p \equiv 5 \pmod 6. \end{cases}$$

The residues of $p\,g(k)$ at k_A and k_B are 1 and -1 if $p \equiv 1 \pmod 6$, and $\frac{1}{2}$ and -1 if $p \equiv 5 \pmod 6$. In the first case $k_A < k_B$ and the two residues cancel. So the partial sums $\sum_{k \leq t} g(k)$ are p-integral for $t < k_A$, and again for $k_B \leq t < p$.

Both first poles are affine in one index. Define k_p by $6k_p = jp + 5$ with $0 < j < 6$. Then $j = 1$ for $p \equiv 1 \pmod 6$ and $j = 5$ for $p \equiv 5 \pmod 6$, $k_p = \frac{j}{6}p + O(1)$, and

$$(k_A, k_B) = \begin{cases} (2k_p - 1,\ 3k_p - 2), & p \equiv 1 \pmod 6, \\ (2k_p - p - 1,\ 3k_p - 2p - 2), & p \equiv 5 \pmod 6. \end{cases}$$

Thus j plays the role of the root class of §4, with q replaced by 6.

7.2. **The local bounds.** Fix a prime $p > 6$ with $6\bar{N} < p^2$. Every pole below $\bar{N}$ is then simple. Call a node t *regular* if $t < \min(k_A, k_B)$, or if $p \equiv 1 \pmod 6$ and t is a border node with $k_B \leq t < p$. At a regular node the partial sum of the increments up to t is p-integral. When t is a common node, its diagonal entry is p-integral too.

Proposition 7.1. *Every member of the alternating geometry of parameter $\frac{1}{2}$ for F_χ satisfies $e_p(r, I) \leq \tau$ for each of the following τ.*

(i) The bounds (T1) and (T1′), namely $2(r+m)$ and $2r + 2m - \|m\|_p$.

(ii) If $\bar{N} \leq 2p$, the bound $2r + 2m - n_{\mathrm{reg}}$. Here n_{reg} is the number of pairs formed by a regular node t and a side on which t lies, row or column, such that $t + p$ is not a node of the other side.

(iii) If $\bar{N} < \min(k_A, k_B)$, the bound 0.

(iv) The bound $2r + 2m - (t_2 - t_1)$ of Theorem 4.7, for every n and every window $[t_1, t_2)$ as there, with the cuts $t \equiv 0$, $t \equiv k_A$, $t \equiv k_B$ and $t \equiv r \pmod p$.

Proof. The bounds (i) are proved as in §4.4.1. For (ii) we apply Lemma 4.1 with the following potentials. At a node that is not regular, the potential is 1 on its row and on its column. At a regular node t, the potential on its row is 1 if $t + p$ is a column node, and 0 otherwise. The potential on its column is 1 if $t + p$ is a row node, and 0 otherwise. Their sum is $2(r+m) - n_{\mathrm{reg}}$, and they dominate the weights W. Indeed, a diagonal weight is 0 at a regular

common node and at most 2 in general. Two nodes that are not congruent modulo p have weight at most 1, since all poles are simple. Their weight is 0 if both are regular. Two distinct congruent nodes have weight at most 2. Since $\bar{N} \le 2p$ and regular nodes are smaller than p, at most one of them is regular. Suppose that the row node t of such an entry is regular. Then its column node is $t+p$, so the potential on the row of t is 1. The case of a regular column node is symmetric.

For (iii), the condition $\bar{N} < \min(k_A, k_B)$ implies $\bar{N} < p$. Then no index below $\bar{N}$ is a pole and no two nodes are congruent modulo p, so every entry is p-integral. For (iv), the proof of Theorem 4.7 uses F only through its poles. The step function V and the diagonal residues are constant between consecutive cuts. This is what the translation between the fibres of the window needs. □

The bounds (T2)–(T5), (LP1), (LP2), (R) and (S) of Theorem 4.5 are not used for F_χ.

7.3. **The denominators.** The tables for Λ of §4.6 and Theorem 4.14 apply to F_χ with q replaced by 6. The root class is $j \in \{1,5\}$ as above, and the bounds are those of Proposition 7.1. These bounds are expressions of the class $\mathcal{E}$ in r, p, m, k_A, k_B. On each class of primes modulo 6 the substitution above makes them expressions in r, p, m, k_p. Their homogenisations are taken with $\kappa = j/6$. The proof of Theorem 4.14 uses F only through these bounds and the root class. It goes through with the bound (iii) in place of (T5) beyond the end of the table.

The table of the formal proof starts at $\theta_0 = \frac{1}{68} = \alpha/17$. Below it the trivial bound $\ell_0 r$ with $\ell_0 = \frac{5}{2} = 2 + 2\alpha$ is used. On $(\alpha/17, \alpha]$ the table uses the bounds (i) and the windows (iv) with $1 \le n \le 16$. Beyond α it uses (i), (ii), the windows (iv) with $n = 0$ between consecutive cuts, and the bound (iii), which ends the table. The table is admissible for the two classes $j = 1, 5$. The arithmetic estimate therefore holds for every even r and every member with $|I| \le 64$, with the denominator exponent

$$\Lambda_\chi = 4.09592776731\ldots$$

This is an explicit rational number, whose numerator and denominator have 117 and 116 digits.

7.4. **Decay and nonvanishing.** The decay estimate is Theorem 6.2 for the function F_χ, with $|\mathcal{T}| = 3$ and $c = \frac{1}{2}$. The node measure $\nu_{1/2}$ has mass $\frac{5}{4}$ and energy

$$\mathcal{I}(\nu_{1/2}) = \frac{75}{32} - \frac{9}{8}\log\frac{3}{2} - \frac{1}{16}\log 2.$$

A trial measure on a polygon, with a cover by cells satisfying the checks of §6.4, gives

$$D = \frac{4643511791526209475773}{10^{21}} = 4.64351\ldots,$$
$$\rho = \frac{37521870783566737}{5\cdot 10^{15}} = 7.50437\ldots$$

Nonvanishing is Theorem 5.1 for F_χ. Its increment Δ_F has its pole at -1 isolated in its class modulo $\mathbb{Z}$, and the depth is $K \le 64$. The deduction of Section 8, at the even scales, gives

$$\mu(\Psi(\tfrac{1}{3},\tfrac{2}{3})) \le \frac{\rho}{D-\Lambda_\chi} = 13.704516\ldots < 13.7046,$$

since $\Psi(\frac{1}{3},\frac{2}{3}) = 36\,\xi_\chi$.

8. Proof of Theorem 1.1

The three estimates combine as follows.

Theorem 8.1 (deduction)**.** *Let F be a function of (2.1) with its geometry. Let K be a depth such that, for every large r and every rational x_0, some member $P_{r,I}$ with $|I| \le K$ satisfies $P_{r,I}(x_0) \neq 0$, as in Theorem 5.1. Suppose that the denominator exponent Λ of Theorem 4.14 and the constants D, ρ of Theorem 6.2 satisfy $\Lambda < D$. Then $\mu(\xi) \le \rho/(D-\Lambda)$.*

Proof. Let $\eta = D - \Lambda$, and let $\mu_0 > \mu_1 > \rho/\eta$. Suppose that $|\xi - a/b| < b^{-\mu_0}$ with b large. Let r be the least admissible scale with $r \ge \mu_1 \log b/\rho$. By Theorem 5.1 with $x_0 = a/b$ there is a member with $P_{r,I}(a/b) \neq 0$. By Lemma 3.4, $|\operatorname{den}(P_{r,I})\,P_{r,I}(a/b)| \ge b^{-r}$. Write $a/b = \xi + h$. Then $|h| < b^{-\mu_0} \le e^{-\rho r}$, because $\rho r \le \mu_1 \log b + O(1) < \mu_0 \log b$ for large b. By Theorems 4.14 and 6.2,

$$|\operatorname{den}(P_{r,I})\,P_{r,I}(a/b)| \le e^{(\Lambda+o(1))r^2} e^{-(D-o(1))r^2} = e^{-(\eta-o(1))r^2}.$$

Comparing the two bounds gives $(\eta - o(1))r^2 \le r\log b$. But $\eta r \ge (\mu_1\eta/\rho)\log b$ with $\mu_1\eta/\rho > 1$. This is a contradiction for large b. □

Proof of Theorem 1.1. For each row of Table 1 the function, the geometry parameter and the constants are those of Table 2. The denominator exponent is given by Theorem 4.14 with the table for Λ of the formal proof. For the pair $(\frac{1}{3},\frac{2}{3})$ it is given by Section 7. The decay rates are given by Proposition 6.3, and nonvanishing by Theorem 5.1. Theorem 8.1, applied at the even scales, gives $\mu(\xi) \le \rho/(D-\Lambda)$, and in every row $\rho/(D-\Lambda) < \mu_{s,t}$. By (2.1), ξ is a nonzero rational multiple of $\Psi(s,t)$, so $\mu(\Psi(s,t)) < \mu_{s,t}$. Since this inequality is strict, the definition of μ gives $|\Psi(s,t) - a/b| > b^{-\mu_{s,t}}$ for all integers a and all sufficiently large b. This proves Theorem 1.1. □

Remark 8.2 (relation to linear independence criteria). The deduction is a one-dimensional version of the familiar scheme behind the criteria of Nesterenko [29] and their use by Ball and Rivoal [2]. The nonzero integer

(s,t)	c	Λ	D	ρ	ρ/η	μ
$(\frac{1}{2},1)$	$\frac{3}{10}$	2.9923824	3.9785713	6.4350874	6.525207	6.5253
$(\frac{k}{3},1)$	$\frac{3}{10}$	3.3104458	4.0291249	6.8272548	9.499725	9.4998
$(\frac{1}{3},\frac{2}{3})$	$\frac{1}{2}$	4.0959278	4.6435117	7.5043742	13.70452	13.7046
$(\frac{k}{4},1)$	$\frac{1}{5}$	3.5149036	3.7447213	7.1904273	31.28751	31.2876
$(\frac{k}{5},1)$	$\frac{2}{9}$	3.5128488	3.8135797	7.1655663	23.82717	23.8272
$(\frac{k}{7},1)$	$\frac{1}{6}$	3.6391158643	3.6424792669	7.4259546075	2207.870	2207.8697

TABLE 2. The data of the proof of Theorem 1.1, with k prime to the denominator. For each pair of shifts the table gives the geometry parameter c, the denominator exponent Λ, the decay rate D, the neighbourhood exponent ρ and the exponent. The constants are the rational numbers of the formal proof, rounded outward to the digits shown: Λ and ρ up, D down. The column ρ/η, with $\eta = D - \Lambda$, is the exact quotient, rounded up.

$b^{|C\setminus I|}\operatorname{den}(P_{r,I})\,P_{r,I}(a/b)$ plays the role of a small linear form with integer coefficients. There are two differences. The lower bound for the form comes from an arithmetic nonvanishing statement rather than from an analytic asymptotic. The size r of the determinant is chosen in terms of b.

8.1. **Notes on the rows.** Except for the pair $(\frac{1}{3},\frac{2}{3})$ of Section 7, the function of each row is F_s of (2.1) with $s = k/q$, $\omega = q^{-2}$ and $\xi_s = -q^{-2}\Psi(\frac{k}{q},1)$. The table for Λ, the trial measure and the cover are data of the formal proof. All shifts with the same denominator share one table for Λ, indexed by the root classes, and one certificate. Indeed the bounds depend on the shift only through the root class, and the decay depends on it only through the separation of the poles from the contour. For $q = 4$ the root classes $j \equiv kp^{-1} \pmod 4$ of $k = 1$ and $k = 3$ exchange the two classes of primes modulo 4. For $q = 3$ the table uses the primes in the classes modulo 6, and the root classes $j = 1, 2$ correspond to first poles near $p/3$ and $2p/3$. For $q = 7$ there are six root classes. Here $D - \Lambda$ is only about $3 \cdot 10^{-3}$, which explains the large exponent. A small further gain in either constant would lower it substantially (Section 9). For $q = 2$ there is one root class, $j = 1$, with first pole $k_p = \frac{p+1}{2}$. The resulting bound is weaker than the known ones for this number.

9. Limits of the method

The exponents are determined by three constants through $\rho/(D - \Lambda)$. Since $\partial\mu/\partial D = -\rho/\eta^2$, the exponent is most sensitive to the decay where η is small. For the shifts $\frac{1}{4}$ and $\frac{3}{4}$, $\rho/\eta^2 \approx 136$. For the shifts $\frac{k}{7}$ it is about $6.6 \cdot 10^5$.

On the arithmetic side, the profiles $\mathcal{P}_j$ are minima of the bounds of Theorem 4.5. Every further local bound lowers Λ by the integral of the amount by which it lies below $\mathcal{P}_j$. The denominator exponent of the method is the integral of the true exponent e_p, which the local bounds approximate from above. At third order this approximation is close to exhaustion for the shifts $\frac{k}{5}$ and $\frac{k}{7}$. We computed the Smith normal forms exactly at scales $r \le 2400$ and primes $p \le 2m$. There the minimum of the local bounds, with (U_n) included, equals the true exponent at all but a few isolated primes. For the shifts $\frac{k}{7}$ the invariants of valuation at least 4 contribute about $5 \cdot 10^{-6}$ to Λ. The bounds (U_n) gain about $7 \cdot 10^{-4}$. So at third order the local bounds are essentially exhausted. These computations are not part of the proof.

On the analytic side, Remark 6.4 explains why a certificate of the present form cannot do better than the constrained equilibrium measure on the chosen contour. A larger decay rate needs a different contour or a different treatment of the energy problem. The geometry parameter c trades the two sides against each other. A longer border increases D, through the larger node measure. It also increases Λ, through the border rows and columns of the assignment bound.

10. Formalization

The proof of Theorem 1.1 is formalized in Lean 4 [10] with Mathlib [39], together with every theorem, proposition and lemma it uses. The remarks and Section 9 are not part of the formalization. The development is available at

https://github.com/slitsey/catalan-irrationality-measure

It combines analytic and arithmetic proofs with generated rational data. Kernel checks verify the tables for Λ of Section 4 and the decay certificates of Section 6. Numerical computations alone are not used as proof.

The statements of Theorem 1.1 are in the file `ChallengeHybrid.lean`. Their proofs are in `SolutionHybrid.lean`, in the namespace `TrigammaDifferences`. Each final theorem concerns one row of Table 1. Its number ξ is written as a series: the classical form where there is one, and $\Psi(s,t)$ otherwise. The theorem states that $b^{-\mu_{s,t}} < |\xi - a/b|$ for all $a \in \mathbb{Z}$ and all sufficiently large b. The theorems are the following.

- `lambda_plus_measure` and `lambda_minus_measure`: the pairs $(\frac{1}{4}, 1)$ and $(\frac{3}{4}, 1)$, with $\mu_{s,t} = 31.2876$.
- `chi3_measure`: the pair $(\frac{1}{3}, \frac{2}{3})$, with $\mu_{s,t} = 13.7046$.
- `chi3_plus_pi2_measure` and `chi3_minus_pi2_measure`: the pairs $(\frac{1}{3}, 1)$ and $(\frac{2}{3}, 1)$, with $\mu_{s,t} = 9.4998$.
- `fifths1_measure`, ..., `fifths4_measure`: the pairs $(\frac{k}{5}, 1)$ for $k = 1, \dots, 4$, with $\mu_{s,t} = 23.8272$.
- `sevenths1_measure`, ..., `sevenths6_measure`: the pairs $(\frac{k}{7}, 1)$ for $k = 1, \dots, 6$, with $\mu_{s,t} = 2207.8697$.

- `pi2_measure`: the pair $(\frac{1}{2}, 1)$, with $\mu_{s,t} = 6.5253$.

The shared results are in the namespace `CatalanCore.Uniform`.

- `Content.BankShared.Check.hasScaledContent_of_check` is Theorem 4.14. The local bounds of Theorem 4.5 are proved in `Check/Catalogue.lean`, `Check/LP2W.lean`, `Check/Prefix.lean` and `Check/UF.lean`, with the closed rank bound and the branches of Theorem 4.8. Its form for functions with several first poles is `Content.BankShared.KTerm.hasScaledContent_of_kcheck`.
- `Content.BankShared.TwoTerm.LP2HypW.gle` is Theorem 4.6.
- `Content.BankShared.TwoTerm.Pre.prefix_gle` is Theorem 4.7.
- `Content.UFSimple.WrapCheck.catValid_wrap` and `Content.UFSimple.WrapPrefixCheck.catValid_prefix_full` are Theorem 4.8 for $n = 0$ and for $n \geq 1$. Lemma 4.11 is `Content.UFSimple.WrapPrefixClosed.HSplit.det_ne` with `HSplit.dir`.
- `BlockGeom.hasNV0` is Theorem 5.1.
- `Decay.AltDecay.altGeom_hasScaledDecay` is Theorem 6.2. It uses the node energy check `Decay/Nodes/EnergyCheck.lean`, the contour check `Decay/ContourCheck.lean` and the cell checks `Decay/PolyWire3`.
- `measure_of_family_nv0_even` is Theorem 8.1 at the even scales.

The solution has no `sorry` and no `native_decide`. The command `#print axioms` reports only the three standard axioms `propext`, `Classical.choice` and `Quot.sound`. The prime number theorem in arithmetic progressions enters the asymptotics of the denominators. Its formal source is the PNT+ project [40], through OpenAI's workspace and compatibility additions. We use its Wiener–Ikehara theorem.

The proofs were checked with Lean `v4.35.0-rc2` and the matching Mathlib release. The vendored files from OpenAI's repository and from the PNT+ project were converted to Lean's module system and ported to the current toolchain. The repository's README and provenance documents describe the changes. The certificate data were generated with the interval arithmetic of Arb [18], through `python-flint`, and Lean re-checks them.

Disclosure of AI use

The mathematics, the Lean formalization, the certificate generators and the text of this paper were produced by AI systems at the request of the author. The author posed the problems, directed the work and is responsible for the paper. The systems were Claude Opus 5.5, Claude Sonnet 5.5 and Claude Haiku 5.5 (Anthropic), running in Claude Code, and GPT-6.1 Sol (OpenAI), running in Codex. The underlying irrationality proof for Catalan's constant and its Lean formalization are due to OpenAI [31].

Acknowledgements. We thank the authors of Mathlib and of the PNT+ project, and OpenAI for making its formalization available.

References

1. Roger Apéry, *Irrationalité de $\zeta(2)$ et $\zeta(3)$*, Astérisque **61** (1979), 11–13.
2. Keith Ball and Tanguy Rivoal, *Irrationalité d'une infinité de valeurs de la fonction zêta aux entiers impairs*, Invent. Math. **146** (2001), no. 1, 193–207.
3. Frits Beukers, *A note on the irrationality of $\zeta(2)$ and $\zeta(3)$*, Bull. London Math. Soc. **11** (1979), no. 3, 268–272.
4. Hrishik Sai Bojnal, *An irrationality exponent below 2402 for Catalan's constant*, GitHub repository, October 15, 2026, https://github.com/proof-factory/catalan-exponent; design search and certification by Claude Fable 5.1 (Anthropic), 2026.
5. Francis Brown, *Mellin transforms, transfinite diameter and rational approximations of integrals*, 2026, Preprint, 22 April 2026, arXiv:2604.20741.
6. Frank Calegari, Vesselin Dimitrov, and Yunqing Tang, *The linear independence of 1, $\zeta(2)$, and $L(2,\chi_{-3})$*, 2024, arXiv:2408.15403.
7. ———, *Arithmetic holonomy bounds and effective Diophantine approximation*, 2025, Theorem 5.1 gives $\mu(L(2,\chi_{-3})) \le 24781$, arXiv:2510.04156.
8. Marcos Costa Santos Carreira, *Determinant constructions for the Dirichlet beta value at four*, Preprint, 9 October 2026, doi:10.5281/zenodo.23326647, 2026.
9. ———, *General designs and an irrationality exponent for Catalan's constant*, Preprint, 9 October 2026, revised 21 October 2026, doi:10.5281/zenodo.23532239; sources at https://github.com/MarcosCarreira/catalan-designs, 2026.
10. Leonardo de Moura and Sebastian Ullrich, *The Lean 4 theorem prover and programming language*, Automated Deduction – CADE 28, Lecture Notes in Computer Science, vol. 12699, Springer, 2021, pp. 625–635.
11. Payman Eskandari, *Improved rational approximations to Catalan's constant*, 2026, Preprint, 22 September 2026, arXiv:2609.26354.
12. Payman Eskandari, Kumar Murty, and Yusuke Nemoto, *Mixed motives and linear forms in the Catalan constant*, 2025, Preprint, October 2025 (v3: 29 August 2026), arXiv:2510.20648.
13. Aabir Fauzan, *$\zeta(5)$ is irrational*, Preprint v1, September 17, 2026, Zenodo, doi:10.5281/zenodo.22826419, 2026.
14. Peter J. Forrester, *Meet Andréief, Bordeaux 1886, and Andreev, Kharkov 1882–83*, 2018, arXiv:1806.10411.
15. David Niedbala Giraudin, *A new upper bound for the irrationality exponent of $\zeta(2)$*, 2026, Preprint, 2 October 2026, arXiv:2610.02912.
16. ———, *A new upper bound for the irrationality exponent of $\zeta(3)$*, 2026, Preprint, 22 September 2026, arXiv:2609.26980.
17. R. K. Jha, *A finite irrationality measure for Catalan's constant*, Preprint v1, October 8, 2026, doi:10.5281/zenodo.23237269; Lean sources at https://github.com/rohitkrjha/catalan-measure, 2026.
18. Fredrik Johansson, *Arb: efficient arbitrary-precision midpoint-radius interval arithmetic*, IEEE Trans. Comput. **66** (2017), no. 8, 1281–1292, arXiv:1611.02831.
19. Roelof Koekoek, Peter A. Lesky, and René F. Swarttouw, *Hypergeometric orthogonal polynomials and their q-analogues*, Springer Monographs in Mathematics, Springer, Berlin, 2010.
20. Christian Krattenthaler and Tanguy Rivoal, *On a linear form for Catalan's constant*, South East Asian J. Math. Math. Sci. **6** (2008), no. 2, 3–15, arXiv:0810.1927.
21. Christian Krattenthaler and Wadim Zudilin, *Hypergeometry inspired by irrationality questions*, Kyushu J. Math. **73** (2019), no. 1, 189–203, arXiv:1802.08856.
22. Li Lai, *The linear independence of 1, $\zeta(3)$, and $\zeta(4)$*, 2026, Preprint, 5 October 2026 (v2: 15 October 2026), arXiv:2610.06838.

23. Elyasheev Leibtag, Hila Barkan, Rotem Kalisch, Shachar Weinbaum, Michael Shalyt, and Ido Kaminer, *New directions in irrationality*, 2026, Preprint, 20 October 2026, arXiv:2610.27799; supplement and code at https://github.com/RamanujanMachine/new-directions-in-irrationality.
24. Sean Litsey, *An irrationality measure for Catalan's constant*, Hexagon preprint hexagon:2610.00187v4, https://hexagonmath.org/2610.00187v4, 2026.
25. Jingwen Liu, Kai Jiang, and Pingwen Zhang, *Irrationality exponents of logarithms of positive rational numbers*, 2026, Preprint, 7 October 2026, revised 8 October 2026 (v2), arXiv:2610.10192.
26. Junqi Liu, Jujian Zhang, and Lihong Zhi, *A formal proof of the irrationality of* $\zeta(3)$ *in Lean 4*, 2025, arXiv:2503.07625.
27. Karl Löwner, *Über monotone Matrixfunktionen*, Math. Z. **38** (1934), 177–216.
28. Raffaele Marcovecchio, *The Rhin–Viola method for* $\log 2$, Acta Arith. **139** (2009), no. 2, 147–184.
29. Yu. V. Nesterenko, *On the linear independence of numbers*, Vestnik Moskov. Univ. Ser. I Mat. Mekh. (1985), no. 1, 46–49, In Russian; English translation: Moscow Univ. Math. Bull. 40 (1985), no. 1, 69–74.
30. E. M. Nikishin and V. N. Sorokin, *Rational approximations and orthogonality*, Translations of Mathematical Monographs, vol. 92, American Mathematical Society, Providence, RI, 1991.
31. OpenAI, *Catalan's constant is irrational*, Preprint dated September 24, 2026, released October 6, 2026, with a Lean formalization, available at the original paper PDF and the Lean formalization (repository commit `fd4aeeb`, October 7, 2026), 2026.
32. ———, *The irrationality exponent of* π *is 2*, Preprint dated September 24, 2026, released October 6, 2026, https://github.com/openai/math/tree/main/preprints/The-irrationality-exponent-of-pi-is-2-September-24-2026, 2026.
33. Thomas Prellberg, *Three irrationality results for the dilogarithm: Rhin–Viola and Viola–Zudilin constructions at* $-1/4$, $1/5$, *and* $-1/3$, 2026, Preprint, 7 September 2026, arXiv:2609.06895.
34. Georges Rhin and Carlo Viola, *On a permutation group related to* $\zeta(2)$, Acta Arith. **77** (1996), no. 1, 23–56.
35. ———, *The group structure for* $\zeta(3)$, Acta Arith. **97** (2001), no. 3, 269–293.
36. Tanguy Rivoal and Wadim Zudilin, *Diophantine properties of numbers related to Catalan's constant*, Math. Ann. **326** (2003), no. 4, 705–721.
37. Edward B. Saff and Vilmos Totik, *Logarithmic potentials with external fields*, Grundlehren der mathematischen Wissenschaften, vol. 316, Springer, Berlin, 1997.
38. Zhi-Wei Sun, *Catalan's constant is irrational*, 2026, arXiv:2609.04176.
39. The mathlib Community, *The Lean mathematical library*, Proceedings of the 9th ACM SIGPLAN International Conference on Certified Programs and Proofs (CPP 2020), 2020, pp. 367–381.
40. The PrimeNumberTheoremAnd contributors, *The PNT+ project: a Lean formalization of the prime number theorem and related results*, https://github.com/AlexKontorovich/PrimeNumberTheoremAnd, The vendored compatibility additions used here are recorded in the repository's `NOTICE` and `PORTING.md`.
41. Dane Wachs, *A note on a recent claimed proof of the irrationality of Catalan's constant*, 2026, arXiv:2609.22339.
42. Ryan Xu, *Improved* $\kappa = 2$ *trial:* $\mu(G) \leq 727.03$, GitHub issue, October 22, 2026, https://github.com/slitsey/catalan-irrationality-measure/issues/5; data and scripts at https://github.com/ry-xu/slightlysupervised, 2026.
43. Doron Zeilberger and Wadim Zudilin, *The irrationality measure of* π *is at most* $7.103205334137\ldots$, Mosc. J. Comb. Number Theory **9** (2020), no. 4, 407–419, arXiv:1912.06345.

44. Wadim Zudilin, *A few remarks on linear forms involving Catalan's constant*, Chebyshevskii Sb. **3** (2002), no. 2(4), 60–70, In Russian, arXiv:math/0210423.
45. ———, *An Apéry-like difference equation for Catalan's constant*, Electron. J. Combin. **10** (2003), Research Paper 14, arXiv:math/0201024.
46. ———, *Two hypergeometric tales and a new irrationality measure of* $\zeta(2)$, Ann. Math. Québec **38** (2014), no. 1, 101–117, arXiv:1310.1526.
47. ———, *A determinantal approach to irrationality*, Constr. Approx. **45** (2017), no. 2, 301–310, arXiv:1507.05697.
48. ———, *Arithmetic of Catalan's constant and its relatives*, Abh. Math. Semin. Univ. Hambg. **89** (2019), no. 1, 45–53, arXiv:1804.09922.

Email address: sean.litsey.math@gmail.com