

# Factorization – a three year journey

Preview Release v2.6

Author: Essbee Vanhoutte

Date: December 2025

To do before release:

1. Check for consistency with the signs in my quadratics.... right side of the congruence should be negative roots and negative linear coefficient. Left side should be negative roots and positive coefficient. And even though on the right side this always produces a positive linear term and visa versa on the left side.. I should still be consistent with it for the sake of clarity.
2. Rewrite chapter IV to show the simpler modular inverse calculation

## Preface

This paper is dedicated to my family and friends, among whom are my former managers who gave me the time and support to grow my interest in cryptography and eventually the math behind it. When I started this journey three years ago, as a high-school drop-out with limited math education, this seemed like an impossible mountain to scale. This paper documents my findings, and proposes an improvement to known factorization algorithms.

## Chapters

- I. Integer to Quadratic (pg. 1)
- II. Fermat's factorization method (pg. 2)
- III. Quadratic to Quadratic congruence (pg. 3)
- IV. Combining modular linear coefficient solutions (pg. 4)
- V. Bridge to Quadratic Sieve (pg. 7)
- VI. Quadratic coefficients (pg. 8)
- VII. Lifting coefficients (pg. 9)
- VIII. Improved Quadratic Sieve (pg. 9)
- IX. Conclusion (pg. 11)
- X. References (pg. 11)

### I. Integer to Quadratic

In attempting to break factorization, it is important to find the correct representation of the problem.

To describe a problem, one of your first steps will be to define some type of algebraic expression with variables representing the solutions you are trying to find. The Number Field Sieve algorithm<sup>1</sup> is a great example of this, where the very beginning of the algorithm is the polynomial selection step.

I would like to preface this chapter, by giving a short review of the steps that led me to my representation of quadratics of the form:

$$x^2 + y_0x + N = 0 \text{ and } x^2 + y_1x - N = 0. \text{ (And eventually } zx^2 + y_0x + N = zx^2 + y_1x - N, \text{ but we ignore the quadratic coefficients until later)}$$

This entire paper is about semi-primes. A semi-prime is a composite number, which has exactly 2 unique prime factors (excluding itself and 1). For the security of the RSA algorithm, this is the only format that matters. Numbers with more factors then this will actually weaken the RSA algorithm<sup>2</sup> (since it reduces the possible search space for factors).

You will need to make your own generalizations if you want to apply this to composites with more factors.

One example of a semi-prime I've used often in the last three years is:

$$p = 41 \text{ and } q = 107 \Rightarrow N = 41 \times 107 = 4387 \text{ (Where semi-prime } N \text{ is the product of factors } p \text{ and } q)$$

One reason I would often use the same semi-prime while doing research, is that you eventually become so familiar with its structure, that any patterns immediately stand out. However, I always make sure my findings generalize to any other semi-prime as well. In this paper I will exclusively use  $p = 41$  and  $q = 107$ . You can use my proof of concepts to verify these findings against arbitrary semi-primes. I am not an educated Mathematician, hence I will not even attempt to write a paper with algebraic letter-soup.

Around three years ago, one of the first things I very quickly started to zero in on, was using modular reduction on  $N$ . My initial thought process was that perhaps I could find some type of pattern in the remainders that would enable me to construct an algorithm.

For example if  $N = 4387$ :

4387	=	47	mod 70
4387	=	40	mod 69
4387	=	35	mod 68
4387	=	32	mod 67
4387	=	31	mod 66 ( $\sqrt{N}$ )
4387	=	32	mod 65

$$\begin{array}{rcl}
4387 & = & 35 \quad \text{mod } 64 \\
4387 & = & 40 \quad \text{mod } 63 \\
4387 & = & 47 \quad \text{mod } 62
\end{array}$$

The rounded  $\sqrt{N}$  (square root of  $N$ ) is a point of symmetry for the remainders.  
In the above example, starting from mod 66, we get the following sequence as remainders in both directions if we increase or decrease the modulus:

$$31, 32, 35, 40, 47, \dots = 31 + \sum_{i=1}^N 2i-1$$

Any trained mathematician will immediately recognize this sequence as simply 31 plus the squares.  
This is as expected, since we are modulo reducing a number close to its square root.

Now we can rephrase factorization as finding some modulus  $m$ , such that  $N = 0 \text{ mod } m$  (aka  $m$  is a divisor of  $N$ ).  
And we also know that the remainders can be calculate by simply adding squares to the initial remainder at the (rounded) square root of  $N$ .  
Around two years ago, I was maybe a couple of weeks in to my research, when I made these realizations and thought :

"Okay, I can do this, this can't be very hard if I know by how much the remainder increases or decreases with each increment or decrement of the modulus, hence it should be easy to figure out when it hits  $0 \text{ mod } m$ ".

Math is full of problems that are very deceptive. Intuitively it would seem like a straight forward mathematical tool to solve this should exist, but the reality of it is much more complicated. At the surface it looks like a shallow pond, but below the surface, are depths of complexity deeper then the deepest oceans.

Thus if we take  $N \text{ mod } m$ , we need to figure out a way to adjust modulus  $m$  such that the remainder reaches  $0 \text{ mod } m$ . At first I would mess around with some algebraic expressions which had a variable in the modulus. Since to make the remainder change in value, we have to increment or decrement the modulus. However, soon after I was able to generalize it to this, removing any unknowns in the modulus:

$$x^2 + yx + N = 0 \text{ mod } m$$

The root,  $x$ , would represent the smallest factor. In this case 41. And we negate the  $x$  so it becomes negative: -41.  
The coefficient  $y$ , would represent the sum of both factors, in this case  $41 + 107$  or 148.

$$\text{Thus we get: } (-41)^2 + 148 \times -41 + 4387 = 0$$

or simplified:

$$41^2 - 148 \times 41 + 4387 = 0$$

And of-course this will also equal 0 for any mod  $m$ .

Note: My motivation for making the root,  $x$ , negative sign, is that it represents the distance to the factor, rather then simply the factor itself. This is related to how I came up originally to represent factorization this way. I went a little bit into the background on this in my original paper, but truth is, its not very relevant, it's simply one of many ways to represent this problem.

We can also flip the sign for the constant in the quadratic, but then the linear coefficient instead of being  $p + q$  becomes  $p - q$ .

$$\text{If } p + q = 148 \text{ then } p - q = -66$$

$$\begin{array}{l} \text{Setting } y \text{ to } -66 \text{ and flipping the sign of the constant the quadratic becomes: } 41^2 + (-66) \times -41 - 4387 = 0 \\ \text{Hence: } 41^2 - 148 \times 41 + 4387 = 41^2 + 66 \times 41 - 4387 \end{array}$$

We have two unknowns, the root and the linear coefficient (ignoring the quadratic coefficient for now). However, isolating the linear coefficient on both sides - which we can do if they share the same root - of both the quadratics and squaring them mod  $N$  we get:

$$148^2 = 66^2 \text{ mod } 4387 \quad (\text{we can ignore the signs when squaring})$$

And because these two squares are congruent mod  $N$ , we can take the GCD (greatest common divisor) of their difference:

$$\begin{array}{l} \text{GCD}(148 + 66, 4387) = 107 \\ \text{GCD}(148 - 66, 4387) = 41 \end{array}$$

Note: Calculating the greatest common divisor can be done very quickly using the Euclidean algorithm<sup>3</sup>.

Because of this property, just finding the correct pair of linear coefficients of a quadratic will result in the factorization of  $N$ .  
We can formalize this as:

$$x^2 + y_0x + N = 0 \text{ and } x^2 + y_1x - N = 0$$

We can factor  $N$  by finding  $y_0$  and  $y_1$  and taking the GCD of their difference.  
This is very similar to what Fermat's factorization method<sup>4</sup> does.

Further down in the paper we will further simplify this to using the roots rather the linear coefficients. This is not simply a math paper. It is a description of my mathematical journey. Starting with my discoveries made in the first year when I was still a novice with no math education, and slowly adding complexity to it in chronological order as I uncovered more and more in the years that followed.

## II. Fermat's factorization method

It is important to now explain Fermat's factorization method, so we can draw some parallels with my own findings, which accidentally ended up converging with Fermat's factorization method. But this just shows how fundamental this is to the factorization problem.

In its most basic form the procedure is as this:

1. Calculate the rounded  $\sqrt{N}$ .
2. Starting from  $x = \sqrt{N}$ , calculate  $y = x^2 - N$  (do so in a loop while incrementing  $x$ ).
3. If  $y$  is also a square, calculate the greatest common divisor (GCD) on the difference.

Taking  $N = 4387$  as an example:

Step 1 (Calculate the +/- square root of  $N$ ):

$$\sqrt{4387} = 66 \text{ (rounded)}$$

Step 2 (Starting from the square root calculate  $y = x^2 - N$ ):

$66^2 - 4387 = -31$  (not a square)  
 $67^2 - 4387 = 102$  (not a square)  
 $68^2 - 4387 = 237$  (not a square)  
 $69^2 - 4387 = 374$  (not a square)  
 $70^2 - 4387 = 513$  (not a square)  
 $71^2 - 4387 = 654$  (not a square)  
 $72^2 - 4387 = 797$  (not a square)  
 $73^2 - 4387 = 942$  (not a square)  
 $74^2 - 4387 = 1089 = 33^2$  (square)

Step 3 (If  $y$  is also a square, calculate the GCD on the difference.):

$$74^2 = 33^2 \text{ mod } 4387$$

$$\text{GCD}(74 + 33, 4387) = 107$$

$$\text{GCD}(74 - 33, 4387) = 41$$

You may notice that 74 and 33 are simply 148 and 66 divided by 2.

This brings us to another important point. Many such square relations can be found mod  $N$ . What will be different is the amount of times  $N$  is in-between both squares.

$$148^2 = 66^2 + 4 \times 4387 \text{ (four times } N \text{ in between)}$$

$$74^2 = 33^2 + 4387 \text{ (one times } N \text{ in between)}$$

This difference of  $N$ , is related to the quadratic coefficient. More on this in chapter VI.

Do be aware, there are two types of square relations mod  $N$ . One which will yield a trivial factorization (1 or  $N$ ) and one which will yield a non-trivial factorization (a prime factor of  $N$ ). The square relations that will always yield a trivial factorization are of the form:

$$a^2 = -a^2 \text{ mod } N$$

We are not interested in these square relations.

Both Quadratic Sieve<sup>5</sup> and Number Field Sieve - the current fastest factorization algorithms - are more elaborate ways of finding these square relations mod  $N$ . Both algorithms were invented by Carl Pomerance<sup>6</sup>. These algorithms are now almost 40 years old, and not much progress aside from a handful of tweaks to these algorithms has been made since. In my opinion, this is not acceptable. A problem as important as factorization should not go without major progress for 40 years. And simply hoping Quantum computing will solve everything is foolish. Thinking as such is the same as thinking AI will replace everything. It is but an excuse to stop trying. We should never stop trying. The day we stop trying, we surrender ourselves to ignorance.

## III. Quadratic to Quadratic congruence

Going back to representing factorization as the following Quadratic:

$$x^2 + yx + N = 0$$

Finding a root and linear coefficient solution in the integers to this is very hard and not easily solve-able.

One approach is to create "fragments" of a possible solution by reducing everything to mod  $p_0, p_1, p_2, \dots, p_{n-1}$  (where  $p$  is prime), finding integer solutions mod  $p_0, p_1, p_2, \dots, p_{n-1}$  and then combining them. This in essence turns the problem into a subset sum<sup>7</sup> type of problem, because then it becomes a matter of which "fragments", aka solutions mod  $p_i$ , to combine.

Our representation now becomes the same quadratic but with modular reduction:

$$x^2 + yx + N = 0 \text{ mod } m$$

Instead of finding solutions in the integers, we reduce the scope to mod  $m$ .

Example:

$$\begin{aligned}N &= 4387 = 6 \bmod 13 \\p &= 41 = 2 \bmod 13 \\q &= 107 = 3 \bmod 13\end{aligned}$$

The residue of  $N \bmod 13$  is the residue of  $pq \bmod 13$  ( $2 \times 3 = 6 \bmod 13$ ).

The residue of linear coefficient  $y \bmod 13$  is  $p + q \bmod 13$  ( $2 + 3 = 5 \bmod 13$ ).

If the coefficient  $y$ , is  $5 \bmod 13$  and we know our residue of  $N \bmod 13$  is 6 then only  $2 + 3$  and  $3 + 2$  can be our two residues for  $p$  and  $q \bmod 13$ .

Thus the root,  $x$ , is either  $-3$  or  $-2$ . (remember we negate the root)

Plugging in for  $y = 5$  and  $x = -3$  or  $-2 \bmod 13$ :

$$\begin{aligned}-2^2 + 5 \times -2 + 4387 &= 0 \bmod 13 \\ \Rightarrow 4 - 10 + 6 &= 0 \bmod 13\end{aligned}$$

$$\begin{aligned}-3^2 + 5 \times -3 + 4387 &= 0 \bmod 13 \\ \Rightarrow 9 - 15 + 6 &= 0 \bmod 13\end{aligned}$$

In real life we don't know  $y$  is 5 (since we do not know  $p + q$ ).

We do know  $N \bmod 13$  is 6.

Thus all possible coefficient solutions  $y \bmod 13$  can be enumerated by summing up each of the two residues  $\bmod 13$  that multiply to 6.

A coefficient solution  $y$  can also be told to exist, if for a given coefficient value  $y$  a root solution  $x$  exists. This can be trivially determined using the Legendre symbol<sup>8</sup> without actually having to find roots.

All root  $x$  and coefficient  $y$  solutions  $\bmod 13$  that solve the quadratic congruence:

$$\begin{aligned}y = 1 \ x = -5 &\Rightarrow -5^2 + 1 \times -5 + 4387 = 0 \bmod 13 \\y = 1 \ x = -9 &\Rightarrow -9^2 + 1 \times -9 + 4387 = 0 \bmod 13 \\y = 5 \ x = -2 &\Rightarrow -2^2 + 5 \times -2 + 4387 = 0 \bmod 13 \\y = 5 \ x = -3 &\Rightarrow -3^2 + 5 \times -3 + 4387 = 0 \bmod 13 \\y = 6 \ x = -7 &\Rightarrow -7^2 + 6 \times -7 + 4387 = 0 \bmod 13 \\y = 6 \ x = -12 &\Rightarrow -12^2 + 6 \times -12 + 4387 = 0 \bmod 13 \\y = 7 \ x = -1 &\Rightarrow -1^2 + 7 \times -1 + 4387 = 0 \bmod 13 \\y = 7 \ x = -6 &\Rightarrow -6^2 + 7 \times -6 + 4387 = 0 \bmod 13 \\y = 8 \ x = -10 &\Rightarrow -10^2 + 8 \times -10 + 4387 = 0 \bmod 13 \\y = 8 \ x = -11 &\Rightarrow -11^2 + 8 \times -11 + 4387 = 0 \bmod 13 \\y = 12 \ x = -4 &\Rightarrow -4^2 + 12 \times -4 + 4387 = 0 \bmod 13 \\y = 12 \ x = -8 &\Rightarrow -8^2 + 12 \times -8 + 4387 = 0 \bmod 13\end{aligned}$$

#### IV. Combining modular linear coefficient solutions

Now that we can find (linear) coefficient solutions  $\bmod p$ . Let us calculate coefficient solutions  $\bmod p_0, p_1, p_2, \dots, p_{n-1}$  now and combine the results into  $\bmod m$  (where  $m = p_0 \times p_1 \times p_2 \times \dots \times p_{n-1}$ )

$$\begin{aligned}y \bmod 3 &= \{1, 2\} \\y \bmod 5 &= \{2, 3\} \\y \bmod 7 &= \{0, 1, 6\} \\y \bmod 11 &= \{1, 2, 5, 6, 9, 10\} \\y \bmod 13 &= \{1, 5, 6, 7, 8, 12\}\end{aligned}$$

These are the solutions  $\bmod 3, 5, 7, 11, 13$  that solve the coefficient of the linear term in:  $x^2 + yx + N = 0 \bmod m$

If we calculate the solution set  $\bmod 15015$  we get:

$$\begin{aligned}y \bmod 15015 &= \{83, 97, 98, 112, 148, 188, 203, 287, 307, 343, 358, 398, 428, 463, 482, 512, 538, 617, 643, 658, 727, 742, 812, 827, 853, 937, 967, 1007, 1022, \\&1028, 1058, 1072, 1112, 1253, 1267, 1282, 1288, 1358, 1373, 1442, 1457, 1462, 1483, 1513, 1553, 1567, 1568, 1618, 1637, 1652, 1813, 1828, 1847, 1897, \\&1912, 1982, 2003, 2008, 2023, 2092, 2107, 2113, 2177, 2183, 2198, 2267, 2282, 2393, 2437, 2443, 2458, 2477, 2528, 2542, 2612, 2638, 2653, 2722, 2723, \\&2737, 2738, 2807, 2822, 2828, 2848, 2932, 2983, 3002, 3023, 3037, 3067, 3158, 3178, 3262, 3268, 3277, 3283, 3353, 3368, 3437, 3452, 3478, 3548, 3563, \\&3613, 3632, 3647, 3697, 3752, 3808, 3823, 3892, 3893, 3907, 3947, 3977, 3983, 3998, 4003, 4087, 4102, 4178, 4192, 4193, 4207, 4388, 4402, 4432, 4438, \\&4453, 4493, 4523, 4577, 4607, 4633, 4648, 4718, 4753, 4802, 4817, 4907, 4922, 4948, 4978, 5032, 5062, 5102, 5117, 5153, 5257, 5312, 5348, 5362, 5363, \\&5377, 5468, 5543, 5557, 5572, 5578, 5608, 5648, 5663, 5732, 5747, 5803, 5818, 5858, 5908, 5923, 5942, 5972, 6007, 6077, 6103, 6118, 6187, 6202, 6272, \\&6287, 6293, 6313, 6397, 6467, 6488, 6518, 6532, 6572, 6623, 6727, 6733, 6742, 6748, 6818, 6832, 6833, 6902, 6917, 6943, 6973, 7013, 7027, 7028, 7078, \\&7097, 7112, 7118, 7162, 7273, 7288, 7357, 7372, 7442, 7448, 7463, 7468, 7547, 7552, 7567, 7573, 7643, 7658, 7727, 7742, 7853, 7897, 7903, 7918, 7937, \\&7987, 7988, 8002, 8042, 8072, 8098, 8113, 8182, 8183, 8197, 8267, 8273, 8282, 8288, 8392, 8443, 8483, 8497, 8527, 8548, 8618, 8702, 8722, 8728, 8743, \\&8813, 8828, 8897, 8912, 8938, 9008, 9043, 9073, 9092, 9107, 9157, 9197, 9212, 9268, 9283, 9352, 9367, 9407, 9437, 9443, 9458, 9472, 9547, 9638, 9652, \\&9653, 9667, 9703, 9758, 9862, 9898, 9913, 9953, 9983, 10037, 10067, 10093, 10108, 10198, 10213, 10262, 10297, 10367, 10382, 10408, 10438, 10492, \\&10522, 10562, 10577, 10583, 10613, 10627, 10808, 10822, 10823, 10837, 10913, 10928, 11012, 11017, 11032, 11038, 11068, 11108, 11122, 11123, 11192, \\&11207, 11263, 11318, 11368, 11383, 11402, 11452, 11467, 11537, 11563, 11578, 11647, 11662, 11732, 11738, 11747, 11753, 11837, 11857, 11948, 11978, \\&11992, 12013, 12032, 12083, 12167, 12187, 12193, 12208, 12277, 12278, 12292, 12293, 12362, 12377, 12403, 12473, 12487, 12538, 12557, 12572, 12578, \\&12622, 12733, 12748, 12817, 12832, 12838, 12902, 12908, 12923, 12992, 13007, 13012, 13033, 13103, 13118, 13168, 13187, 13202, 13363, 13378, 13397, \\&13447, 13448, 13462, 13502, 13532, 13553, 13558, 13573, 13642, 13657, 13727, 13733, 13748, 13762, 13903, 13943, 13957, 13987, 13993, 14008, 14048, \\&14078, 14162, 14188, 14203, 14273, 14288, 14357, 14372, 14398, 14477, 14503, 14533, 14552, 14587, 14617, 14657, 14672, 14708, 14728, 14812, 14827,\end{aligned}$$

14867, 14903, 14917, 14918, 14932 }

Each of these solutions mod 15015 represents a unique combination of solutions mod 3, 5, 7, 11 and 13 (Cartesian product). There is a number theoretical trick we can use to make finding these solutions mod 15015 a lot easier.

Lets say that we want to calculate the following combination for mod 15015:

$$\begin{aligned}y &= 1 \bmod 3 \\y &= 3 \bmod 5 \\y &= 1 \bmod 7 \\y &= 5 \bmod 11 \\y &= 5 \bmod 13\end{aligned}$$

For each solutions mod  $p$ , we divide and then multiple by every other prime. Using the following procedure:

$$\begin{aligned}m &= \text{total modulus} \\p &= \text{prime}\end{aligned}$$

$$\begin{aligned}a &= m / p \\b &= y \times a^{-1} \bmod p \\y &= a \times b \bmod m\end{aligned}$$

For  $y = 1 \bmod 3$  we get:

$$\begin{aligned}a &= 15015 / 3 = 5005 \\b &= 1 \times 1 = 1 \bmod 3 \\y &= 5005 \times 1 = 5005 \bmod 15015\end{aligned}$$

This gives us 5005 mod 15015

$$5005 = 5 \times 7 \times 11 \times 13 \text{ and } 5005 \bmod 3 = 1$$

Hence this simply changes  $1 \bmod 3$  to mod 15015 by adding multiples of mod 5,7,11,13 while keeping it congruent to  $1 \bmod 3$ .

Now to do the rest:

For  $y = 3 \bmod 5$  we get:

$$\begin{aligned}a &= 15015 / 5 = 3003 \\b &= 3 \times 2 = 1 \bmod 5 \\y &= 3003 \times 1 = 3003 \bmod 15015\end{aligned}$$

This gives us 3003 mod 15015

For  $y = 1 \bmod 7$  we get:

$$\begin{aligned}a &= 15015 / 7 = 2145 \\b &= 1 \times 5 = 5 \bmod 7 \\y &= 2145 \times 5 = 10725 \bmod 15015\end{aligned}$$

This give us 10725 mod 15015

For  $y = 5 \bmod 11$  we get:

$$\begin{aligned}a &= 15015 / 11 = 1365 \\b &= 5 \times 1 = 5 \bmod 11 \\y &= 1365 \times 5 = 6825 \bmod 15015\end{aligned}$$

This gives us 6825 mod 15015

For  $y = 5 \bmod 13$  we get:

$$\begin{aligned}a &= 15015 / 13 = 1155 \\b &= 5 \times 6 = 4 \bmod 13 \\y &= 1155 \times 4 = 4620 \bmod 15015\end{aligned}$$

This gives us 4620 mod 15015

$$\text{Adding the results for mod 3,5,7,11,13 together: } 5005 + 3003 + 10725 + 6825 + 4620 = 148 \bmod 15015$$

In my original paper I would call these intermediate results, partial results. I think that is a fitting name. When we sum up these partial results, we get 148 mod 15015. This way we can reduce finding combinations modulo a composite number to summing together partial results constructed from the prime factors of the composite modulus.

Now what we could do is, calculate all the possible coefficient solutions mod 3,5,7,11,13 and then use the above calculations to create partial results from them mod 15015:

Before:

$y \bmod 3 = \{ 1, 2 \}$   
 $y \bmod 5 = \{ 2, 3 \}$   
 $y \bmod 7 = \{ 0, 1, 6 \}$   
 $y \bmod 11 = \{ 1, 2, 5, 6, 9, 10 \}$   
 $y \bmod 13 = \{ 1, 5, 6, 7, 8, 12 \}$

After:

$y \bmod 3 = \{ 5005, 10010 \}$   
 $y \bmod 5 = \{ 12012, 3003 \}$   
 $y \bmod 7 = \{ 0, 10725, 4290 \}$   
 $y \bmod 11 = \{ 1365, 2730, 6825, 8190, 12285, 13650 \}$   
 $y \bmod 13 = \{ 6930, 4620, 11550, 3465, 10395, 8085 \}$

One of my original research approaches was the insight that if we could grow the modulus by adding more and more primes, eventually  $p + q$  will end up being the smallest solution mod  $m$ . The other solutions will keep growing to a number bigger than  $N$ .

So my initial idea is, if we generate these partial results, and we select one partial result from each prime modulus and sum them together mod  $m$ , how do we find the smallest sum mod  $m$ ? This however is a modular multiple-choice subset-sum problem. Not easily solve-able.

In my original paper I also discussed constructing partial results in mod  $m_0$  and  $m_1$  each from unique primes and as long as  $m_0$  and  $m_1$  are large enough, then we can use the intersection between both sets of solutions to further narrow down the possible set of solutions.

For example:

Mod  $m_0 = 3 \times 7 \times 13 \times 19$  (5187)

$y \bmod 3 = \{ 1729, 3458 \}$   
 $y \bmod 7 = \{ 0, 4446, 741 \}$   
 $y \bmod 13 = \{ 1197, 798, 1995, 3192, 4389, 3990 \}$   
 $y \bmod 19 = \{ 3003, 3822, 1638, 2457, 273, 4914, 2730, 3549, 1365, 2184 \}$

Mod  $m_1 = 5 \times 11 \times 17 \times 23$  (21505)

$y \bmod 5 = \{ 8602, 12903 \}$   
 $y \bmod 11 = \{ 13685, 5865, 3910, 17595, 15640, 7820 \}$   
 $y \bmod 17 = \{ 0, 12650, 10120, 16445, 7590, 13915, 5060, 11385, 8855 \}$   
 $y \bmod 23 = \{ 0, 18700, 7480, 1870, 17765, 14960, 6545, 3740, 19635, 14025, 2805 \}$

From  $m_0$  we select:  $1729 + 4446 + 798 + 3549 = 148 \bmod 5187$  and from  $m_1$  we select:  $12903 + 3910 + 11385 + 14960 = 148 \bmod 21505$

As you can see,  $148 (p + q)$  can be found as a sum mod  $m_i$ , this will hold true for any modulus bigger than  $p + q$ . Hence by inspecting intersections between solution sets, we can quickly narrow it down to one single solution. But in practice, since we need a modulus bigger than  $p + q$ , the amount of possible sums, aka the order of the Cartesian product, quickly grows. Hence this is not feasible, but nonetheless, it is an interesting direction to approach this problem.

In my first paper I attempted to find these intersections using the LLL algorithm<sup>9</sup>. However many improvement can be made there, and would I write it today, there would be many things I would change and simplify further.

## V. Bridge to Quadratic Sieve

After my attempt at finding intersections between solutions sets in mod  $m_i$  using LLL, I instead used these findings to generate smooth candidates for the Quadratic Sieve algorithm. I will now quickly describe the transformations I used to achieve this.

Let us factor 4387.

We set the factor base  $b$  to:

$b = \{ 3, 5, 7, 11, 13 \}$

Next using the quadratic formula:  $x^2 + yx + N \bmod b_i$  we calculate all the possible coefficient solutions for each prime in the factor base:

$y \bmod 3 = \{ 1, 2 \}$   
 $y \bmod 5 = \{ 2, 3 \}$   
 $y \bmod 7 = \{ 0, 1, 6 \}$   
 $y \bmod 11 = \{ 1, 2, 5, 6, 9, 10 \}$   
 $y \bmod 13 = \{ 1, 5, 6, 7, 8, 12 \}$

Next we create a hashmap and go over each coefficient and calculate the following linear congruence:

$$x \times N = y^2 \bmod b_i$$

For coefficient solution  $y = 5 \bmod 11$  we would get:

$$x \times 4387 = 5^2 \bmod 11$$
$$\Rightarrow x = 4$$

We save the  $x$  solution as key in the hashmap and the coefficient solution as value:

```

mod 3 = 1 : { 1, 2 }
mod 5 = 2 : { 2, 3 }
mod 7 = 0 : { 0 }, 3 : { 1, 6 }
mod 11 = 5 : { 1, 10 }, 9 : { 2, 9 }, 4 : { 5, 6 }
mod 13 = 11 : { 1, 12 }, 2 : { 5, 8 }, 6 : { 6, 7 }

```

Next we iterate sieve interval  $i$  from 0 to  $i_{n-1}$

For example when  $i = 97$  we check if  $97 \bmod 3, 97 \bmod 5, \dots$  is a key in the hashmap and we collect the results.

```

97 mod 3 = 1 : { 1, 2 }
97 mod 5 = 2 : { 2, 3 }
97 mod 7 = /
97 mod 11 = 9 : { 2, 9 }
97 mod 13 = 6 : { 6, 7 }

```

At  $i = 97$  we found results in 4 out 5 elements in the coefficient solution set for factor base  $b$ .  
We multiply the moduli together for which we found a result:

$$3 \times 5 \times 11 \times 13 = 2145.$$

And if this is bigger then  $\sqrt{i \times N}$  we continue.

Next we calculate the partial results for the coefficient solutions we just collected (Chapter IV):

```

y mod 3 = { 715, 1430 }
y mod 5 = { 1287, 858 }
y mod 11 = { 585, 1560 }
y mod 13 = { 825, 1320 }

```

Next we generate combinations mod 2145 choosing at most one partial result per modulus.

For example:  $715 + 1287 + 585 + 825 = 1267 \bmod 2145$

Lets call this the coefficient candidate  $y$ .

The useful thing about this setup is that if we now calculate  $y^2 - i \times N$  we know the result will be divisible by the moduli from which we collected the partial results.

$$\text{Hence } 1267^2 - 97 \times 4387 = 1179750 \quad (2 \times 3 \times 5^3 \times 11^2 \times 13)$$

All but one of the factors are in the factor base (2) in this example.

The closer  $y^2$  is to  $i \times N$ , the smaller the smooth candidate will be.

Once enough such smooth candidates are found, you finish the rest of the algorithm using the default Quadratic Sieve proceedings.

Which I won't reiterate as this is widely documented. But in short you would use Gaussian Elimination<sup>10</sup> or Block Lanczos<sup>11</sup> to find a combination of smooths that can be multiplied together to form a square relation on both sides of the congruence mod  $N$ .

## VI. Quadratic coefficients

In the above, we generate possible linear coefficients  $y$  and then square them. By subtracting  $i \times N$ , which I shall henceforth refer to as simply  $iN$ , we can then predict at-least some of the factors of the smooth candidates.

Let us explore how this  $iN$  value relates to the coefficient of the quadratic term, an important subject we have not touched on yet.

For example if we have  $y_0 = 148$  we see that  $148^2 - 4 \times 4387 = 66^2$  ( $i = 4$  thus  $iN = 4 \times 4387$ ) satisfies our square relation mod  $N$ .

Also note that  $148^2 - 4 \times 4387 = 66^2$  is the formula for the quadratic discriminant, which makes sense.

We can see where  $i = 4$  comes from when we subtract and add both linear coefficients from each-other and look at the factorization of the result:

$$\begin{aligned}
148 - 66 &= 82 \quad (41 \times 2) \\
148 + 66 &= 214 \quad (107 \times 2)
\end{aligned}$$

When subtracting we get 2 times the lower factor and when adding we get 2 times the upper factor.

We multiply the factors we found, excluding the factors of  $N$  and we get:  $2 \times 2 = 4$ .

Another example when  $y_1 = 3$  and  $y_0 = 1602$ :

$$\begin{aligned}
1602^2 &= 3^2 \bmod N \\
1602 - 3 &= 1599 \quad (41 \times 39) \\
1602 + 3 &= 1605 \quad (107 \times 15)
\end{aligned}$$

Hence we get  $i = 39 \times 15 = 585$  and verifying this:

$$1602^2 - 4387 \times 585 = 3^2$$

And when  $y_1 = 1$  and  $y_0 = 534$ :

$$1^2 = 534^2 \bmod N$$

$$534 - 1 = 41 \times 13$$

$$534 + 1 = 107 \times 5$$

Hence we get  $i = 13 \times 5 = 65$  and verifying this:

$$534^2 - 4387 \times 65 = 1^2$$

This  $i$  value is actually the coefficient for the quadratic term (you may recognize this in the discriminant formula).

In the example that  $y_1 = 1$  and  $y_0 = 534$  we have a  $i$  value of  $13 \times 5$ .

We know from the above explanation in the first chapters that the roots represent the factors of  $N$ .

We can see that the following holds:

$$13 \times (-41)^2 - 1 \times (-41) + 41 = 0 \bmod 4387$$

$$5 \times (-107)^2 + 1 \times (-107) - 107 = 0 \bmod 4387$$

I will define the quadratic coefficient as  $z$ .

However, in the above example we note that if we have an odd linear coefficient we get quadratics of the shape:  $zx^2 + yx + x$ . Because the quadratic for even coefficients is simpler ( $zx^2 + yx$ ) we shall restrict ourselves to these alone.

Thus working with even coefficient if we have  $y_1 = 2$  and  $y_0 = 1068$ , we see that:

$$1068 - 2 = 26 \times 41$$

$$1068 + 2 = 10 \times 107$$

Since we are now working with two even coefficients we get a quadratic of the shape:  $zx^2 + yx$

This means the quadratic coefficients become  $26/2$  and  $10/2$  (since when we take the derivative, it gets multiplied by 2 from the quadratic exponent).

$$13 \times (-41)^2 - 2 \times (-41) = 5 \times 4387 \text{ or } 0 \bmod 4387$$

$$5 \times (-107)^2 + 2 \times (-107) = 13 \times 4387 \text{ or } 0 \bmod 4387$$

And the derivative reveals the other linear coefficient:

$$26 \times (-41) - 2 = -1068$$

And the quadratic for  $y_0 = 1068$ :

$$13 \times (-41)^2 + 1068 \times (-41) = -5 \times 4387 \text{ or } 0 \bmod 4387$$

$$5 \times (-107)^2 + 1068 \times (-107) = -13 \times 4387 \text{ or } 0 \bmod 4387$$

Our quadratic with the addition of quadratic coefficients is now:  $zx^2 + y_0x + N = zx^2 + y_1x - N$

## VII. Lifting coefficients

As we build up to the final algorithm, another important step must first be explained. This is the lifting of our coefficients mod  $p^e$ , where  $e$  is the exponent.

This is very straight forward, so I will just show you a single example.

Let us say we want to lift  $y_0 = 5$  and  $z = 1 \bmod 11$  to mod  $11^2$  with  $N = 4387$ .

We will only focus on lifting solutions where the resulting  $y_1$  is equal to  $0 \bmod p^e$ .

The very first calculation we must perform is figuring out the root for our polynomial:

$$zx^2 + y_0x + N = 0 \bmod 11$$

Plugging in  $z = 1$  and  $y_0 = 5$ :

$$x^2 + 5x + N = 0 \bmod 11$$

The procedure is as following:

We multiply  $y_0$  by the inverse of  $z \bmod p$  and then multiply that result by the inverse of  $2 \bmod p$ .

Inverse of  $z = 1$  is  $1$ . So nothing changes. Inverse of  $2 \bmod 11$  is  $6$  so we get:

$$5 \times 6 = 8 \bmod 11.$$

And plugging in this root we see that the result is correct:

$$8^2 - 5 \times 8 + 4387 = 0 \bmod 11 \quad (\text{note: remember that we need to use negative roots, hence flipping the sign for the linear term})$$

If we have our statement:  $zx^2 + y_0x + N = zx^2 + y_1x - N$

Then any  $y_0$  and  $y_1$  pairings must also share a common root.



Calculating the derivative to get  $y_1$  will yield a 0 solution mod 11 as expected:

$$y_1 = 2 \times 8 - 5 = 0 \text{ mod } 11$$

And the polynomial for  $y_1$ :

$$8^2 - 0 \times 8 - 4387 = 0 \text{ mod } 11$$

If we now set  $y_1 = 0 \text{ mod } 11^2$ , then finding the new root is simply a matter of finding  $8 + 11 \times a \text{ mod } 11^2$

We can apply Hensel's lifting lemma for this.

$$\begin{aligned} x &= 8 \\ s &= (2x)^{-1} = 9 \text{ mod } 11 \\ t &= (zN - x^2) / 11 = 8 \text{ mod } 11 \\ v &= t \times s = 6 \text{ mod } 11 \\ \text{lifted } x &= z^{-1} \times (x + v \times 11) = 74 \text{ mod } 11^2 \end{aligned}$$

And our root for modulo  $11^2$  thus becomes  $x = 74$ .

We can now take the derivative to generate a new  $y_0$ :

$$y_0 = 2 \times 74 + 0 = 148 \text{ mod } 11^2$$

And verifying the polynomial for  $y_0 = 148$ :

$$74^2 - 148 \times 74 + 4387 = 0 \text{ mod } 11^2$$

Hence lifting linear coefficients is very straight forward. The same principles would apply to lifting quadratic coefficients.

### VIII. Improved Quadratic Sieve

We ended the previous chapter having found the following root and linear coefficient mod  $11^2$

$$74^2 - 148 \times 74 + 4387 = 0 \text{ mod } 11^2$$

Setting  $y_1$  to 0 mod  $11^2$  and calculating the quadratic polynomial yields the following result in the integers:

$$74^2 + 0 \times (-74) - 4387 = 33^2$$

Since  $74^2$  and  $33^2$  is square we can take the GCD and we find the factorization of 4387.

Of-course if we have our quadratic  $zx^2 + y_1x - N$ , then  $zx^2$  isn't always automatically square such as happens to be the case in the above example.

However, we can perform trial factorization on  $zx^2$  and the result generated by the quadratic polynomial and solve for a square on both sides using linear algebra. Furthermore, we would only need to factorize  $z$ , since  $x^2$  is always square and we can ignore squares during the linear algebra step when it is performed in  $gf(2)$ .

This setup generates two smooth candidates.

Let us define smooth candidate left, namely  $z$  as  $L$ .

Let us define smooth candidate right, namely  $zx^2 + y_1x - N$  as  $R$ .

Since we will be generating square moduli and setting  $y_1$  to 0,  $R$  can thus also be shortened to  $zx^2 - N$ .

The way we would approach this next, is by using very large square moduli to reduce the bit length of smooth candidates as much as possible. This is the only way to punch past the theoretical 110 digit ceiling for Quadratic Sieve based algorithms. Since we can use the quadratic coefficient to resize the root. We can achieve both fine grain control of the size of the root and the modulus.

To do: Go over a full numerical example once the code is finished.

## IX. Conclusion

Note to self: rewrite this chapter once paper is finished.

We have managed to make many reductions in complexity in the factorization problem. Where traditionally modern variants of Fermat's factorization method, which includes Number Field sieve and Quadratic sieve, had to find smooth numbers within a factor base, to then hopefully complete a square relation mod  $N$ . We have now found a way to instead enumerate the roots of the squares in this square relation by calculating coefficients of the linear term of a quadratic mod  $p_i$ . This is done in an attempt to break the almost 40-year stalemate in factorization algorithms. I would now encourage the reader, to continue this work, as this is merely the beginning, and many unknown lands of modular magic and polynomials lay ahead of us to explore.

## X. References

1. Number Field Sieve: [https://en.wikipedia.org/wiki/General\\_number\\_field\\_sieve](https://en.wikipedia.org/wiki/General_number_field_sieve)
2. RSA algorithm: [https://en.wikipedia.org/wiki/RSA\\_cryptosystem](https://en.wikipedia.org/wiki/RSA_cryptosystem)
3. Euclidean algorithm: [https://en.wikipedia.org/wiki/Euclidean\\_algorithm](https://en.wikipedia.org/wiki/Euclidean_algorithm)
4. Fermat's factorization method: [https://en.wikipedia.org/wiki/Fermat's\\_factorization\\_method](https://en.wikipedia.org/wiki/Fermat's_factorization_method)
5. Quadratic sieve: [https://en.wikipedia.org/wiki/Quadratic\\_sieve](https://en.wikipedia.org/wiki/Quadratic_sieve)
6. Carl Pomerance: [https://en.wikipedia.org/wiki/Carl\\_Pomerance](https://en.wikipedia.org/wiki/Carl_Pomerance)
7. Subset-sum problem: [https://en.wikipedia.org/wiki/Subset\\_sum\\_problem](https://en.wikipedia.org/wiki/Subset_sum_problem)
8. Legendre Symbol: [https://en.wikipedia.org/wiki/Legendre\\_symbol](https://en.wikipedia.org/wiki/Legendre_symbol)
9. LLL algorithm: [https://en.wikipedia.org/wiki/Lenstra–Lenstra–Lovász\\_lattice\\_basis\\_reduction\\_algorithm](https://en.wikipedia.org/wiki/Lenstra–Lenstra–Lovász_lattice_basis_reduction_algorithm)
10. Gaussian Elimination: [https://en.wikipedia.org/wiki/Gaussian\\_elimination](https://en.wikipedia.org/wiki/Gaussian_elimination)
11. Block Lanczos: [https://en.wikipedia.org/wiki/Block\\_Lanczos\\_algorithm](https://en.wikipedia.org/wiki/Block_Lanczos_algorithm)