

Semi-prime factorization – Analysis of Fermat’s Factorization method

By Essbee Vanhoutte
Version 4.5

Foreword

I am self-taught without formal math education as I dropped out of high-school. This represents 3 years of research and teaching myself mathematics.

Before this I was a self-taught software vulnerability researcher who briefly worked at companies such as Microsoft and Trend Micro and has well over 50+ CVEs in software such as Windows OS, OpenSSL and Adobe Reader attributed to their name (often under the handle SandboxEscaper or Polar Bear).

No AI was used for this research, I have only used it in the last couple of weeks to review my paper and get feedback and occasionally to produce code for well documented mathematical functions.

This paper mainly illustrates findings by example and companion code rather than rigorous mathematical notation and proofs. So perhaps rather than a formal academic paper, view this as a mathematical exploration by an enthusiast with prior experience in IT security.

To do: Add references and closing thoughts and clean everything up.

Chapter I – Fermat’s Factorization method

In its most basic form the procedure for Fermat’s Factorization method is as follows (where N is the semi-prime being factored).

1. Calculate $\lceil\sqrt{N}\rceil$
2. Starting from $x = \lceil\sqrt{N}\rceil$, calculate $f(x) = x^2 - N$ in a loop, incrementing x .
3. If $f(x)$ evaluates to a square, y^2 , compute $GCD(x + y, N)$ and $GCD(x - y, N)$

It is plain to see why this works, because if $x^2 - N = y^2$ then it also follows that $N = x^2 - y^2 = (x - y)(x + y)$ hence anytime we have $x^2 \equiv y^2 \pmod{N}$ where $x \not\equiv \pm y \pmod{N}$ the factors can be computed via GCD

Taking $N = 4387$ as an example:

Step 1 (Calculate $\lceil\sqrt{N}\rceil$)

$$\lceil\sqrt{4387}\rceil = 67$$

Step 2 (Starting from $x = \lceil\sqrt{N}\rceil$, calculate $f(x) = x^2 - N$):

$$67^2 - 4387 = 102 \text{ (not a square)}$$

$$\begin{aligned}
68^2 - 4387 &= 237 \text{ (not a square)} \\
69^2 - 4387 &= 374 \text{ (not a square)} \\
70^2 - 4387 &= 513 \text{ (not a square)} \\
71^2 - 4387 &= 654 \text{ (not a square)} \\
72^2 - 4387 &= 797 \text{ (not a square)} \\
73^2 - 4387 &= 942 \text{ (not a square)} \\
74^2 - 4387 &= 1089 = 33^2 \text{ (square)}
\end{aligned}$$

Step 3 (If the result is also a square, calculate the *GCD* on the difference.):

$$74^2 \equiv 33^2 \pmod{4387}$$

$$GCD(74 + 33, 4387) = 107$$

$$GCD(74 - 33, 4387) = 41$$

Both the Quadratic Sieve and Number Field Sieve algorithms (to do: insert references) are expansions on this method.

Chapter II – Analysis

a. Binomial expansion

As seen in chapter 1 the goal is to find the following numbers that we can use to take the *GCD* with:

$$GCD(x + y, N) = q$$

$$GCD(x - y, N) = p$$

x and y can thus be rewritten as the differences of the factors of N divided by 2, $(p + q)/2$ or $(p - q)/2$.

In addition, many such possible x and y solutions exist created by $(ap + bq)/2$ and $(ap - bq)/2$ where a and b are multipliers.

In practice we will not know the factors p and q , but treating them as unknowns we can calculate residues in Z/ℓ (for a prime ℓ) for them using the framework we're establishing here.

If $a = 13$, $b = 5$ and $N = 4387$ (and thus factors as $p = 41$ and $q = 107$)

We get:

$$(13 \times 41 + 5 \times 107)/2 = 534$$

$$(13 \times 41 - 5 \times 107)/2 = -1$$

And we can see this creates the following expression when we square $534 \pmod{N}$:

$$534^2 - 4387 \times 13 \times 5 = 1^2$$

One way to represent x and y is through binomial expansion.

I will shorten binomial expansion as $BE(x-y)^e$.

For example $BE(x-74)^2 = x^2 - 148x + 5476$.

Substituting the binomial term x in $BE(x-74)^e$, with for example 33, we get:

$$BE(33-74)^1 = 33-74 = -41$$

$$BE(33-74)^2 = 33^2 - 148 \times 33 + 5476 = -41^2$$

$$BE(33-74)^3 = 33^3 - 222 \times 33^2 + 16428 \times 33 - 405224 = -41^3$$

$$BE(33-74)^4 = 33^4 - 296 \times 33^3 + 32856 \times 33^2 - 1620896 \times 33 + 29986576 = -41^4$$

Conversely we also see the following occurring if we substitute in the factor as root instead (or swapping a binomial term with the binomial value):

$$BE(41-74)^1 = 41-74 = -33$$

$$BE(41-74)^2 = 41^2 - 148 \times 41 + 5476 = -33^2$$

$$BE(41-74)^3 = 41^3 - 222 \times 41^2 + 16428 \times 41 - 405224 = -33^3$$

$$BE(41-74)^4 = 41^4 - 296 \times 41^3 + 32856 \times 41^2 - 1620896 \times 41 + 29986576 = -33^4$$

In addition, if we have even exponents we can substitute the constant by a multiple of N and generate a polynomial $g(x)$ to solve for 0.

$$BE(41-74)^2 = 41^2 - 148 \times 41 + 5476 \rightarrow g(x) = 41^2 - 148 \times 41 + 4387 = 0$$

$$BE(41-74)^4 = 41^4 - 296 \times 41^3 + 32856 \times 41^2 - 1620896 \times 41 + 29986576 \rightarrow g(x) = 41^4 - 296 \times 41^3 + 32856 \times 41^2 - 1620896 \times 41 + 4387 \times 6565 = 0$$

b. The discriminant

Of-course we cannot blindly perform binomial expansion on random numbers hoping to find a difference of squares. We will need to construct a way to sieve this.

Take for example the following binomial expansion $BE(x-74)^2 = x^2 - 148x + 5476$ and on this quadratic polynomial, we substitute the constant with N (for example $N = 4387$):

$$g(x) = x^2 - 148x + 4387$$

If we now calculate the discriminant of $g(x)$, $DISC(g(x)) = 148^2 - 4 \times 4387 = 66^2$, we see that the result is also square.

And we now have the square relation $148^2 \equiv 66^2 \pmod{4387}$, where $GCD(148-66, 4387) = 41$.

An observation:

From this we can make the following generalization that we can derive the factorization of a semi-prime N , if we find a quadratic polynomial with a constant set to N , or a multiple thereof, and whose discriminant yields a square. And from this it is also true that the discriminant would yield a quadratic residue modulo any prime if the discriminant is square in Z . Hence we can use this property to calculate residues in Z/ℓ where valid solutions in Z/ℓ are a quadratic residue and discard residues for quadratics where this is not the case. Now it is also true, that if we find a polynomial whose discriminant is square in Z then it must have a multiple of the factor of N as root for polynomials $g(x)$ at degree 2. That relation between polynomial and discriminant is still something I am researching. Perhaps this will be useful in a type of number field type of construction, hence why I feel it is important to make note of this finding. In addition, it also links everything back to the polynomials generated with binomial expansion.

c. An extended example

Let $N = 4387$

Let binomial $b = x-534$.

Hence $BE(x-534)^2 = x^2-1068x + 534^2$

We now need to substitute the constant in the resulting polynomial with a multiple of N .

For example let N multiplier $k = 1$:

$$k = 1 \rightarrow g(x) = x^2-1068x + 4387$$

$DISC(g(x))$ is not a square in Z hence we know $k = 1$ cannot be a valid solution.

However if $k = 65$:

$$k = 65 \rightarrow g(x) = x^2-1068x + 4387 \times 65 \text{ and } DISC(g(x)) = 2^2$$

The discriminant is defined as $DISC(g(x)) = (2t)^2$, where t is the unknown binomial term, we can now substitute this into binomial b .

$b(1) = 1-534 = -533$ and $g(533) = 0$. Since $g(x)$ evaluates to 0 at 533 we know we have a correct solution and we can take the gcd using our binomial terms:

$$GCD(1 + 534, 4387) = 107$$

$$GCD(534-1, 4387) = 41$$

In the next chapter we will see how to find these solutions in Z/ℓ

Chapter III – Improvements

a. Calculating polynomial residues in $\mathbb{Z}/\ell$

Let us restrict ourselves to binomial expansions to the second degree.

The way we calculate polynomial residues is that we must consider all possible coefficients whose discriminant evaluates to a quadratic residue.

Here we calculate all possible residues for the quadratic polynomial $ax^2 + bx + Nk$ where $N = 4387$ (you can also calculate them for $ax^2 + bx - Nk$ instead, it just changes the side we are calculating in the expression $x^2 = y^2 \pmod{N}$). We will ignore the cases where the leading coefficient is 0, as this would downgrade the degree of our polynomial. And in addition we will also ignore the case where $k = 0$ for the sake of brevity.

Mod 3

$k = 1, a = 1$ and $b = -1 \rightarrow x = 2, k = 1, a = 1$ and $b = -2 \rightarrow x = 1, k = 1, a = 2$ and $b = 0 \rightarrow x = 1, 2$
 $k = 2, a = 1$ and $b = 0 \rightarrow x = 2, 1, k = 2, a = 2$ and $b = -1 \rightarrow x = 1, k = 2, a = 2$ and $b = -2 \rightarrow x = 2$

Mod 5

$k = 1, a = 1$ and $b = -2 \rightarrow x = 4, 3, k = 1, a = 1$ and $b = -3 \rightarrow x = 2, 1, k = 1, a = 2$ and $b = 0 \rightarrow x = 3, 2,$
 $k = 1, a = 2$ and $b = -1 \rightarrow x = 4, k = 1, a = 2$ and $b = -4 \rightarrow x = 1, k = 1, a = 3$ and $b = 0 \rightarrow x = 1, 4,$
 $k = 1, a = 3$ and $b = -2 \rightarrow x = 2, k = 1, a = 3$ and $b = -3 \rightarrow x = 3, k = 1, a = 4$ and $b = -1 \rightarrow x = 3, 1,$
 $k = 1, a = 4$ and $b = -4 \rightarrow x = 4, 2$
 $k = 2, a = 1$ and $b = 0 \rightarrow x = 4, 1, k = 2, a = 1$ and $b = -1 \rightarrow x = 3, k = 2, a = 1$ and $b = -4 \rightarrow x = 2,$
 $k = 2, a = 2$ and $b = -1 \rightarrow x = 2, 1, k = 2, a = 2$ and $b = -4 \rightarrow x = 3, 4, k = 2, a = 3$ and $b = -2 \rightarrow x = 3, 1,$
 $k = 2, a = 3$ and $b = -3 \rightarrow x = 4, 2, k = 2, a = 4$ and $b = 0 \rightarrow x = 2, 3, k = 2, a = 4$ and $b = -2 \rightarrow x = 4,$
 $k = 2, a = 4$ and $b = -3 \rightarrow x = 1$
 $k = 3, a = 1$ and $b = 0 \rightarrow x = 3, 2, k = 3, a = 1$ and $b = -2 \rightarrow x = 1, k = 3, a = 1$ and $b = -3 \rightarrow x = 4,$
 $k = 3, a = 2$ and $b = -2 \rightarrow x = 2, 4, k = 3, a = 2$ and $b = -3 \rightarrow x = 1, 3, k = 3, a = 3$ and $b = -1 \rightarrow x = 3, 4,$
 $k = 3, a = 3$ and $b = -4 \rightarrow x = 2, 1, k = 3, a = 4$ and $b = 0 \rightarrow x = 4, 1, k = 3, a = 4$ and $b = -1 \rightarrow x = 2,$
 $k = 3, a = 4$ and $b = -4 \rightarrow x = 3$
 $k = 4, a = 1$ and $b = -1 \rightarrow x = 2, 4, k = 4, a = 1$ and $b = -4 \rightarrow x = 1, 3, k = 4, a = 2$ and $b = 0 \rightarrow x = 4, 1,$
 $k = 4, a = 2$ and $b = -2 \rightarrow x = 3, k = 4, a = 2$ and $b = -3 \rightarrow x = 2, k = 4, a = 3$ and $b = 0 \rightarrow x = 2, 3,$
 $k = 4, a = 3$ and $b = -1 \rightarrow x = 1, k = 4, a = 3$ and $b = -4 \rightarrow x = 4, k = 4, a = 4$ and $b = -2 \rightarrow x = 1, 2,$
 $k = 4, a = 4$ and $b = -3 \rightarrow x = 3, 4$

The problem is, that we can Chinese remainder these together to modulo 15, but then the solution set of possible residues grows quickly (the product of the size of both sets of solutions). So we need to come up with a smart way to generate solutions with this setup.

b. Singular quadratic roots vs non-singular quadratic roots

In the above residue mapping, we will observe that sometimes a set of coefficients will have a single root solution, and sometimes it will have two root solutions.

For example, looking at Mod 5 we see the following two coefficient, root pairings:

$k = 1, a = 4$ and $b = -1 \rightarrow x = 3, 1$ Non-singular roots
 $k = 1, a = 3$ and $b = -2 \rightarrow x = 2$ Singular root

Singular roots indicate that the corresponding binomial term has prime ℓ (5 here) as divisor and that the discriminant is also divided by prime ℓ .

We have calculated the above residues using polynomials of the shape $ax^2 + bx + Nk$.

If we calculate the derivative in Z/ℓ we get the residue for linear coefficient b in corresponding to a quadratic with the signs flipped $ax^2 + bx - Nk$ (the other side of the expression $x^2 = y^2 \pmod{N}$ that we are trying to solve for).

Calculating the derivative for coefficients with a singular root in Z/ℓ gives us a derivative that is 0 in Z/ℓ . Since a singular root means $DISC(g(x)) = 0 \pmod{\ell}$ and since $DISC(g(x)) = (2t)^2$, we have that $t \equiv 0 \pmod{\ell}$. (strictly speaking about degree 2 polynomials here).

For example:

Using $k = 1, a = 3$ and $b = -2 \rightarrow x = 2$ we construct the polynomial $g(x) = 3 \times 2^2 - 2 \times 2 + 2$ in $Z/5$.

And we see that the derivative $g(x)' \equiv 2 \times 3 \times 2 - 2 \equiv 0 \pmod{5}$ (or in other words 0 in $Z/5$).

This implies that the binomial term y , in binomial $b = t - y$ equals to 0 in $Z/5$.

Non-singular roots always yield non-zero residues for the binomial terms in Z/ℓ .

C. p-adic lifting of polynomials in Z/ℓ

Let us say we have the following coefficient and root pairing from our residue mapping:

$k = 4, a = 3$ and $b = -1 \rightarrow x = 1$ in $Z/5$ gives us polynomial $g(x) = 3 \times 1^2 - 1 \times 1 + 4387 \times 4 = 0 \pmod{5}$

This gives us all of the following possible solutions in $Z/5^2$ where also the derivative yields 0 in $Z/5^2$, for the sake of brevity we will focus on these cases only:

$k = 4, a = 3$ and $b = -1 \rightarrow x = 21, k = 4, a = 8$ and $b = -6 \rightarrow x = 16, k = 4, a = 13$ and $b = -11 \rightarrow x = 11$
 $k = 4, a = 18$ and $b = -16 \rightarrow x = 6, k = 4, a = 23$ and $b = -21 \rightarrow x = 1$
 $k = 9, a = 3$ and $b = -11 \rightarrow x = 6, k = 9, a = 8$ and $b = -16 \rightarrow x = 1, k = 9, a = 13$ and $b = -21 \rightarrow x = 21$
 $k = 9, a = 18$ and $b = -1 \rightarrow x = 16, k = 9, a = 23$ and $b = -6 \rightarrow x = 11$
 $k = 14, a = 3$ and $b = -21 \rightarrow x = 16, k = 14, a = 8$ and $b = -1 \rightarrow x = 11, k = 14, a = 13$ and $b = -6 \rightarrow x = 6$
 $k = 14, a = 18$ and $b = -11 \rightarrow x = 1, k = 14, a = 23$ and $b = -16 \rightarrow x = 21$
 $k = 19, a = 3$ and $b = -6 \rightarrow x = 1, k = 19, a = 8$ and $b = -11 \rightarrow x = 21, k = 19, a = 13$ and $b = -16 \rightarrow x = 16$
 $k = 19, a = 18$ and $b = -21 \rightarrow x = 11, k = 19, a = 23$ and $b = -1 \rightarrow x = 6$
 $k = 24, a = 3$ and $b = -16 \rightarrow x = 11, k = 24, a = 8$ and $b = -21 \rightarrow x = 6, k = 24, a = 13$ and $b = -1 \rightarrow x = 1$
 $k = 24, a = 18$ and $b = -6 \rightarrow x = 21, k = 24, a = 23$ and $b = -11 \rightarrow x = 16$

Each of these solutions map to the same solution in $Z/5$. We see a total of 25 (or 125 if we also count those with non-zero derivatives $Z/5^2$) possible solutions in $Z/5^2$

generated from a single solution in $Z/5$. I refer to p-adic lifting, but usually it is used in the context of lifting roots of polynomials, but in the context that I use this term, it is also used to lift coefficient residues in Z/ℓ p-adically. So please keep that in mind. I tend to create these abstractions in my head and the names I give these abstract concepts may not be strictly correct in formal mathematical terms. But to me it would seem, that whether we are lifting root residues or coefficient residues, the underlying mechanism are very similar and connected.

D. Converting singular roots to non-singular roots and visa versa

Because we are searching for two squares such that $x^2 = y^2 \pmod{N}$, and since both x and y can also be represented as binomials which can be expanded to the second degree, we can substitute the expression $x^2 = y^2 \pmod{N}$ with the more elaborate: $ax^2 - b_0x + Nk = ax^2 + b_1x - Nk$.

This also means that we can convert singular roots to non-singular roots temporarily, simply by transferring a root to the other side of the equation in Z/ℓ . As long as the side we transfer to has a discriminant that is not divisible by ℓ . (there is a few other edge cases that will complicate things, for example when k divides by ℓ).

For example $1x^2 + 66x - 4387 = 0 \pmod{37}$ has a singular root, while the other side, $1x^2 - 0x + 4387 = 0 \pmod{37}$ has two roots in Z/ℓ .

$$b_1 = 66 \rightarrow 1 \times 4^2 + 66 \times 4 - 4387 = 0 \pmod{37}$$

$$b_0 = 0 \rightarrow 1 \times 4^2 - 0 \times 33 + 4387 = 0 \pmod{37}$$

$$b_0 = 0 \rightarrow 1 \times 33^2 - 0 \times 33 + 4387 = 0 \pmod{37}$$

A polynomial that produces the other singular root can also be found, in this case all we have to do is flip the sign on linear coefficient b_1 , then the singular root becomes the other non-singular root for b_0 (we must also swap a and k , but both are 1 here):

$$b_1 = -66 \rightarrow 1 \times 33^2 - 66 \times 33 - 4387 = 0 \pmod{37}$$

One way this ends up being useful is that with this construction we never have to p-adically lift singular roots (except for a couple of edge cases like $Z/2^e$ or when the leading coefficient or k is divisible by ℓ), as we can just change them to non-singular, lift them and use them as lifted root for the singular side.

Chapter IV - Residue and two-sided sieving

a. Residue sieving

As mentioned earlier, we cannot simply precompute residues and combine them with Chinese Remainder Theorem hoping to find a valid solution, because the possible set of solutions grows too quickly. The easiest way to take advantage of these residues is with a Quadratic Sieve style sieving setup. In addition, due to the setup we have build in prior chapter, there is a unique two-sided variation that we can construct that are not seen in standard SIQS/MPQS variants.

Let us say that $N = 4387$

We can setup our sieve easily such that we always have a guaranteed factor up to a maximum of $\sqrt{N}$. This can be achieved in two different ways, either we can do it the way SIQS/MPQS does by encoding the modulus into the leading coefficient or through using a modulus instead.

Lets say at each interval step, we always want 11 as a divisor for every B-smooth candidate generated. SIQS/MPQS uses the following procedure:

$N = 4387$ (semi-prime to be factored)

$m = 11$ (modulus)

Next we generate a quadratic polynomial this way:

First find x such that $x^2 - 4387 \equiv 0 \pmod{m}$

This gives us that $x = \pm 3 \pmod{11}$

Calculate the coefficient a, b and constant c as following:

$$a = m$$

$$b = 2x$$

$$c = (x^2 - N)/m$$

This gives us:

$$g(x) = 11x^2 + 6x - 398$$

And we sieve the roots of this polynomial. This is a non-monic quadratic, meaning it can be translated into a monic by multiplying the root and constant with the leading coefficient. This in essence multiplies the polynomial value with the leading coefficient.

For example if evaluating $g(x)$ with 5:

$$g(5) = -93$$

Translating non-monic $g(x)$ into monic $h(x)$, $h(x)$ becomes:

$h(x) = z^2 + 6z - (398 \times 11)$ and evaluating with $z = x \times a$ we get $h(55) = 11 \times -93$, and we see this effectively transfers the leading coefficient to the polynomial value.

Furthermore we can add linear coefficient b to the constant c , using the formula $c - (b/2)^2$, making c equal to N again, and now evaluating $u^2 - 4387$ with $u = z + (b/2)$ gives use $58^2 - 4387 = 11 \times -93$ which is exactly the format that we require.

The problem with this is, that the smallest polynomial values that we can generate with $x^2 - N$, live near $x = \lceil \sqrt{N} \rceil$, and with each increment of x , the polynomial value changes by roughly $2\sqrt{N}$, meaning if we are to sieve that

range, our polynomial values will be roughly of size $2\sqrt{N}$. So no matter what we do, we cannot strip away more known factors than roughly $\sqrt{N}$, since we're in essence just dividing the polynomial values in advance.

But I independently also (re)discovered that we can simply use a modulus instead without manipulating the quadratic (although this is still a fairly basic number theoretical setup and I assume would be known already, hence I do not want to claim novelty here):

to do: show procedure

This was something that I achieved at around 1.5 years into my research. This gives us performance similar to SIQS, while still allowing for the same transformations on-top of this. The next question I was then left with was: How to improve from here, can we use these SIQS/MPQS type transformations, which we have not utilized yet, to add more performance?

b. Two-sided sieving in the rationals with linear poly split of the unknown part

Without the addition of number fields, you really cannot sieve with a modulus/known factors greater than $\sqrt{N}$

What I did stumble on, was a mechanism to split the unknown part of the B-smooth into two linear polynomials, that we can presieve and re-arrange how they get paired up.

to do: show procedure

c. Benchmarks

to do

Closing thoughts and future work

to do