



Therapeutic Diet and/or Fluids Procedure

1. Scope

Site	Operational Area	Applicable to
Armada Kalamunda Group	All services	Catering Staff, Dietetics, Nursing, Speech Pathology

2. Purpose

To ensure correct processes are followed when ordering therapeutic diets or fluids for patients.

The *Nutrition Standards for adult inpatients in WA hospitals* (2018) states “WA Health has a duty of care to ensure inpatients have access to safe, appropriate and adequate food and fluids”.

Multiple departments and staff are involved in ensuring patients receive their correct meal. This procedure will ensure all staff are aware of their roles and responsibilities in the process of ordering therapeutic diet and fluids.

The delivery of incorrect meals and/or fluids may occur within the hospital setting and consequently place the patient at risk, e.g. If a patient receives regular soup instead of level 3 moderately thickened soup, they may be at risk of aspiration.

Risk: Patients receiving incorrect meals can result in adverse and potentially life-threatening reactions that may impact the length of inpatient stay and increased costs to the health service. For this reason, there is a “no diet no meal” approach, whereby if there is no diet entered into i-Soft, the kitchen will not provide a meal

3. Procedure

3.1 Diet orders

- Nursing staff to ensure all patients have a diet entered into iSOFT before 0500, 0900, and 1400. Nursing to identify potential food allergies or need for therapeutic diet on admission based on patient/facility handover and enter into iSoft using approved diet terminology [Appendix1](#). When no special diet required as desired (AD) to be entered.
- Catering staff print out ward diet lists at 0500, 0900, and 1400. Catering staff will check for missing diet orders and contact the shift coordinator via page or phone to request missing diets to be entered into iSOFT immediately
- Changes to diets or new admissions (after 0500, 0900 and 1400), to be phoned (x2021), or paged (#445) to catering
- Nursing staff are to notify catering as soon as possible via fax, phone or page regarding any bed changes or ward transfers

3.2 Altering Therapeutic Diets

- Nursing, DT or SP identify need for therapeutic diet
- DT or SP inform the relevant nursing staff about changes
- SP completes a safe swallow chart using template [Appendix 2](#) and places above patient's bedhead for modified diet/fluids
- Nursing, DT or SP change diet in iSOFT using approved diet terminology [Appendix 1](#) and contact catering staff as per 3.1.3

3.3 Process if patient receives incorrect therapeutic diet or fluid

- If patient received an incorrect meal/fluid, review the meal/fluid and match it with the current recommendations (including iSOFT, ward lists, nutrition standards)
- Nurse Coordinator or DT/SP to contact Catering Coordinator, Head Chef, or Chef – Corporate Services
- If able, the Catering Coordinator will come up to the ward and assess the meal/rectify the issue and provide the correct meal/fluid for the patient. SP and/or DT may be consulted if required
- Person who identified the meal as incorrect to complete clinical incident form online via Datix CIMS, noting error, update iSOFT as indicated, and appoint SP or DT as third-party investigator (if completed by nursing staff)
- All clinical incidents to be tabled at Nutrition Management Sub-Committee for review

4. Legislative context

The following documents are required to give effect to this procedure:

- Department of Health Western Australia (HDWA) [Clinical Incident Management Policy MP 0122/19](#)
- HDWA [Clinical Handover Policy MP 0095](#)
- East Metropolitan Health Service [Patient Identification and Procedure Matching EMHS: 21](#)

5. Supporting information

The following documents support the implementation of this procedure:

- Dietitians Association of Australia Nutrition Manual 2009

6. Compliance, monitoring and evaluation

Compliance against this procedure will be evaluated by monitoring of Datix CIMS by the Management of Nutrition Committee.

7. Relevant standards

National Safety and Quality Health Service (NSQHS) Standards (second edition):

- Standard 1 - Action 1.11: The health service organisation has organisation-wide incident management and investigation systems, and:
 - Supports the workforce to recognise and report incidents
 - Supports patients, carers, and families to communicate concerns or incidents
 - Involves the workforce and consumers in the review of incidents

- Provides timely feedback on the analysis of incidents to the governing body, the workforce, and consumers
- Uses the information from the analysis of incidents to improve safety and quality
- Incorporates risks identified in the analysis of incidents into the risk management system
- Regularly reviews and acts to improve the effectiveness of the incident management and investigation systems
- Standard 5 – Action 5.27: The health service organisation that admits patients overnight has systems for the preparation and distribution of food and fluids that include nutrition care plans based on current evidence and best practice
- Standard 5 – Action 5.28: The workforce uses the systems for preparation and distribution of food and fluids to:
 - Meet patients' nutritional needs and requirements
 - Monitor the nutritional care of patients at risk
 - Identify, and provide access to, nutritional support for patients who cannot meet their nutritional requirements with food alone
 - Support patients who require assistance with eating and drinking
- Standard 6 – Action 6.8: Clinicians use structured clinical handover processes that include:
 - Preparing and scheduling clinical handover
 - Having the relevant information at clinical handover
 - Organising relevant clinicians and others to participate in clinical handover
 - Being aware of the patient's goals and preferences
 - Supporting patients, carers, and families to be involved in clinical handover, in accordance with the wishes of the patient
 - Ensuring that clinical handover results in the transfer of responsibility and accountability for care

8. References

1. Department of Health, Western Australia. Nutrition Standards Guidelines for Adult Inpatients in WA Hospitals. 2018
2. Agency for Clinical Innovation, New South Wales. Therapeutic Diet Specifications for Adult Inpatients. 2011

9. Glossary

The following definitions are relevant to this policy.

Term	Definition
Therapeutic Diet	Includes texture modified diets e.g. Level 4 - Pureed diet, – Level 5 - Minced and moist diet, Level 6 – Soft and bite sized diet and nutritionally indicated diets e.g. renal, low fibre, low fat, high energy high protein.
Fluid Levels	Level 2 – mildly thick, Level 3– moderately thick, Level 4 – extremely thick. Includes consistency of all fluids e.g. sauces, and soups.

10. Policy contact (relating to these procedures)

Enquiries relating to this procedure may be directed to:

Title: Coordinator Speech Pathology
 Department: Speech Pathology
 Email: Renee.M.Davis@health.wa.gov.au

11. Review

This procedure will be reviewed and evaluated as required to ensure relevance and currency. At a minimum it will be reviewed within one (1) year after first issue and at least every three (3) years thereafter.

Version	Effective from	Effective to	Amendment(s)
Version 3	19/10/2023	19/10/2026	Scheduled review

12. Authorisation

This procedure has been authorised and issued by the Director Allied Health.

Authorised by	Danielle Kilmurray - Director Allied Health
Authorisation date	19/10/2023
Published date	19/10/2023

Appendix 1: Diet Terminology

Diet	DAA Nut Manual	Description
As Desired (L7 Regular Diet)		Full ward diet with no restrictions
Clear Fluids	P 30	Includes foods that are liquid (fat containing liquids are excluded i.e. all dairy products); Highly restrictive and offers little nutritive value. Fluids include water, clear soup, plain jelly, cordial, soft drinks, flavoured ice blocks, Forti juice.
Diabetic	P 67	Low in sugar and fat. Includes dessert options suitable for people with diabetes
Fasting		No diet; patient will not receive meal(s) due to upcoming procedure/medical test
Gluten Free	P 155	Eliminates all gluten from the diet (wheat, rye, barley, oats) including commercial foods containing gluten and is suitable for patients with Coeliac Disease.
Halal	P 13	A diet suitable for specific religious requirements
High Protein High Energy	P 194	Provides an increased level of protein & total energy to the diet & includes fortification of foods and may also include nourishing snacks/drinks ordered by the Dietitian. Also see Dietetics Prescribing High Energy High Protein (HEHP) Diets Guideline.
Lactose Free	P 170	Eliminates foods that contain lactose including milk/milk products, foods containing whey, milk solids, casein.
Level 1 Slightly Thick Fluids		Thickened Fluids to Level 1 – Slightly thick consistency
Level 2 Mildly thick fluids (L150)		Thickened Fluids to Level 2 – Mildly thick consistency
Level 3 Moderately thick fluids (L400)		Thickened Fluids to Level 3 – Moderately thick consistency
Level 4 Extremely thick fluids (L900)		Thickened Fluids to Level 4 – Extremely thick consistency
Level 3 Liquidised Diet		Can be eaten with a spoon or drunk from a cup. Cannot be eaten with a fork because it drips through the fork prongs. Has a smooth texture with no 'bits' (lumps, fibres, husk, shell or skin, particles of gristle or bone).
Level 4 Pureed Diet		Has a smooth texture with no lumps that does not require chewing, holds shape on a spoon and falls off a spoon in a single spoonful when tilted. Not sticky. Liquids (i.e. sauces) must not separate from the solids.
Level 5 Minced and Moist Diet		Soft and moist, but with no liquid leaking/dripping from the food. Biting is not required, minimal chewing required, lumps of 4mm in size which can be mashed with the tongue. Food can be easily mashed with just a little pressure from a fork. Should be able to scoop food onto a fork with no liquid dripping and no crumbs falling off the fork.
Level 6 Soft and Bite Sized Diet		Soft, tender and moist, but with no thin liquid leaking/dripping from the food. Ability to 'bite off' a piece of food is not required; pieces of food should be no bigger than 1.5cm x 1.5cm in size. Food can be mashed/broken down with pressure from the fork and a knife is not required.
Level 7 Easy Chew Diet		Normal/everyday foods of soft/tender textures, with no hard, tough, chewy, sticky, fibrous, stringy, crunchy, or crumbly bits, pips, seeds, fibrous parts of fruit, husks, or bones. No restriction on particle size.
Light Diet	Appendix 3	Excludes foods that are fatty, strongly flavoured and gas-forming; can be used post-surgery/medical procedure, with nausea and as a transition from liquid to normal diet.






Therapeutic Diet and/or Fluids Procedure

Low Bacteria	Appendix 4	Avoids food with high bacterial load; for patients with neutropenia
Low Fat	P 164	Low in fat (especially saturated fat) less than 45g/day, and low in sugar
Low Fibre	P 57	Eliminates foods that are high in fibre.
Low Phosphate	Appendix 5	Diet avoids/limits foods high in phosphate (milk, cheese, processed meat, etc.) to < 1200mg/day
Low Potassium	P 150 Appendix 6	Diet avoids/limits foods high in potassium (milk, meat, some fruit/vegetables etc.) to < 70mmol/day
Low Residue	Appendix 7	Eliminates food that provides significant amounts of dietary fibre or that produce residue in the intestine.
Low Sodium	P 188 Appendix 8	Diet to reduce sodium intake including avoiding high sodium foods & adding salt to food in cooking & on serving
Nil By Mouth		No food orally; not safe/medically able to eat.
Nourishing Fluids	P 35	Includes fluids that also contain fat- all clear fluids plus all nutrition supplements & milk-based drinks.
Renal Diet	Appendix 9	Used for patients with renal failure that are pre-dialysis (↓Na, ↓K ↓PO4)
Renal Haemodialysis	P 134 Appendix 10	Used for patients who have haemodialysis. No soup unless specified by the dietitian. (Na, ↓K, ↓PO4)
Haemodialysis Unit	Appendix 11	Used for non-admitted patients who are having haemodialysis at AHS. Includes light meals and snacks only.
Vegetarian	P 22 & 26	No meat provided; need to specify if patient is vegan (no eggs or dairy also)
Weight Management	P 100	Planned to achieve weight loss by reducing the patient's total energy intake; can include use of Optifast

Appendix 2: Safe Swallow Chart

Safe Swallowing Chart

Name: [Click here to enter text.](#)**Date:** 13/01/2023**Ward:** Choose an item.

	Choose an item.		Choose an item.
	Choose an item.		
	Choose an item. Choose an item.		
	Choose an item. Choose an item.		Choose an item.

Standard safe swallow strategies:

	Must be fully awake & alert		Minimise distractions during mealtime
	Must be sitting fully upright		Stay upright for at least 20 mins after meals

For this patient:

	Make sure each mouthful is swallowed before giving the next		Remove any pocketed food from left/right cheek after meals
	Alternate food & drink during meals		Ensure small mouthfuls
	Regular oral care		Full oral care after meals
	May have small sips of thin drinks with supervision		Patient allowed Level 7 Regular snacks with supervision

Rehabilitation Strategies

	Double swallows for each mouthful		Keep chin down when swallowing
	Effortful swallow ("Swallow hard, with all your muscles")		Cough, clear & swallow every 4-5 mouthfuls
	Place food in the right side of the mouth		Place food in the left side of the mouth

**Stop feeding if coughing or tired.**

If necessary, please refer to speech pathology or administer the Dysphagia Screening Tool (DST).

Speech Pathologist: [Click here to enter text.](#) Contact Number: [Click here to enter text.](#)

Appendix 3: Light Diet

Aim:

- To provide a diet containing foods lower in fat and less gas-forming foods and spices.

Characteristics:

- Physiologically this diet has no advantages over the standard full diet, but traditionally it has been used for patients who cannot tolerate the full diet and want mildly flavoured, easily digested food. This diet does not have to be soft, bland or low fibre. Small meals selected from a normal low-fat menu may be a suitable alternative.

Indications:

- post-surgery / medical procedures
- nausea

Precautions:

- Individual likes and dislikes need to be considered when planning this diet.

Food Choices Chart

Foods to Choose	Foods to Avoid
Breads & Cereals	
All other cereals & all breads All other sandwiches Plain rice & pasta Plain cake All biscuits Canned spaghetti	Bran-based breakfast cereals Pickles, chutney, gherkin
Fruits	
All other fruits	Rhubarb, prunes, dates, pineapple
Vegetables	
All other vegetables Starchy vegetables (boiled, mashed, roasted potato)	Corn, broccoli, brussel sprouts, cauliflower, cabbage, capsicum Salad & salad dressings Fried potato chips or wedges
Milk/Milk Products	
All other milk/milk products	Cream
Meat & Meat Alternatives	
All other meat/meat alternative dishes Egg dishes	Strongly spiced dishes (e.g. Curries, satays) High fat dishes >15g fat per serve Bacon, sausages, baked beans Dishes containing nuts or legumes
Miscellaneous	

Foods to Choose	Foods to Avoid
All other soups All other desserts All beverages	Minestrone, lentil, broccoli, potato & leek soups Any desserts >10g fat per serve (e.g. cheesecake, pie, pastries)

Appendix 4: Low Bacteria Diet

Aim:

- To provide a diet limiting foods at high risk of carrying food-borne disease (e.g. Salmonella, E. coli, Listeria monocytogenes).

Characteristics:

- Avoids food with a high bacterial or fungal load. Sterile diets are no longer used.

Indications:

- Patients with neutropenia (neutrophil count $\leq 1000 \text{ cell}/\mu\text{L}$)
- Severely immunosuppressed patients (e.g. post-transplant; undergoing some cancer treatments)

Precautions:

- All salads and sandwiches served to patients on low bacteria diet must be prepared on the same day. Purchased pre-prepared salads and sandwiches are not allowed. These patients are often malnourished and may require nutritional supplements as per dietitian assessment. Raw fruit and vegetables are to be sanitised (washed or soaked for the appropriate amount of time) using validated chemical preparations, e.g. sodium hyperchlorite solution or electrolysed water.

Notes:

- Nutritionally adequate. Due to improvements in food safety, the standard hospital diet may be considered safe for most immune-compromised patients.

Food Choices Chart

Foods to Choose	Foods to Avoid
Breads & Cereals	
All breads, processed ready-to-eat Cooked breakfast cereals (e.g. porridge, toasted muesli) All other breakfast cereals Cooked rice & pasta Plain cake All biscuits Canned spaghetti	Raw oats and muesli
Fruits	
Sanitised fresh fruit Portion controlled packs of fruit	Fresh fruit with a rough texture (e.g. raspberries, rockmelon)
Vegetables	
All cooked vegetables and starchy vegetables Sanitised vegetables only	Unwashed raw vegetables including leafy green vegetables (e.g. lettuce, raw sprouts) & mushrooms Pre-prepared or pre-packaged salads
Milk/Milk Products	
All pasteurised milk & milk products	All yoghurt

Foods to Choose	Foods to Avoid
Commercially packaged mild cheese (e.g. cheddar) Cottage cheese if served on same day as package opened	Soft, semi-soft and surface-ripened cheese (e.g. brie, camembert, feta, blue)
Meat & Meat Alternatives	
All well-cooked meat, poultry, fish and boiled eggs Canned fish Pasteurised meats (cooked in bag) and served on day of opening	Undercooked meat, poultry, fish or eggs Raw tofu, smoked meat & fish, tempeh Sliced process meats (e.g. ham, chicken roll) Pates & meat spreads Peeled, ready-to-eat prawns
Miscellaneous	
All cooked sauces and gravies Portion packs commercial mayonnaise and dressings Pasteurised fruit juices in sealed portions Tea, coffee, soft drink All hot soups Jam, peanut butter, Vegemite	Salad dressings made with blue cheese or raw eggs Fresh and unpasteurised juices Herbal teas Cold soups, miso soup Pates & honey
Desserts	
Hot desserts Heat-treated portion-controlled cold desserts Refrigerated, commercial and homemade cakes, pastries and puddings Fresh, pasteurised custard Normal frozen ice-cream	Unrefrigerated cream-filled pastries Commercial products containing cream or cold custard Soft serve ice-cream

Appendix 5: Low Phosphate Diet

Aim:

- To limit phosphate intake to 1200mg per day.

Characteristics:

- Limit foods high in phosphate.

Indications:

- Renal failure
- Hyperphosphataemia
- Secondary hyperparathyroidism
- Chronic kidney disease

Precautions:

- Meat should be limited to two serves per day. Milk products should be limited to two serves per day.
- Phosphate binders are usually used in conjunction with a low-phosphate diet. In some cases a low-phosphate diet is needed together with a high-protein diet, and the dietitian may then prescribe higher levels of protein foods than normally allowed on this diet.

Notes:

- Nutritionally adequate.

Food Choices Chart

Foods to Choose	Foods to Avoid
Breads & Cereals	
All other breads Cornflakes Porridge made with water Special K Weetbix White rice Couscous Cornflour Sao crackers Fresh & dried white pasta White plain flour	Fruit bread Porridge made with milk Muesli Brown rice Barley Raw oats Wholemeal pasta
Fruits	
All fresh and canned fruit	Dried fruit
Vegetables	
All others	Mixed beans Vegetables served with a cheese or white sauce
Milk/Milk Products	
ONLY 2 SERVES PER DAY#: Milk for cereal and hot drinks Cream	All milk (including soy), other than for cereal and hot drinks Semi-hard and hard cheeses

Foods to Choose	Foods to Avoid
Sour cream Cream cheese Ricotta Cottage cheese Ice-cream Fruché Yoghurt – Limit to 100g tub	Condensed milk Evaporated milk Milk powder Yoghurt – 200g tub Custard Creamed rice
Meat & Meat Alternatives	
ONLY 2 SERVES PER DAY+: Beef Lamb Chicken Turkey Pork Fish (not crumbed) Deli meats Egg Tofu	Other fish Sardines Bacon Corned beef Sausages Crumbed fish Legumes Baked beans
Miscellaneous	
Plain biscuits Milk Arrowroot biscuits Milk Coffee biscuits Cracker biscuits Shortbread Scotch fingers Jam filled biscuit Cream biscuit Coffee Tea	Cola flavoured drinks Chocolate Chocolate cake Fruit cake Oatmeal or chocolate biscuits Cheese and white sauces Cream soups made with milk Soups with legumes e.g. minestrone Peanut butter Vegemite Nuts Cocoa based drinks e.g. Milo

Desserts

1 serve dairy = ½ cup milk, rice milk, creamed rice
 30 g cheese
 100 g yoghurt
 + 1 serve meat = 90 g meat, chicken
 100 g fish
 2 eggs

Appendix 6: Low Potassium Diet

Aim:

- To limit potassium intake to 70mmol (2730mg) per day.

Characteristics:

- Limits foods high in potassium.

Indications:

- Hyperkalaemia
- Patients on haemodialysis
- End stage renal disease

Precautions:

- Fruit is limited to 2 serves per day and vegetables limited to 1 or 4 serves per day depending on the type.

Notes:

This diet may not be nutritionally adequate. Nutrients that may require additional supplementation include fibre, magnesium, calcium and zinc.

Food Choices Chart

Foods to Choose	Foods to Avoid
Breads & Cereals	
All breads Cornflakes Special K Weetbix Porridge Fresh & dried white pasta White Rice Couscous Noodles Barley Sao crackers	Fruit Bread Cereals containing dried fruit Wholemeal pasta Brown rice
Fruit	
Fruit juice – 125mL/day maximum ONLY 2 SERVES PER DAY °: Apple Pear Strawberries Blackberries Blueberries Raspberries Boysenberries Tinned and drained:	All dried fruit Apricot Nectarine Peach Cherries Grapefruit Grapes Pineapple Kiwifruit Banana Honeydew melon

Foods to Choose	Foods to Avoid
Apple Pear Peach Pineapple Mandarin Plums Paw paw Papaya	Orange Lemon Plum Mango Watermelon Rockmelon Rhubarb Avocado
Vegetables	
ONLY 4 SERVES PER DAY^: Cabbage Cauliflower Carrot Green beans Bean sprouts Snow peas Celery Capsicum Cucumber Lettuce Peas Zucchini Mushroom Spinach - raw Garlic Spring onion Radish Turnip Swede Eggplant Fresh-made unsalted soups – if made from vegetables in the above list ONLY 1 SERVE / DAY^: Potato – boiled Pumpkin – boiled Sweet potato - boiled Asparagus Broad beans Beetroot	Potato – other than boiled Potato chips Pumpkin – other than boiled Sweet potato – other than boiled Tomato Broccoli Silverbeet Avocado Dried beans Kidney beans Baked beans Canned soup

Foods to Choose	Foods to Avoid
Sweetcorn Leek Parsnip Brussels Sprouts Onion Lentils Chickpeas Four bean mix	
Milk/Milk Products	
150ml milk per day for cereal, hot drinks or as a beverage only (e.g. cow or dairy alternatives) All cheese Cream cheese Sour cream Cream Creamed rice Ice-cream Custard Fruché Plain or vanilla yoghurt – 100g tub only	All milk above allowance Condensed milk Evaporated milk Milk powder Soy milk Yoghurt – 200g tub
Meat & Meat Alternatives	
Chicken Beef Pork Tuna – tinned Salmon – tinned Ham Egg Salami Tofu	Fish - except tinned Sardines Turkey Lamb Bacon
Miscellaneous	
Macadamia nuts Sunflower seeds Sesame seeds Soy sauce Oyster sauce BBQ Sauce Curry powder	Almonds Cashews Peanuts Vegemite Worcestershire sauce Coffee - ground Iced coffee

Foods to Choose	Foods to Avoid
Brown sugar Golden syrup Vegetable improver Gravy Fats Tomato sauce Coffee – instant Milo Drinking chocolate Shortbread Scotch fingers Sponge cake Jam filled biscuit Cream biscuit Swiss roll Carrot cake Oatmeal biscuit	Liquorice Fruit & nut biscuit Cheesecake Chocolate Desiccated coconut Creamed rice Chocolate cake Fruit cake Dried Fruits Peanut butter Vegemite Baked beans

- To reduce the potassium content of vegetables, boil in a large volume of water. Do not roast, bake, steam or microwave vegetables

^1 serve vegetables = 1 cup raw/salad vegetables
 ½ cup cooked vegetables

° 1 serve fruit = 1 medium piece
 ½ cup diced
 125 ml fruit juice

Appendix 7: Low Residue Diet

Aim:

- To provide a diet low in dietary fibre.

Characteristics:

- A diet providing less than 10g of dietary fibre per day by avoiding foods that are high in dietary fibre and resistant starch and cause residue in the bowel.

Indications:

- Preparation for colonoscopy; specific gastrointestinal disorders (e.g. fistulae, stenosis)
- Acute phases of inflammatory bowel disease and diverticulitis

Precautions:

- A dietitian should be consulted if this diet is followed for more than three days.

Notes:

- Not nutritionally adequate. Nutrients that may be inadequate include fibre, folate and magnesium.

Food Choices Chart

Foods to Choose	Foods to Avoid
Breads & Cereals	
White bread & bread rolls Breakfast cereals providing <1.5g fibre per serve (e.g. Cornflakes, Rice bubbles, Special K) White rice, pasta & noodles Plain dry, white crackers All other biscuits (e.g. Milk Arrowroot, Milk Coffee)	Wholemeal, wholegrain, Hi-fibre white or rye bread Other cereals (Weetbix, rolled oats, porridge, muesli, bran cereals) Brown rice or wholemeal pasta; canned spaghetti Biscuits made with wholemeal flour, fruit, nuts, coconut or bran
Fruits	
One serve only per day of canned fruit <2.5g fibre per serve (e.g. peaches, apples, two fruits)	All fresh and dried fruit Canned pineapple, pears, apricots, plums, fruit salad, cherries, berries, prunes
Vegetables	
Boiled or mashed potato <1.5g fibre per serve (e.g. well-cooked cauliflower, pumpkin, zucchini, carrots)	Jacket and roasted potato Salads All other vegetables including peas, tomato, mushrooms, pickles
Milk/Milk Products	
Plain yoghurt All milk and cheese – limit to two serves per day	Fruit yoghurt
Meat & Meat Alternatives	
Plain roasted and grilled meats, poultry and fish Eggs & plain omelettes	All casseroles & stir-fry's Sausages & bacon All beans & lentils including baked beans
Miscellaneous	
All sauces, mayonnaise Strained fruit juices (e.g. apple) Tea, coffee, soft drink, cordial, Milo Clear broths, chicken noodle soup, strained cream soups	All salad dressings Prune and tomato juice All other soups

Foods to Choose	Foods to Avoid
Honey, Vegemite	Jam, marmalade, peanut butter Unprocessed bran, nuts, seeds
Desserts	
Plain milk puddings (e.g. custards, creamy rice) Ice-cream, jelly Plain cakes and puddings	Any desserts containing fruit (e.g. sultanas) Cakes made with wholemeal flour, fruit, nuts, coconut or bran

Appendix 8: Low Sodium Diet

Aim:

- To limit total sodium intake to 100mmol (2300mg) per day.

Characteristics:

- Limit foods high in sodium.

Indications:

- Hypertension
- Congestive heart failure
- Acute or chronic renal failure
- Liver disease with ascites

Precautions:

- No salt sachets are provided on meal trays.
- If more than four slices of bread are served per day, a lower-sodium bread may be needed to meet the daily restriction.

Notes:

- Nutritionally adequate.

Food Choices Chart

Foods to Choose	Foods to Avoid
Breads & Cereals	
Special K Weetbix Porridge Natural muesli Bread Fruit bread Fresh & dried pasta Rice Couscous Plain flour Noodles Barley Sao crackers	Cornflakes Wraps Tinned spaghetti Instant rice, pasta, noodles
Fruit	
All other	Olives
Vegetables	
Fresh Frozen Legumes (lentil, chickpeas, soup mix, soybeans, kidney beans)	Baked beans / salt reduced baked beans Pickled & canned vegetables Commercial tomato / vegetable juice

Foods to Choose	Foods to Avoid
'No added salt' canned vegetables 'No added salt' baked beans	Vegetables served with white sauce or added salt
Milk/Milk Products	
Milk – all types Yoghurt Custard Dairy desserts Ice-cream Ricotta Cottage cheese Swiss cheese Fruché	All other semi-hard and hard cheeses – cheddar, brie, camembert, feta, parmesan Cream cheese Cheese spreads
Meat & Meat Alternatives	
Fresh, unprocessed meat & fish 'No added salt' tinned fish e.g. in spring water Egg Tofu	Processed meats e.g. deli meats, bacon Tinned fish in brine / flavouring, sausages, smoked fish, corned beef
Miscellaneous	
As desired – unsalted varieties only Unsalted butter Salt reduced margarine e.g. Meadow Lea Oil Mint sauce – no added salt Apple sauce – no added salt Home-made unsalted soups Jam Honey Plain biscuits Milk Arrowroot biscuits Milk Coffee biscuits	Salted butter Margarine Most sauces e.g. Soy sauce Tomato sauce BBQ sauce Worcestershire sauce Teriyaki sauce Tartare sauce Vegetable improver White or cheese sauce Gravy Stock cubes Meat & fish paste Bonox Soup Mustard Pickles Relish Vegemite Peanut butter

Foods to Choose	Foods to Avoid
	Salty crackers e.g. Jatz® crackers Salted nuts

- No added salt/stock or salt substitutes to cooking – beware of commercial spice mixes that often contain a high salt content.
- Do not provide salt sachets on meal tray.

Appendix 9: Renal Diet

Aim:

- To limit intake of sodium, potassium, and phosphate.

Characteristics:

- Limits foods high in sodium, potassium, and phosphate.

Indications:

- End stage renal disease
- Acute or chronic kidney disease

Precautions:

- No salt sachets are provided on meal trays.
- If more than four slices of bread are served per day, a lower-sodium bread may be needed to meet the daily restriction.
- Meat, fruit and milk products should be limited to two serves per day. Vegetables are limited to 1 or 4 serves per day depending on the type.
- The dietitian may prescribe higher levels of protein foods than normally allowed on this diet.

Notes:

- This diet may not be nutritionally adequate. Nutrients that may require additional supplementation include fibre, magnesium, calcium and zinc.

Food Choices Chart

Foods to Choose	Foods to Avoid
Breads & Cereals	
All other bread Special K Weetbix Porridge made with water Natural muesli Fresh & dried white pasta White Rice Couscous Noodles Barley Sao crackers	Wholegrain bread Fruit bread Cereals containing dried fruit Wraps Porridge made with milk Cornflakes Tinned spaghetti Instant rice, pasta, noodles Wholemeal pasta Brown rice
Fruit	
Juice – 125mL/day maximum ONLY 2 SERVES PER DAY °: Apple Pear	Orange Lemon Plum Mango Watermelon

Foods to Choose	Foods to Avoid
Berries – strawberries, blackberries, blueberries, raspberries Tinned and drained: Apple Pear Peach Pineapple Mandarin Plums Paw paw Papaya	Olives All fresh stone fruit e.g. Apricot Nectarine Peach Grapefruit Grapes Pineapple Kiwifruit Banana Honeydew melon Rockmelon Rhubarb Avocado Dried fruit
Vegetables	
ONLY 4 SERVES PER DAY^: Cabbage Cauliflower Carrot Green beans Bean sprouts Snow peas Celery Capsicum Cucumber Lettuce Peas Zucchini Mushroom Garlic Spring onion Radish Turnip Swede Eggplant Canned vegetables as per above list – must be ‘no added salt’ varieties Home-made unsalted soups – if made from vegetables in the above list	Baked beans / salt reduced baked beans Pickled & canned vegetables Commercial tomato / vegetable juice Tomato paste Potato – other than boiled Potato chips Pumpkin – other than boiled Sweet potato – other than boiled Tomato Broccoli Silverbeet Spinach Avocado Dried beans Kidney beans Baked beans Mixed beans Canned soup Vegetables in a cheese or white sauce

Foods to Choose	Foods to Avoid
ONLY 1 SERVE PER DAY^: Potato – boiled Pumpkin – boiled Sweet potato - boiled Asparagus Broad beans Beetroot Sweetcorn Leek Parsnip Brussels Sprouts Onion Tomato Lentils Chickpeas	
Milk/Milk Products	
ONLY 2 SERVES PER DAY#: 150 ml milk per day for cereal and hot drinks or as a beverage (e.g. cow or dairy alternatives) Custard Dairy desserts Ice-cream Cottage cheese Ricotta Sour cream Cream Creamed rice Fruché Yoghurt – Limit to 100g tub	All milk above allowance All other semi-hard and hard cheeses – cheddar, gouda, brie, camembert, feta, parmesan Cream cheese Cheese spreads Condensed milk Evaporated milk Milk powder Yoghurt – 200g tub
Meat & Meat Alternatives	
ONLY 2 SERVES PER DAY +: Chicken Beef Pork Tuna – tinned ‘No added salt’ tinned fish e.g. in spring water Salmon – tinned ‘No added salt’ tinned fish e.g. in spring water Egg Tofu	Lamb Turkey Sardines Processed meats e.g. deli meats, bacon, sausages Tinned fish in brine / flavouring Fish – except tinned Corned beef Baked beans Legumes

Foods to Choose	Foods to Avoid
Miscellaneous	
Macadamia nuts Sunflower seeds Sesame seeds Unsalted butter Salt reduced margarine e.g. Meadow Lea Oil Mint sauce – no added salt Apple sauce – no added salt Jam Honey Marmalade Lemon spread Sugar Golden syrup Herbs & spices Curry powder Pepper Vinegar Garlic Lemon juice Chilli Plain biscuits Milk Arrowroot biscuits Milk Coffee biscuits Coffee – instant Tea Shortbread Scotch fingers Sponge cake Jam filled biscuit Cream biscuit Swiss roll Carrot cake Oatmeal biscuit	Salted butter Margarine Almonds Cashews Peanuts Salted nuts Salted butter Margarine Most sauces e.g. Soy sauce Tomato sauce BBQ sauce Worcestershire sauce Teriyaki sauce Tartare sauce Vegetable improver Gravy Tomato paste Stock cubes Meat & fish paste White or cheese sauce Bonox Soup Mustard Pickles, chutney Relish Peanut butter Vegemite Worcestershire sauce Salty crackers e.g. Jatz crackers Coffee - ground Iced coffee Liquorice Fruit & nut biscuit Cheesecake Chocolate Desiccated coconut Creamed rice Chocolate cake

Foods to Choose	Foods to Avoid
	Fruit cake Cola or Cocoa based drinks e.g. Milo

- To reduce the potassium content of vegetables, boil in a large volume of water. Do not roast, bake, steam or microwave vegetables
 - ^1 serve vegetables = 1 cup raw/salad vegetables
½ cup cooked vegetables
 - ° 1 serve fruit = 1 medium piece
½ cup diced
125 ml fruit juice
 - # 1 serve dairy = ½ cup milk, rice milk, creamed rice
30 g cheese
100 g yoghurt
 - + 1 serve meat = 90 g meat, chicken
100 g fish
2 eggs

Appendix 10: Haemodialysis Diet

Aim:

- To provide a diet higher in energy and protein and restricted in sodium, potassium, and phosphate.

Characteristics:

- Limits foods high in sodium, potassium and phosphate whilst providing high biological value protein choices.

Indications:

- Patients on haemodialysis

Precautions:

- Not all patients on haemodialysis require this diet specification. For example, a low-potassium diet may not be required; nor will it be needed if a patient is eating poorly.
- Monitoring by a dietitian is required if this diet specification is to be removed, combined with other diet specifications, or used long term.

Notes:

- This diet may not meet individual nutrient requirements due to the individual needs and treatment of this patient group. Supplementation may be required.

Food Choices Chart

Foods to Choose	Foods to Avoid
Breads & Cereals	
White bread Rice Cornflour Couscous Sao crackers® Cornflakes Special K Weetbix Porridge made with water Fresh & dried white pasta	Bread – wholegrain and wholemeal Fruit bread Wraps Cereals containing dried fruit Bran-based cereals e.g. All-Bran Porridge made with milk Natural muesli Tinned spaghetti Instant rice, pasta, noodles Barley Wholemeal pasta Brown rice
Fruit	
ONLY 2 SERVES PER DAY °: Apple Pear Berries – strawberries, blackberries, blueberries, raspberries Tinned: Apple Pear Peach Pineapple	Orange Lemon Plum Mango Watermelon Juice – 125mL/day maximum Olives All fresh stone fruit e.g.

Foods to Choose	Foods to Avoid
Mandarin Plums Paw paw Papaya	Apricot Nectarine Peach Grapefruit Grapes Pineapple Kiwifruit Banana Honeydew melon Rockmelon Rhubarb Avocado All dried fruit
Vegetables	
ONLY 4 SERVES PER DAY^: Cabbage Cauliflower Carrot Green beans Bean sprouts Snow peas Celery Capsicum Cucumber Lettuce Peas Mushroom Zucchini Garlic Spring onion Radish Turnip Swede Eggplant Canned vegetables as per above list – must be ‘no added salt’ varieties Home-made unsalted soups – if made from vegetables in the above list ONLY 1 SERVE / DAY^: Potato – boiled	Baked beans / salt reduced baked beans Pickled & canned vegetables Commercial tomato / vegetable juice Tomato paste Potato – other than boiled Hot chips Pumpkin – other than boiled Sweet potato – other than boiled Broccoli Silverbeet Spinach Avocado Dried beans Kidney beans Baked beans Bean mix Canned soup Any vegetables served with cheese or white sauce

Foods to Choose	Foods to Avoid
Pumpkin – boiled Sweet potato - boiled Asparagus Broad beans Beetroot Sweetcorn Leek Parsnip Brussels Sprouts Onion Tomato Lentils Chickpeas	
Milk/Milk Products	
<250ml per day milk for cereal and hot drinks (e.g. cow or dairy alternatives) 1 slice cheese per day Hard cheese allowed in hot mains and sauces Cream Sour cream Creamed rice Ricotta Cottage cheese Ice-cream Fruché Yoghurt – Limit to 100g tub per day	All milk above allowance Hard cheese above limits Cream cheese Condensed milk Evaporated milk Milk powder Soy milk Yoghurt – 200g tub Custard
Meat & Meat Alternatives	
Beef Lamb Chicken Turkey Pork Tuna & Salmon – no added salt varieties Egg Tofu	Processed meats e.g. deli meats, bacon Fish – except tinned in brine/flavouring Sardines Legumes and lentils (unless vegetarian)
Miscellaneous	
Macadamia nuts Sunflower seeds	Almonds Cashews

Foods to Choose	Foods to Avoid
Sesame seeds Unsalted butter Salt reduced margarine e.g. Meadow Lea Oil Mint sauce – no added salt Apple sauce – no added salt Jam Honey Marmalade Lemon spread Sugar Golden syrup Herbs & spices Curry powder Pepper Vinegar Garlic Lemon juice Chilli Pickles, chutney (small amounts only) Plain biscuits Milk Arrowroot biscuits Milk Coffee biscuits Coffee – instant Shortbread Scotch fingers Jam filled biscuit Cream biscuit	Peanuts Salted butter Margarine Most sauces e.g. Soy sauce Tomato sauce BBQ sauce Worcestershire sauce Teriyaki sauce Tartare sauce Vegetable improver Gravy Tomato paste Stock cubes Meat & fish paste Bonox, Bovril Soup Coconut – cream, milk, desiccated etc. Mustard Relish Vegemite Peanut butter Vegemite / Marmite Worcestershire sauce Salty crackers e.g. Jatz crackers Coffee - ground Iced coffee Liquorice Fruit & nut biscuit Cake – all varieties Chocolate Desiccated coconut Creamed rice Oatmeal biscuit Dried fruit Milo Drinking chocolate

- To reduce the potassium content of vegetables, boil in a large volume of water. Do not roast, bake, steam or microwave vegetables

^1 serve vegetables = 1 cup raw/salad vegetables
½ cup cooked vegetables

° 1 serve fruit = 1 medium piece
½ cup diced
125 ml fruit juice

Appendix 11: Haemodialysis Unit Light Meals and Snack Menu

- Patients booked for morning sessions can choose high protein porridge or a sandwich.
- For all other sessions times patients will receive a sandwich only.
- Available sandwich fillings include: chicken, beef, cream cheese, tuna or eggs. This will ensure patients on haemodialysis receive sandwiches that are in line with the Australian Guide to Healthy Eating and evidence-based guidelines for chronic renal failure.
- Snacks include savoury crackers with cream cheese.

Food Choices Chart

Morning Dialysis Session
High protein porridge Or sandwich (with or without salad, white or wholemeal bread) Chicken Beef Cream cheese Tuna Egg
All Other Sessions
Sandwich (with or without salad, white or wholemeal bread) Chicken Beef Cream cheese Tuna Egg
Available Snacks
Savoury crackers with cream cheese Fresh fruit Stewed fruit