



# Single Room Prioritisation and Cohorting of Inpatients on Transmission-based Precautions Procedure

| Site                   | Operational Area    | Applicable to       |
|------------------------|---------------------|---------------------|
| Armada Kalamunda Group | All inpatient areas | Medical and Nursing |

## 1. Purpose

This document is intended to provide clinical guidance for nursing and medical staff of appropriate utilisation of single rooms and bed placement for inpatients on transmission-based precautions. This procedure outlines the decision-making for cohorting of inpatients on transmission-based precautions (suspected or confirmed cases) in a shared room or cohort care area (multi-bedded room/bay/ward/unit), in instances where single rooms are unavailable, and in the context of a local outbreak or pandemic.

The formation of this procedure is in accordance with the current [national](#) and [state](#) guidelines, and used in conjunction with relevant [AKG Infection Prevention and Management guideline and policy documents](#).

## 2. Prioritisation for single-room and negative-pressure isolation room (NPIR) placement

- Inpatients requiring transmission-based precautions, those with behavioural/safety concerns, and palliative care must be given priority for a single room placement.
- The use of negative pressure isolation room must be prioritised to inpatients with highly transmissible airborne infectious illness.
- The decision for prioritising single room allocations should be individually assessed with consideration for any foreseeable risks in relation to non-placement in a single room, changes to patient's risk status and surge in demand for single rooms.

| Priority | Infectious Disease       | Type of Precautions  | Placement and Cohorting Guidelines                                                                                                                                                                  |
|----------|--------------------------|----------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 1        | Viral Haemorrhagic Fever | Contact and Airborne | <ul style="list-style-type: none"><li>▪ <b>NPIR (Negative Pressure Isolation Room).</b></li><li>▪ <b>Do not Cohort.</b></li><li>▪ Referral and transfer to appropriate tertiary facility.</li></ul> |
|          | Pulmonary Tuberculosis   | Airborne             | <ul style="list-style-type: none"><li>▪ NPIR.</li><li>▪ <b>Do not Cohort.</b></li><li>▪ Referral and transfer to appropriate tertiary facility if no available NPIR .</li></ul>                     |
|          | Chicken pox (Varicella)  | Contact and Airborne | <ul style="list-style-type: none"><li>▪ NPIR.</li><li>▪ <b>Do not Cohort.</b></li><li>▪ Referral and transfer to appropriate tertiary facility if no available NPIR.</li></ul>                      |

Before referencing this procedure document please ensure that you have the latest version from the [Policy Library](#) information hub page.

# Single Room Prioritisation and Cohorting of Inpatients on Transmission-based Precautions Procedure

|          |                                                                                                                                                                                                                                                             |                                                                            |                                                                                                                                                                                                                                                                                                                                    |
|----------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|          |                                                                                                                                                                                                                                                             |                                                                            | If transfer not feasible, place in single room with door closed and place an <b>air purifier unit (APU) inside room.</b>                                                                                                                                                                                                           |
|          | <ul style="list-style-type: none"> <li>SARS</li> <li>MERS CoV</li> <li>Pandemic influenza</li> </ul>                                                                                                                                                        | <b>Contact and Airborne</b>                                                | <ul style="list-style-type: none"> <li>NPIR</li> <li><b>Do not Cohort</b></li> <li>Referral and transfer to appropriate tertiary facility if no available NPIR.</li> <li>If transfer not feasible, place in single room with door closed and place an <b>air purifier unit (APU) inside room</b></li> </ul>                        |
|          | Measles                                                                                                                                                                                                                                                     |                                                                            |                                                                                                                                                                                                                                                                                                                                    |
|          | Disseminated HZV (shingles)<br><i>(rash in multiple sites of the body)</i>                                                                                                                                                                                  |                                                                            |                                                                                                                                                                                                                                                                                                                                    |
|          | <ul style="list-style-type: none"> <li>Meningococcal disease</li> <li>Mumps</li> <li>Undiagnosed rash/fever</li> <li>Parvovirus B19</li> </ul>                                                                                                              | <b>Droplet</b>                                                             | <ul style="list-style-type: none"> <li>Single room with ensuite or dedicated bathroom</li> <li><b>Do not cohort</b></li> </ul>                                                                                                                                                                                                     |
|          | Invasive group A streptococcus disease (iGAS)                                                                                                                                                                                                               | <b>Contact</b><br>(Add DROPLET precautions if with respiratory symptoms)   | <ul style="list-style-type: none"> <li>Single room with ensuite or dedicated bathroom</li> <li><b>Do not cohort</b></li> </ul>                                                                                                                                                                                                     |
| <b>2</b> | <ul style="list-style-type: none"> <li>Micro- C (Healthcare Associated MRSA)</li> <li>Micro- G (CRE/CPO)</li> </ul>                                                                                                                                         | <b>Contact</b>                                                             | <ul style="list-style-type: none"> <li>Single room with ensuite or dedicated bathroom.</li> <li><b>Do not cohort.</b></li> <li>Contact IPM if no single room available</li> </ul>                                                                                                                                                  |
|          | <ul style="list-style-type: none"> <li>SARS-CoV-2 (COVID-19), including symptomatic and asymptomatic close contacts.</li> <li>Influenza/Influenza-like illness</li> <li>Other confirmed respiratory viruses</li> <li>Suspected respiratory virus</li> </ul> | <b>Droplet</b><br>(Add contact precautions if with respiratory secretions) | <ul style="list-style-type: none"> <li>NPIR/Single Room with en suite or dedicated bathroom</li> <li>For single room – place air purifier unit inside room.</li> <li>Do not cohort if laboratory result is pending.</li> <li>Only cohort when single rooms exhausted.</li> <li>Only cohort if the viruses are the same.</li> </ul> |
|          | <ul style="list-style-type: none"> <li>Respiratory Syncytial Virus (RSV)</li> <li>Rhinovirus</li> </ul>                                                                                                                                                     | <b>Contact and Droplet</b>                                                 |                                                                                                                                                                                                                                                                                                                                    |
|          | <ul style="list-style-type: none"> <li>Micro-Alert B – MRSA B (<b>Community associated</b>)</li> <li>Micro-Alert J – <i>C. auris</i></li> <li>Micro-Alert K – <i>C. auris</i> (<i>Contact</i>)</li> <li>Micro-Alert V - VRE</li> </ul>                      | <b>Contact</b>                                                             | <ul style="list-style-type: none"> <li>Single room with en suite or dedicated bathroom</li> <li>Cohort room with en suite or dedicated bathroom</li> <li><b>Cohort patients with the same micro-alert and with no other risk factors</b></li> </ul>                                                                                |
|          | <i>Clostridioides difficile</i> or infective gastroenteritis                                                                                                                                                                                                | <b>Contact</b>                                                             | <ul style="list-style-type: none"> <li>Single room with ensuite or dedicated bathroom</li> <li>Cohort room with ensuite</li> </ul>                                                                                                                                                                                                 |

|   |                                                                                                                                                                |                                                                                                                                 |                                                                                                                                                                                                                                                                                                                 |
|---|----------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|   |                                                                                                                                                                |                                                                                                                                 | <ul style="list-style-type: none"> <li>▪ <b>Cohort patients with same confirmed organism and no other risk factors</b></li> </ul>                                                                                                                                                                               |
| 3 | Shingles (Localised) (where the lesions cannot be covered)                                                                                                     | Contact                                                                                                                         | <ul style="list-style-type: none"> <li>▪ Single room with ensuite or dedicated bathroom</li> <li>▪ Cohort room with ensuite or with dedicated bathroom</li> <li>▪ Cohort only if no other risk factors and infections</li> </ul>                                                                                |
|   | Hospitalised <u>Outside Australia</u> (direct hospital transfer)                                                                                               | Contact Precaution                                                                                                              | <ul style="list-style-type: none"> <li>▪ Single room with ensuite or dedicated bathroom</li> <li>▪ Cohort room with ensuite or with dedicated bathroom</li> <li>▪ <b>Contact Precautions until screening results are negative</b></li> </ul>                                                                    |
|   | Hospitalised <u>outside Western Australia</u> within the last 12 months                                                                                        | Standard                                                                                                                        | <ul style="list-style-type: none"> <li>▪ Single room with ensuite or dedicated bathroom</li> <li>▪ Cohort room with ensuite or with dedicated bathroom</li> </ul>                                                                                                                                               |
|   | Multi-resistant Gram-Negative Bacteria (ESBL)                                                                                                                  | <b>Standard</b><br>(Contact precautions if with risk factors -e.g., incontinence, indwelling devices, full assistance required) | <ul style="list-style-type: none"> <li>▪ Single room with ensuite or dedicated bathroom</li> <li>▪ Cohort room with ensuite or with dedicated bathroom</li> </ul>                                                                                                                                               |
|   | <ul style="list-style-type: none"> <li>▪ Micro-Alert F – VRE contact</li> <li>▪ Micro-Alert H – CPO contact</li> <li>▪ Micro-Alert W – MRSA contact</li> </ul> | Standard                                                                                                                        | <ul style="list-style-type: none"> <li>▪ Single room with ensuite or dedicated bathroom</li> <li>▪ Cohort room with ensuite or dedicated bathroom               <ul style="list-style-type: none"> <li>• <b>Cohort patients with the same micro-alert and with no other risk factors</b></li> </ul> </li> </ul> |

### 3. Cohorting

#### 3.1 Principle

- Cohorting of inpatients on transmission-based precautions will be initiated in response to bed management when there is a shortage of single room availability.
- Cohorting will only be undertaken for interim until further options become available and the demand for single room subsided.
- Cohorting of inpatients will be in consultation with the Clinical Microbiologist and IP&M team and will only be initiated after thorough case review/risk assessment.

Refer to Appendix 1 for [Risk Assessment Guide for Patient Placement](#)

### 3.2 Suitability criteria for cohorting

| Suitable for Cohorting                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           | Not Suitable for Cohorting                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <ol style="list-style-type: none"> <li><b>Laboratory confirmed respiratory virus diseases.</b> <ul style="list-style-type: none"> <li>Patients with same type and strain</li> <li>Patients without other infectious illness or micro-alerts.</li> </ul> </li> <li><b>Laboratory confirmed <i>Clostridioides difficile</i> or infective gastroenteritis.</b> <ul style="list-style-type: none"> <li>Patients with the same confirmed causative organisms and no other risk factors.</li> </ul> </li> <li>Patients with same micro-alert (<b>except for Micro C and G</b>) with no other risk factors.</li> <li><b>Laboratory confirmed respiratory virus diseases.</b> <ul style="list-style-type: none"> <li>Patients with same laboratory confirmed infection and no other risk factors.</li> </ul> </li> </ol> | <ol style="list-style-type: none"> <li>Unconfirmed infectious illness including viral respiratory illness with pending laboratory test result.</li> <li>Requiring airborne precautions (<b>Priority for NPIR use</b>).</li> <li>Unknown cause of diarrhoea.</li> <li>Dual infections.</li> <li>Infectious illness with laboratory test result that has not detected a known virus.</li> <li>Direct transfer from hospital outside of Western Australia.</li> <li>High risks (<b>neonates, burns patients, pregnant women, multiple morbidities</b>) and severely immunocompromised.</li> </ol> |
| Risk factors to consider                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
| <ul style="list-style-type: none"> <li>Wandering and cognitive impairment</li> <li>Safety/Security issue</li> <li>Incontinence</li> <li>Non-intact skin (open/draining wounds)</li> <li>Invasive devices</li> <li>Poor hygiene practices</li> <li>Aerosol generating behaviours</li> <li>Aerosol generating procedures</li> <li>Clinical symptoms (Diarrhoea, vomiting, coughing, sneezing)</li> <li>Micro-alert</li> <li>Suppressed immune state</li> </ul>                                                                                                                                                                                                                                                                                                                                                     |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |

### 3.3 Process flow for cohorting

- All alternative options must be considered before cohorting patients.
- Ward nurse coordinator or nurse unit manager must notify IP&M team during normal business hours or the after-hours nurse manager (AHNM)/clinical nurse specialist (ACNS) for the need to cohort patients.
- IP&M team will conduct a risk assessment to determine cohorting suitability during normal business hours. For after-hours, AHNM/CNS will conduct the risk assessment and inform IP&M by email.
- IPM team or AHNM/ACNS will contact the on-call clinical microbiologist and discuss the cohorting plan.
- IP&M team or AHNM/CNS will notify the director of nursing or executive on-call.
- Once risk assessment and clinical consultation have been completed, patient can be cohorted.
- Cohorted patients must be re-evaluated daily, and appropriate placement be considered once alternative options become available.

#### 4. Management of Inpatients in cohort room/area

|                                      | Actions/Procedures                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
|--------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Room/Bathroom</b>                 | <ul style="list-style-type: none"> <li>▪ To fulfill the principles of isolation, the cohorting room must have a closing door and preferably en-suite facilities.</li> <li>▪ If no en-suite facilities, a dedicated bathroom must be assigned with appropriate precaution card displayed on the door that is clearly visible.</li> <li>▪ Bedside commode can be provided if necessary.</li> <li>▪ Curtains must be drawn to create a partition between patients.</li> <li>▪ Patient must be distanced/separated for at least 1.5 metres apart.</li> </ul> |
| <b>Precautions Cards</b>             | Nursing staff must ensure that the appropriate precaution card is displayed clearly on the main door.                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| <b>Personal Protective Equipment</b> | <ul style="list-style-type: none"> <li>▪ Clean PPE must be placed outside the cohort room/area and be safe from contamination risk.</li> <li>▪ If gowns and gloves are required in the care of cohorted patients, these must be changed in between each patient.</li> <li>▪ Masks and eye protection (face shield or goggles) do not need to be changed between each patient, except when soiled or touched.</li> <li>▪ PPE must also be used as per standard precautions, when there is a risk of a body fluid exposure.</li> </ul>                     |
| <b>Cleaning and disinfection</b>     | <ul style="list-style-type: none"> <li>▪ Two (2) times daily cleaning and disinfection of shared bathroom.</li> <li>▪ Frequent cleaning and disinfection of commonly touched surfaces.</li> <li>▪ Only use the hospital approved disinfectant wipes/solutions.</li> <li>▪ Follow the hospital cleaning guidelines.</li> <li>▪ Follow AKG linen management.</li> </ul>                                                                                                                                                                                    |
| <b>Hand hygiene</b>                  | <ul style="list-style-type: none"> <li>▪ Follow the 5 moments for hand hygiene.</li> <li>▪ Use soap and water for hand hygiene when managing patient with enteric infections (e.g., norovirus, <i>C. diff</i>).</li> </ul>                                                                                                                                                                                                                                                                                                                               |
| <b>Staffing</b>                      | <ul style="list-style-type: none"> <li>▪ There should be limited number of staff/persons entering the cohort room/area.</li> <li>▪ Staff who are not immune to vaccine preventable diseases and those who are pregnant should not be allocated for the care of cohorted patients.</li> </ul>                                                                                                                                                                                                                                                             |
| <b>Patient equipment</b>             | <ul style="list-style-type: none"> <li>▪ Patient equipment should be disposable where possible or dedicated to the cohort care area.</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                          |

|                              |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
|------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                              | <ul style="list-style-type: none"> <li>All shared equipment must be thoroughly cleaned and disinfected between each patient use.</li> </ul>                                                                                                                                                                                                                                                                                                                                                                     |
| <b>Patient education</b>     | <ul style="list-style-type: none"> <li>Patients must be educated on the importance of hand hygiene, especially after using the bathroom.</li> <li>Patients must be educated about cough etiquette.</li> <li>Patients should be educated on the importance of personal hygiene.</li> <li>Patients should be advised to inform nursing staff when they have used the bathroom facility or if the area is noted to be visibly soiled, so cleaning and disinfecting can be arranged as soon as possible.</li> </ul> |
| <b>Transfer and movement</b> | <ul style="list-style-type: none"> <li>Patient transfer and movement should be restricted as much as possible. Non-urgent medical procedures should be delayed until the infectious period has lapsed.</li> <li>Medical procedures (e.g., X-rays, ultrasound) should be done in the cohort room/area.</li> </ul>                                                                                                                                                                                                |
| <b>Visitors and Family</b>   | <ul style="list-style-type: none"> <li>Visitors' restrictions should be in-place during the infectious period.</li> <li>Educate visitors and family about hand-hygiene practices and correct PPE use if required.</li> </ul>                                                                                                                                                                                                                                                                                    |

## 5. Legislative context

The following documents are required to give effect to this procedure [i.e. the documents included are mandatory]:

- [Health Services Act 2016](#)
- [Public Health Act 2016](#)
- [Public Health Regulations 2017](#)
- Australian Commission on Safety and Quality in Health Care (ACSQHC) [Recommendations for the control of Multi-drug resistant Gram-negatives: Carbapenem resistant Enterobacteriaceae](#)
- Health Department of Western Australia (HDWA) [Guidelines for the Screening and Management of Multi-resistant Organisms in Healthcare Facilities](#)
- East Metropolitan Health Service (EMHS) [Infection Prevention and Management Policy \(EMHS: 34\)](#)
- EMHS [Immunisation for Health Care Workers \(EMHS: 11\)](#)
- EMHS [Cleaning Standard 2022](#)
- Communicable Disease Network Australia (CDNA) [Series of National Guidelines \(SoNGs\)](#)
- National Health and Medical Research Council (NHMRC) [Australian Guidelines for the Prevention and Control of Infection in Healthcare](#)

## 6. Supporting information

The following documents support the implementation of this procedure:

- [Influenza-Like Illness Infection Prevention Guidelines \(AKG-GUI-0208-15\)](#)
- [Hand Hygiene \(AKG-POL-0187-17\)](#)
- [Infection Prevention and Management for Multi-resistant Organisms \(AKG-GUI-0621/23\)](#)
- [Standard and Transmission-based Precautions \(AKG-PRO-0513-19\)](#)
- [Cleaning of Reusable Clinical Non-invasive Equipment Procedure \(AKG-PRO-0383-17\)](#)
- [Health Care Worker Immunisation and Health \(AKG-PL-0188-14\)](#)

## 7. Compliance, monitoring and evaluation.

Compliance against this procedure will be monitored by the Infection Prevention and Management Committee.

Evaluation and monitoring activities will include:

- Observational audits in the clinical areas regarding adherence to: Transmission-based and standard precautions, correct PPE use, hand hygiene, bare below the elbow practices, waste management, aseptic technique, cleaning and disinfection practices and hand hygiene product placement.
- Reporting of infection rates through the Infection Prevention, Policy, and Surveillance Unit (IPPSU).
- Reporting of occupational exposures to blood and body fluids
- Reporting of any breaches and non-compliances to infection prevention practices.
- PPE training monitoring and assessment.
- Escalation of issues identified/concerns raised to the relevant committees for review and quality improvement.

## 8. Relevant standards

[National Safety and Quality Health Service \(NSQHS\) Standards](#) (second edition):

- Standard 3 – Action 3.6
- Standard 3 – Action 3.7

Standards Australia/New Zealand Standards (AS/NZS):

- [AS/NZS 3666.3.2000 Air Handling and Water Systems of Buildings](#)
- [AS/NZS 4187:2003 Cleaning, Disinfecting and Sterilising Reusable Medical and Surgical Instruments and Equipment](#)

## 9. Policy owner (relating to these procedures)

Enquiries relating to this procedure may be directed to:

Department: Infection Prevention and Management  
Email: [ahsinfection.controlnurses@health.wa.gov.au](mailto:ahsinfection.controlnurses@health.wa.gov.au)

## 10. Review

This procedure will be reviewed and evaluated as required to ensure relevance and currency. At a minimum it will be reviewed within one (1) year after first issue and at least every three (3) years thereafter.

| Version | Effective from | Effective to | Amendment(s) |
|---------|----------------|--------------|--------------|
| V3      | 29/04/2024     | 29/04/2027   |              |

## 11 Authorisation

This procedure has been authorised and issued by the AKG Executive Director.

|                    |                                                     |
|--------------------|-----------------------------------------------------|
| Authorised by      | Helen Fullarton – Director of Nursing and Midwifery |
| Authorisation date | 29/04/2024                                          |
| Published date     | 02/05/2024                                          |

## Appendix 1: Risk Assessment Guide for Patient Placement

| Risk Factors | Source and modes of transmission | Clinical predictors of transmission | Clinical impact of transmission | Room availability |
|--------------|----------------------------------|-------------------------------------|---------------------------------|-------------------|
|--------------|----------------------------------|-------------------------------------|---------------------------------|-------------------|



## Single Room Prioritisation and Cohorting of Inpatients on Transmission-based Precautions Procedure

| Question for Consideration | <ul style="list-style-type: none"> <li>Is human to human transmission known?</li> <li>Is/are the mode/s of transmission known?</li> <li>Has the person recently returned from overseas travel?</li> <li>What is the infectivity of the organism?</li> </ul> | <ul style="list-style-type: none"> <li>Does the patient have factors that would increase the risk of transmission?</li> </ul>                                                                                                                                                                                                                                                                                   | <ul style="list-style-type: none"> <li>How susceptible are the other patients in the area?</li> <li>What is the morbidity and mortality associated with the organism/condition disease?</li> <li>Will the safety of the individual who is to be isolated be affected?</li> </ul>                         | <ul style="list-style-type: none"> <li>What is the availability of negative pressure isolation rooms?</li> <li>What competing priorities exist for single room provision?</li> <li>Are single rooms with designated toilet facilities available?</li> <li>Are there other patients with the same organism species and/or strain that could be cohorted?</li> </ul> |
|----------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Examples                   | <ul style="list-style-type: none"> <li>Suspected or confirmed acute respiratory infection</li> <li>Public health notification</li> </ul>                                                                                                                    | <ul style="list-style-type: none"> <li>Wandering</li> <li>Cognitive impairment</li> <li>Incontinence</li> <li>Broken skin</li> <li>Open/draining wounds</li> <li>Invasive devices</li> <li>Poor hygiene practices</li> <li>Clinical symptoms such as:                             <ul style="list-style-type: none"> <li>Diarrhoea</li> <li>Vomiting</li> <li>Coughing</li> <li>Sneezing</li> </ul> </li> </ul> | <ul style="list-style-type: none"> <li>Organism not easily transmitted but associated with high mortality rate</li> <li>Immunosuppressed patients</li> <li>Neonates and young children</li> <li>Elderly patients</li> <li>Patients with burns</li> <li>Renal patients</li> <li>Pregnant women</li> </ul> | <ul style="list-style-type: none"> <li>Patients requiring high security or one-on-one observation</li> <li>Patient requiring end-of-life care</li> <li>Privacy and dignity issues</li> <li>Existing cohorts</li> </ul>                                                                                                                                             |

Adopted from: [Clinical Excellence Commission 2016](#)