



Structured Administration & Supply Arrangement (SASA)

ColonLYTELY® (Pre Admission Clinic)

Issuing Authority

Health Service	Sir Charles Gairdner Osborne Park Health Care Group (SCGOPHCG)
Address	Hospital Avenue, Nedlands, Western Australia
Contact	6457 2348

Authorisations

Practitioners:	SCGOPHCG Registered Nurse SCGOPHCG Medication Competent Enrolled Nurse
Practice Setting:	Pre Admission Clinic
Qualification:	Registered with the Nursing & Midwifery Board of Australia Enrolled Nurse who has completed a Board-approved qualification in administration of medicines
Authorisation:	Supply
Risk rating:	Low
Patients	Eligible outpatients as outlined within inclusion criteria of this SASA

Approved Sites

Site	Site Address
1.	Sir Charles Gairdner Hospital, Hospital Avenue, Nedlands, WA
2.	Osborne Park Hospital, Osborne Place, Stirling, WA

ColonLYTELY®

Route:	Per Oral	Brand:	ColonLYTELY®
Form:	Powder for oral solution	Strength:	Each 70 g sachet of ColonLYTELY® contains: 5.68 g Sodium sulfate 59 g Macrogol 3350 1.69 g Sodium bicarbonate 1.46 g Sodium chloride 0.75 g Potassium chloride
Dose:	Up to 4 sachets (4 L)	Quantity:	Maximum of 4 sachets
Instructions:	Detailed written instructions for food & drink consumption prior to surgery/procedure should be given to the patient. <i>The timing and number of sachets to be taken varies depending on the surgery/procedure undertaken. Ensure patient is given specific written instructions regarding the timing and number of sachets to be taken.</i> In general: Dissolve the contents of 1 sachet in 1 L of water. Repeat the process until the required volume is prepared. The made-up solution may be chilled in the refrigerator to increase palatability. Bowel motion usually starts after about 1 hour, usually within 3 hours, which normally removes the bowel contents.		
Administration Notes:	Advise patient to: • Keep medicine in the sachet until it is time to mix and take a dose. • Keep ColonLYTELY® powder in a cool, dry place where it stays below 25°C. • See ColonLYTELY® consumer information for more detailed information regarding preparation and administration. Oral medication taken within one hour of the commencement of the administration of ColonLYTELY® may be flushed from the gastrointestinal tract and not absorbed.		

Patient Inclusion:	Patients requiring emptying and cleansing of the large bowel in preparation for gastrointestinal diagnostic or surgical procedures.
Patient Exclusion:	Known documented allergy to the active substance or to any of the excipients. Gastrointestinal obstruction (partial or complete), gastric retention, bowel perforation, severe inflammatory bowel disease (i.e. toxic colitis or toxic megacolon) - contraindicated.
Patient Precautions:	Use with caution in patients with impaired renal function, pre-existing electrolyte disturbances (eg hypokalaemia), active inflammatory bowel disease, heart conditions and in the elderly – fluid and electrolyte disturbances can occur (monitor closely). Diabetic patients: a liquid diet may affect blood glucose levels and diabetic medications may require adjustment. ColonLYTELY® inactivates the low dose contraceptive pill, discuss risks with patients taking contraceptive medication.
Safety in Pregnancy/ Breastfeeding:	Pregnancy: safe to use. Breastfeeding: safe to use.
Documentation:	Outpatients: • Outpatient notes inclusive of MR 410 • Electronically via approved electronic software (i.e. hospital approved DMR)
Adverse Effects:	Adverse effects are transient and subside rapidly. ColonLYTELY® administration is associated with a low incidence of adverse effects, primarily nausea, abdominal fullness and bloating. Abdominal cramps, vomiting, and anal irritation occur less frequently.



Related policies, procedures, and guidelines

SCGOPHCG Guidelines

- [Medicines Management](#)
- [Clinical Practice Guideline – Gastrointestinal Endoscopic Procedures](#)

References

1. Australian Medicines Handbook. Macrogol laxatives. In: Australian Medicines Handbook [Internet]. Adelaide (South Australia): Australian Medicines Handbook; 2025 [cited 2025 Aug 04] Available from: [Australian Medicines Handbook \(health.wa.gov.au\)](http://health.wa.gov.au)
2. Structured Administration and Supply Arrangements, Department of Health. Available at https://ww2.health.wa.gov.au/Articles/S_T/Structured-Administration-and-Supply-Arrangements
3. [SCGOPHCG Structured Administration and Supply Arrangements \(SASAs\)](#)
4. Pregnancy and Breastfeeding Medicines Guide [Internet]. Melbourne (VIC): The Royal Women's Hospital; Macrogols (Polyethylene Glycol (Peg)) [cited 2025 Aug 04]. Available from: [macrogols \(Polyethylene glycol \(PEG\)\) | PBMG](#)

Document ID	SCGOPHCG.SASA.Colonlytely_PAC	Date of Issue:	9/09/2025
Record No.	MMS33	Date of Expiry:	9/09/2027
Poisons Permit Holder			
Name	Michael Petrovski (SCGH) Melinda Leeder (OPH)	Endorsed	8/09/2025
Authorising Committee			
Committee	Medicines and Therapeutics Committee (MTC)		
Chairperson	Dr Deepan Krishnasivam	Endorsed	8/09/2025
Nursing			
Name	Dr Suzanne Sinni	Signed:	
Position	A/Area Director Nursing and Midwifery Services	Date:	16/09/2025
Senior Medical Practitioner			
Name	Dr Deepan Krishnasivam	Signed:	
Position	A/Area Director Clinical Services	Date:	15/09/2025
Executive			
Name	Dr Jodi Graham	Signed:	
Position	Executive Director	Date:	18/09/2025
Keywords:	Bowel preparation, ColonLYTELY, Saline laxatives, Macrogol laxatives		
Owner:	Chief Pharmacist		
Standards	NSQHS Standards: 1  Governance, 4  Medication Safety		
Printed or personally saved electronic copies of this document are considered uncontrolled. Access the current version from the SCGOPHCG intranet. For any enquiries relating to this guideline, please email SCGOPHCG.MTC@health.wa.gov.au			