



Structured Administration & Supply Arrangement (SASA)

Mometasone 0.1% Cream (WHS)

Issuing Authority

Health Service:	Sir Charles Gairdner Osborne Park Health Care Group
Address:	Hospital Avenue, Nedlands, Western Australia
Contact:	6457 2348

Authorisations

Practitioners:	Nurses employed by SCGOPHCG
Practice Setting:	Work Health and Safety (WHS)
Qualification:	Nurse registered with the Nursing & Midwifery Board of Australia
Authorisation:	Supply
Risk rating:	Low
Patients:	Eligible staff members as outlined within inclusion criteria

Approved Sites

Site	Site Address
1.	Sir Charles Gairdner Hospital, Hospital Avenue, Nedlands, WA
2.	Osborne Park Hospital, Osborne Place, Stirling, WA

Mometasone 0.1% Cream 15g

Route:	Topical	Brand:	Novasone® Elocon®
Form:	Cream	Strength:	0.1% w/w
Dose:	Apply sparingly to the affected skin area(s) once daily.	Quantity:	15g (1 tube)
Instructions:	Apply to the affected area(s) on the hands once a day in the evening for up to 7 days. Sufficient cream should be applied so that the entire affected area is covered. The cream can then be rubbed or massaged into the inflamed skin. Note: One fingertip unit is enough to cover an area of skin twice the size of a flat adult with the fingers together. Use for a MAXIMUM of 7 days. If symptoms persist after 7 days, see WHS nurse for review. If supply is exhausted within 7 days of initial provision, see WHS nurse for review.		
Administration Notes:	Use after bathing and cleaning hands. Apply enough to cover affected areas. For external use only. Use in conjunction with recommended non-pharmacological treatment e.g. soap-free wash, barrier cream and moisturisers. If using in conjunction with a barrier cream and/or moisturiser, it is recommended to apply the corticosteroid cream first.		
Patient Inclusion:	Short term management of contact dermatitis of the hand(s) related to workplace exposure to hand sanitiser and/or chemicals.		
Patient Exclusion:	History of allergic reaction to mometasone furoate, any other topical corticosteroid, or any of the excipients listed at the end of the product Consumer Medicines Information (CMI). Infected skin – do not use topical corticosteroid. Broken or ulcerated skin - do not use topical corticosteroid. Women who are pregnant – Medical Officer review required.		

Safety in Pregnancy/Breastfeeding:	Pregnancy: Refer to Medical Officer. Short term use of topical mometasone is considered safe during pregnancy if it is the medicine of choice. However, mild to moderate potency topical corticosteroids are preferred in pregnancy, especially in the first trimester. It is recommended to use the lowest appropriate potency of topical corticosteroid for the shortest time necessary where moisturisers and other simple measures are inadequate. Mometasone 0.1% is considered a potent corticosteroid. Breastfeeding: Safe to use on the hands. Ensure breast area is free of corticosteroid before breastfeeding.
Documentation:	Entry of supply into WHS Staff Health Database.
Adverse Effects:	Transient burning, stinging or pain on application. Adverse effects which may arise from prolonged use include skin atrophy, delayed wound healing, striae, purpura, telangiectasia, acneiform eruptions.

Related Documents

Department of Health:

- Structured Administration and Supply Arrangements



SCGOPHCG:

- Healthcare Worker Health and Immunisation Policy
- Structured Administration and Supply Arrangements Policy

References

1. Australian Medicines Handbook (online). Adelaide: Australian Medicines Handbook Pty Ltd [2025 Jan]. Available from [Australian Medicines Handbook \(health.wa.gov.au\)](https://health.wa.gov.au)
2. Elocon® Product Information [Internet]. AusDI; c2023. [cited 2025 Oct 13]. Available from [Elocon® - AusDI \(health.wa.gov.au\)](https://health.wa.gov.au)
3. Women's PBMG. Pregnancy and Breastfeeding Medicines Guide [Internet]. Melbourne (AU): The Royal Women's Hospital; [cited 2025 Oct 13]. Available from: <https://thewomenspbmg.org.au>
4. Australasian College of Dermatologists. How do I apply topical corticosteroids for the treatment of eczema? [Internet]. Sydney: ACD; 2022 Sep [cited 2025 Oct 13]. Available from: <https://www.dermcoll.edu.au/wp-content/uploads/2022/09/ACD-Fact-Sheet-How-do-I-apply-topical-corticosteroids-for-the-treatment-of-eczema-September-2022.pdf> [dermcoll.edu.au]

Authorisation & Version Control

Document ID:	SCGOPHCG.SASA.Mometasone_0.1% Cream_WHS		Date of Issue:	08/12/2025	
Record No.	MMS41	Version No.	1.0	Date of Expiry:	08/12/2027
Poisons Permit Holder					
Name:	Michael Petrovski (SCGH) Melinda Leeder (OPH)		Endorsed:	08/12/2025	
Authorising Committee					
Committee:	SCGOPHCG Medicines and Therapeutics Committee				
Chairperson:	Michael Petrovski		Endorsed:	08/12/2025	
Nursing					
Name:	Dr Joanne Reid		Signed:		
Position:	A/Director Nursing and Midwifery		Date:	5/1/2026	
Senior Medical Practitioner					
Name:	Dr Simon Wood		Signed:		
Position:	Director Clinical Services		Date:	5/1/2026	
Executive					
Name:	Jodi Graham		Signed:		
Position:	Executive Director		Date:	5/1/2026	
Keywords:	Mometasone, steroid cream, dermatitis, WHS, work health and safety				
Owner:	Chief Pharmacist				
Standards:	NSQHS Standards: 1  Governance, 4  Medication Safety				
Printed or personally saved electronic copies of this document are considered uncontrolled. Access the current version from the SCGOPHCG Policy Hub. For any enquiries relating to this guideline, please email SCGOPHCG.MTC@health.wa.gov.au					