



Structured Administration & Supply Arrangement (SASA)

Mupirocin nasal ointment 2%

Issuing Authority

Health Service	Sir Charles Gairdner Osborne Park Health Care Group
Address	Hospital Avenue, Nedlands, Western Australia
Contact	6457 2348

Authorisations

Practitioners:	SCGOPHCG Registered Nurse SCGOPHCG Medication Competent Enrolled Nurse
Practice Setting:	• Pre-admission Clinic (PAC) Inpatient wards where surgery or procedure is applicable. This includes: ◦ G52, G53, G62, G63, G65, G66, G71, G73 and OPH Ward 6 • Haemodialysis Centre Patients and Peritoneal Dialysis Inpatient and Outpatients • Day Therapy Unit Haematology and Oncology patients • Radiology department
Qualification:	Registered with the Nursing & Midwifery Board of Australia Enrolled Nurse who has completed a Board-approved qualification in administration of medicines
Authorisation:	Initiation, administration, and supply
Risk rating:	Low
Patients	Eligible patients as outlined in SCGOPHCG Mupirocin Nasal Ointment for Decolonisation

Approved Sites

Site	Site Address
1.	Sir Charles Gairdner Hospital, Hospital Avenue, Nedlands, WA
2.	Osborne Park Hospital, Osborne Place, Stirling, WA

Mupirocin Nasal Ointment 2%


Route:	Intranasal	Brand:	Any stocked brand
Form:	Ointment	Strength:	2%
Dose:	Small amount "match-head size"	Quantity:	5g
Instructions:	Refer to: SCGOPHCG Mupirocin Nasal Ointment for Decolonisation <i>Application of nasal mupirocin 2% ointment TWICE daily inside BOTH nostrils for a total of FIVE days, where possible starting 5 days prior to procedure</i> <u>Note:</u> For insertion of non-tunnelled large bore catheters, tunnelled (subcutaneous) large bore catheters and peritoneal dialysis catheter, continue to apply nasal mupirocin 2% ointment inside BOTH nostrils ONCE a week following insertion and for the duration that the catheter remains in situ.		
Administration Notes:	• Apply "match head" sized quantity of ointment to inside of each nostril with a clean cotton bud. • Spread the ointment around the nasal vestibule by squeezing the nose between thumb and forefinger and rubbing them together. • Mupirocin may be tasted at the back of the throat after application. • Hand hygiene is essential to minimise the spread of infection.		
Patient Inclusion:	Patients undergoing a surgery or procedure admitted under: • Orthopaedic surgery (procedures involving implantation of prosthetic material) • Cardiothoracic surgery (all) • Neurosurgery (all) Patients for insertion of any of the following: • Non-tunnelled large bore central venous catheters (e.g. Vas-Cath™) • Tunnelled (subcutaneous) large bore central venous catheters (e.g. Hickman™, Broviac™, Groshong™) • Totally Implantable Venous Access Device (e.g. Infusaport™, Port-a-cath™, PAS-Port™) • Peritoneal Dialysis (PD) catheter)		

Patient Exclusion:	Known or documented allergy to mupirocin or components of the formulation. Non high-risk surgical procedures/interventions outside the patient inclusion criteria of this SASA Patients undergoing routine insertion of short to medium term catheter insertions e.g. PICCs (peripherally inserted central catheter) or non-tunnelled, temporary CVCs
Documentation:	Product to be provided is pre-labelled with directions. Nurse supplying to patient must complete relevant fields on label by hand. Inpatients: • Nursing Admission Assessment MR852.2/MR132.2 additional Information section • WA Hospital Medication chart MR410/805 • Electronically via approved electronic software (i.e. MOSAIQ or hospital approved DMR) Outpatients: • Outpatients notes inclusive of MR 410 • Electronically via approved electronic software (i.e. MOSAIQ or hospital approved DMR) Pre admission clinic: • Day Procedure Checklist MR710 • Nursing Admission Assessment MR852.2/MR132.2 additional Information section • Pre-Admission Clinic MR419
Adverse Effects:	Irritation, stinging
Clinical Guidelines	SCGOPHCG Mupirocin Nasal Ointment for Decolonisation SCGOPHCG Radiological Examination and Neuro Interventional Procedures

Related policies, procedures and guidelines	
SCGOPHCG Medicines Management Infection Prevention and Control of Methicillin-resistant Staphylococcus aureus (MRSA) in Western Australian Healthcare Facilities NPG Preoperative Care SCGOPHCG Hip Fracture Management Orthopaedic implant and cardiothoracic surgery Patient Pamphlet PAC	

References

1. Structured Administration and Supply Arrangements, Department of Health.
Available at https://www2.health.wa.gov.au/Articles/S_T/Structured-Administration-and-Supply-Arrangements

Document ID	SCGOPHCG.SASA.Muprocín Nasal	Date of Issue:	3/06/2025
Record No.	MMS29	Date of Expiry:	3/06/2027
Poisons Permit Holder			
Name	Michael Petrovski (SCGH) Melinda Leeder (OPH)	Endorsed	12/05/2025
Authorising Committee			
Committee	SCGOPHCG Medicines and Therapeutics Committee (MTC)		
Chairperson	Dr Deepan Krishnasivam	Endorsed	12/05/2025
Nursing			
Name	Annie Thompson	Signed:	
Position	Area Director of Nursing	Date:	16/06/2025
Senior Medical Practitioner			
Name	Dr Simon Wood	Signed:	
Position	A/Director of Clinical Services	Date:	16/06/2025
Executive			
Name	Dr Jodi Graham	Signed:	
Position	Executive Director	Date:	16/06/2025
Keywords:	Mupirocin, Bactroban, surgery, MRSA, decolonisation, staphylococcus aureus, staph		
Owner:	Chief Pharmacist		
Standards	NSQHS Standards: 1 Governance  , 4 Medication Safety  ,		
Printed or personally saved electronic copies of this document are considered uncontrolled. Access the current version from the SCGOPHCG intranet. For any enquiries relating to this guideline, please email SCGH.DTC@health.wa.gov.au			