



Structured Administration & Supply Arrangement (SASA)

Nicotine Replacement Therapy

Issuing Authority

Health Service	Sir Charles Gairdner Osborne Park Health Care Group
Address	Hospital Avenue, Nedlands, Western Australia
Contact	6457 2348

Authorisations

Practitioners:	SCGOPHCG Registered Nurses and Pharmacists
Practice Setting:	Inpatient, Outpatient
Qualification:	Nurses and Pharmacists registered with respective Health Professional Board of Australia
Authorisation:	Initiate, Administer (Nurse Only) & Supply
Patients:	Eligible patients as outlined in SCGOPHCG Clinical Practice Guideline: Management of Nicotine Dependence

Approved Sites

Site	Site Address
1.	Sir Charles Gairdner Hospital, Hospital Avenue, Nedlands, WA
2.	Osborne Park Hospital, Osborne Place, Stirling, WA

Nicotine Replacement Therapy

Route:	Topical, buccal, sublingual	Brand:	Any stocked brands
Form:	Patch	Strength:	Patch: 21mg/24hr
	Gum		Gum: 2mg, 4mg
	Lozenge		Lozenge: 2mg, 4mg
	Mouth Spray		Mouth Spray: 1mg/dose
Dose:	Refer to: SCGOPHCG Clinical Practice Guideline: Management of Nicotine Dependence	Quantity:	<i>Nurse initiation – Inpatient:</i> Maximum of 2 consecutive doses
			<i>Supply – Outpatient:</i> Patches: 1 box (quantity of 28) Gum: 1 box (quantity of 24) Lozenges: 1 box (quantity of 72) Mouth Spray: 1 box (quantity of 1)
Instructions:	Offer an appropriate level of NRT in accordance with the algorithm in Appendix 2: SCGOPHCG Clinical Practice Guideline: Management of Nicotine Dependence . Alternative regimens must be prescribed by a medical officer. Ongoing supply in an inpatient setting must be charted by a Medical Officer or Pharmacist Initiated Medicines (PIM) credentialed or Partnered Medication Charting (PPMC) Pharmacist. Ongoing supply for outpatients can be facilitated by outpatient prescription by a Medical Officer or via GP.		
Administration Notes:	Patch: • Apply to a non-hairy, clean, dry place on the upper body. Rotate site of application each day to avoid irritation – do not use the same site for at least 3 days. • Note: if patches cause vivid dreams, 24-hour patches can be removed before bed and reapplied in the morning. Fold patch in half on the adhesive side and dispose of safely out of reach of children and pets. • Printable patient leaflet Gum: • Chew slowly; bite gum until tingling or bitter taste, then park gum between cheek and upper gum until tingling subsides.		

	Then chew again. Each piece lasts about 30mins (when it no longer causes tingling). • If the gum is too bulky, pieces may be cut in half and used more often. • Avoid eating or drinking whilst using this product as this can reduce the amount of nicotine absorbed in the mouth. • Printable patient leaflet Lozenges: • Place lozenge in the mouth between gum & cheek and allow to dissolve slowly (do not chew or swallow). • Move the lozenge from one side of the mouth to another until completely dissolved. • A lozenge may take up to 30 mins to completely dissolve. • Printable patient leaflet Mouth Spray: • Hold the spray nozzle as close as possible to the mouth, avoiding the lips and press down on the top of the dispenser firmly to release one spray into the mouth. • Do NOT inhale or swallow while spraying but hold the spray in the mouth for 2-3 seconds before swallowing. • If cravings are still present after 1 minute, repeat once more. • Has the fastest onset of action of all intermittent formulations so can also be used when cravings occur. • Printable patient leaflet		
Patient Inclusion:	Patients who are nicotine-dependent, as determined by an assessment using the Heaviness of Smoking Index		
Patient Exclusion: [Note: these relate to scenarios which are excluded from this SASA – treatment may be appropriate under the direction of a medical officer]	Women who are pregnant or breastfeeding – medical officer consultation required. Recent MI or stroke, unstable angina, severe arrhythmias Dentures – avoid use of gum Oral, oesophageal, pharyngeal or gastric inflammation – may be worsened by oral or inhaled nicotine; avoid oral/inhaled products Generalised skin disease – avoid use of patches Phenylketonuria – avoid use of lozenge (contains aspartame)		
Documentation:	Inpatient administration: <table border="1"> <tr> <td>Nurse-initiated</td><td>“Once only, pre-medication and nurse-initiated medicines” section of the WA Hospital Medication</td></tr> </table>	Nurse-initiated	“Once only, pre-medication and nurse-initiated medicines” section of the WA Hospital Medication
Nurse-initiated	“Once only, pre-medication and nurse-initiated medicines” section of the WA Hospital Medication		

		Chart (WA HMC) MR 410/805, Long-stay Medication Chart MR810.03 OR Pre-/Post-operative Protocol for Day Surgery Patients MR810.06
	Pharmacist-initiated	As per SCGOPHCG Partnered Charting Pharmacist Program General Information OR Pharmacist Initiated Medicines (PIM) Framework and Procedures
	Outpatient supply:	
	Anaesthesia Perioperative Clinic (Nurse)	Clinic letter, uploaded to iClinical Manager
	OPH Pre-Admission Clinic (Nurse)	Nursing Admission Assessment MR852.2/MR132.2, in the 'Additional Information' section
	SCGH Outpatient Clinics (Nurse, Pharmacist)	SCGH MR410 Outpatient Notes OR Electronically via approved electronic software (i.e. MOSAIQ or hospital approved DMR)
	Pre-Admission Clinic (Pharmacist)	Medication History and Management Plan MR410/804
Adverse Effects:	<i>Oral and inhaled products:</i> hiccups, irritation of throat or mouth and sinusitis <i>Patch:</i> application site reactions may occur <i>Other common adverse effects:</i> dizziness, headache, nausea, vomiting, burning-lip sensation (mouth spray), hiccups, indigestion, abdominal pain, myalgia, cough (inhalation), vivid dreams (especially 24-hour patch)	
Clinical Guidelines	SCGOPHCG Clinical Practice Guideline: Management of Nicotine Dependence.	

Related policies, procedures and guidelines

[SCGOPHCG Medication Management](#)

[NMHS Smoke Free Policy](#)

[SCGOPHCG Smoke Free Procedure](#)

[Guidelines for the WA Hospital Medication Chart](#)

[SCGOPHCG Clinical Practice Guideline: Management of Nicotine Dependence](#)

[SCGOPHCG Pharmacist Initiated Medicines \(PIMs\) Procedure](#)

References

1. Australian Medicines Handbook. [Internet]. Adelaide (South Australia): Australian Medicines Handbook; 2022 [cited 2022 April 13] Available from: <https://amhonline-amh-net-au.kelibresources.health.wa.gov.au>
2. MIMS Australia. In: MIMS Online [Internet]. St Leonards (New South Wales): MIMS Australia; 2022 [cited 2022 April 13]. Available from: <https://www-mimsonline-com-au.kelibresources.health.wa.gov.au>
3. Structured Administration and Supply Arrangements, Department of Health. Available at [https://ww2.health.wa.gov.au/Articles/S T/Structured-Administration-and-Supply-Arrangements](https://ww2.health.wa.gov.au/Articles/S_T/Structured-Administration-and-Supply-Arrangements)

Document ID	SCGOPHCG.SASA.NicotineReplacement Therapy	Date of Issue:	16/06/2025
Record No.	MMS02	Date of Expiry:	16/06/2027
Health Service Permit Holders			
Name	Michael Petrovski (SCGH) Melinda Leeder (OPH)	Endorsed	09/06/2025
Authorising Committee			
Committee	Medicines and Therapeutics Committee		
Chairperson	Dr Deepan Krishnasivam	Endorsed	09/06/2025
Nursing			
Name	Annie Thompson	Signed:	
Position	Area Director of Nursing	Date:	16/06/2025
Senior Medical Practitioner			
Name	Dr Simon Wood	Signed:	
Position	A/Director of Clinical Services	Date:	16/06/2025
Executive			
Name	Dr Jodi Graham	Signed:	
Position	Executive Director	Date:	16/06/2025
Keywords:	Nicotine Replacement Therapy, NRT, smoking cessation, nicotine dependence, SASA		
Owner:	Chief Pharmacist		
Standards	NSQHS Standards: 1  Governance, 4  Medication Safety,		
Printed or personally saved electronic copies of this document are considered uncontrolled. Access the current version from the SCGOPHCG intranet. For any enquiries relating to this guideline, please email SCGOPHCG.MTC@health.wa.gov.au			