



Structured Administration & Supply Arrangement (SASA)

Oxybuprocaine 0.4% Eye Drops for Eye Clinic

Issuing Authority

Health Service	Sir Charles Gardiner Osborne Park Health Care Group
Address	Hospital Avenue, Nedlands, Western Australia
Contact	6457 2348

Authorisations

Practitioners:	SCGOPHCG Registered Nurse SCGOPHCG Medication Competent Enrolled Nurse
Practice Setting:	Sir Charles Gairdner Hospital (SCGH) Ophthalmology Outpatient Clinic
Qualification:	Nurse registered with the Nursing & Midwifery Board of Australia who have completed the SCGOPHCG Ophthalmology Outpatient Clinic Orientation Competency Enrolled Nurse who has completed a Board-approved qualification in administration of medicines and completed the SCGOPHCG Ophthalmology Outpatient Clinic Orientation Competency
Authorisation:	Administration
Risk rating:	Low
Patients:	Eligible outpatients as outlined in the SCGH Administration of Eye Drops in the Ophthalmology Outpatient Clinic Guideline

Approved Sites

Site	Site Address
1.	Sir Charles Gardiner Hospital, Hospital Avenue, Nedlands, WA

Oxybuprocaine 0.4% Eye Drops

Route:	Topical	Brand:	Any stocked brand
Form:	Eye drop	Strength:	0.4%
Dose:	1 drop	Quantity:	0.5 mL (1 x single use Minim) Up to 4 drops
Instructions:	Instil 1 drop into affected eye(s) as required for consultation.		
Administration Notes:	- Tilt patient's head up and gently pull down lower eyelid to form a pouch. Avoid letting the tip of the dropper touch the eyes, skin, or any other surface. - To reduce systemic effects, advise patients to close their eyes and apply punctual occlusion after administration. Administer 1 drop into the procedural eye(s). Wait 5 minutes, then administer an additional 1 drop. Repeat this process up to a total of 4 drops (per eye). If urgent, then up to 3 drops can be administered in less than 5 minutes.		
Patient Inclusion:	Patients aged 16 years and over requiring topical anaesthesia for general consults/intravitreal injections/minor procedures.		
Patient Exclusion:	- Patient past history of allergy / sensitivity to Oxybuprocaine - Patients with history of corneal damage		
Safety in Pregnancy / Breastfeeding:	Pregnancy: Safe to use Breastfeeding: Safe to use		
Documentation:	Nurse to sign, date and time the Outpatient Ophthalmic Notes and Medication Chart MR 473 medication administration section.		
Adverse Effects:	<u>Common:</u> - Localised stinging <u>Rare:</u> - Allergic reaction		
Clinical Guidelines	Administration of Eye Drops in the Ophthalmology Outpatient Clinic		

Related policies, procedures and guidelines

[SCGOPHCG Medication Management](#)

[Guidelines for the WA Hospital Medication Chart](#)

[SCGOPHCG Structured Administration and Supply Arrangements \(SASAs\)](#)

References

1. [Structured Administration and Supply Arrangements \(health.wa.gov.au\)](https://health.wa.gov.au)
2. Australian Medicines Handbook. Adelaide: Australian Medicines Handbook Pty Ltd; 2024. Available [Australian Medicines Handbook](#)

Document ID	SCGOPHCG.SASA.Oxybuprocaine 0.4% Eye Drops	Date of Issue:	23/07/2025
Record No.	MMS24	Date of Expiry:	23/07/2027
Poisons Permit Holder			
Name	Michael Petrovski	Endorsed	14/07/2025
Authorising Committee			
Committee	Medicines and Therapeutics Committee		
Chairperson	Dr Deepan Krishnasivam	Endorsed	14/07/2025
Nursing			
Name	Annie Thompson	Signed:	
Position	Area Director of Nursing and Midwifery Services	Date:	08/08/2025
Senior Medical Practitioner			
Name	Dr Simon Wood	Signed:	
Position	A/Director Clinical Services	Date:	08/08/2025
Executive			
Name	Dr Jodi Graham	Signed:	
Position	Executive Director	Date:	08/08/2025
Keywords:	Oxybuprocaine 0.4% Eye Drops, SASA		
Owner:	Chief Pharmacist		
Standards	NSQHS Standards: 1  Governance, 4  Medication Safety		
Printed or personally saved electronic copies of this document are considered uncontrolled. Access the current version from the SCGOPHCG intranet. For any enquiries relating to this guideline, please email SCGOPHCG.MTC@health.wa.gov.au			