

Gate-level Minimization

CS207 Chapter 3

James YU

yujq3@sustech.edu.cn

Department of Computer Science and Engineering
Southern University of Science and Technology

Sept. 21, 2022



南方科技大学
SOUTHERN UNIVERSITY OF SCIENCE AND TECHNOLOGY



Gate-level minimization

- The complexity of digital logic gates to implement a Boolean function is directly related to the complexity of algebraic expression.
- Gate-level minimization is the design task of finding an optimal gate-level implementation of Boolean functions describing a digital circuit.
 - Difficult by hand for more than few inputs.
 - Typically by computer, need to understand the underlying principle.

The map method

- The map method, first proposed by Veitch and slightly improvised by Karnaugh, provides a simple, straightforward procedure for the simplification of Boolean functions.
 - Called *Karnaugh map*.
- The map is a diagram consisting of *squares*. For n variables on a Karnaugh map there are 2^n numbers of squares. *2ⁿ个格子*.
 - Each square or cell represents one of the minterms.
 - Since any Boolean function can be expressed as a sum of minterms, it is possible to recognize a Boolean function graphically in the map from the area enclosed by those squares whose minterms appear in the function.

Two-variable K-map

- A two-variable system can form four minterms

| $\backslash B$ A | 0 | 1 |
|-----------------------|-------|-------|
| 0 | m_0 | m_1 |
| 1 | m_2 | m_3 |

| $\backslash B$ A | 0 | 1 |
|-----------------------|--------|-------|
| 0 | $A'B'$ | $A'B$ |
| 1 | AB' | AB |

| $\backslash B$ A | 0 | 1 |
|-----------------------|---|---|
| 0 | 0 | 1 |
| 1 | 1 | 1 |

- The two-variable Karnaugh map is a useful way to represent any of the 16 Boolean functions.
 - Example:

$$\begin{aligned}A + B &= A(B + B') + B(A + A') \\&= AB + AB' + AB + A'B = AB + AB' + A'B\end{aligned}$$

- So the squares corresponding to AB , AB' , and $A'B$ are marked with 1.

Three-variable K-map

- Since there are eight minterms for three variables, the map consists of eight cells or squares.
 - Minterms are arranged, not according to the binary sequence, but according to the sequence similar to the gray code.
 - **Between two consecutive rows or columns, only one single variable changes its logic value from 0 to 1 or from 1 to 0.**
连续两行或两列只能改变一个数位上的数

| $A \backslash BC$ | | | | | |
|-------------------|--|-------|-------|-------|-------|
| | | 00 | 01 | 11 | 10 |
| 0 | | m_0 | m_1 | m_3 | m_2 |
| | | m_4 | m_5 | m_7 | m_6 |

Three-variable K-map



- To understand the usefulness of the map for simplifying the Boolean functions, we must observe the basic properties of the adjacent squares.
 - Any two adjacent squares in the Karnaugh map differ by only one variable, which is complemented in one square and uncomplemented in one of the adjacent squares. 在 karnaugh 映射中, 任何两个相邻的正方形只相差一个变量, 该变量在一个正方形中互补, 在另一个正方形中不互补。
 - The sum of two minterms can be simplified to a single AND term consisting of less number of literals.
 - $m_1 + m_5 = A'B'C + AB'C = (A' + A)B'C = B'C$

| | | BC | | | |
|---|---|-------|-------|-------|-------|
| | | 00 | 01 | 11 | 10 |
| A | 0 | m_0 | m_1 | m_3 | m_2 |
| | 1 | m_4 | m_5 | m_7 | m_6 |

Three-variable K-map

- Example: Simplify the Boolean function $F = A'BC + A'BC' + AB'C' + AB'C$.

| $A \backslash BC$ | 00 | 01 | 11 | 10 |
|-------------------|----|----|----|----|
| 0 | 0 | 0 | 1 | 1 |
| 1 | 1 | 1 | 0 | 0 |

- The first row: $A'BC + A'BC' = A'B$.
- The second row: $AB'C' + AB'C = AB'$.
- $F = A'B + AB'$.

Three-variable K-map

- Example: Simplify the Boolean function $F = A'BC + AB'C' + ABC + ABC'$.

| $A \backslash BC$ | 00 | 01 | 11 | 10 |
|-------------------|----|----|----|----|
| 0 | 0 | 0 | 1 | 0 |
| 1 | 1 | 0 | 1 | 1 |

同样相连

- The third column: $A'BC + ABC = BC$.
- The second row: $AB'C' + ABC' = AC'$.
- $F = BC + AC'$.

Three-variable K-map

- Example: Simplify the Boolean function $F = \sum(1, 2, 3, 5, 7)$.

| $A \backslash BC$ | 00 | 01 | 11 | 10 |
|-------------------|----|----|----|----|
| 0 | 0 | 1 | 1 | 1 |
| 1 | 0 | 1 | 1 | 0 |

- $F = C + A'B$.

Three-variable K-map

- Example: Simplify the Boolean function $F = \sum(0, 2, 4, 5, 6)$.

| $A \backslash BC$ | 00 | 01 | 11 | 10 |
|-------------------|----|----|----|----|
| 0 | 1 | 0 | 0 | 1 |
| 1 | 1 | 1 | 0 | 1 |

- $F = C' + AB'$.

Four-variable K-map



- Similar to the method used for two-variable and three-variable Karnaugh maps, four-variable Karnaugh maps may be constructed with 16 squares consisting of 16 minterms.

| $AB \backslash CD$ | | CD | | | |
|--------------------|----|----------|----------|----------|----------|
| | | 00 | 01 | 11 | 10 |
| AB | 00 | m_0 | m_1 | m_3 | m_2 |
| | 01 | m_4 | m_5 | m_7 | m_6 |
| | 11 | m_{12} | m_{13} | m_{15} | m_{14} |
| | 10 | m_8 | m_9 | m_{11} | m_{10} |

Four-variable K-map



- Two, four, or eight adjacent squares can be combined to reduce the number of literals in a function. 两个, 四个, 八个可以组合在一起.
- The squares of the top and bottom rows as well as leftmost and rightmost columns may be combined. 左右, 顶部与底部可以组合
 - When two adjacent squares are combined, it is called a *pair* and represents a term with three literals. 3 两个相邻: Pair
 - Four adjacent squares, when combined, are called a *quad* and its number of literals is two. 2 四个相邻: Quad
 - If eight adjacent squares are combined, it is called an *octet* and represents a term with one literal. 1 八个相邻: Octet
 - If, in the case all sixteen squares can be combined, the function will be reduced to 1.

Four-variable K-map



- Example: Simplify the Boolean function
 $F = m_1 + m_5 + m_{10} + m_{11} + m_{12} + m_{13} + m_{15}$.

- $A'B'C'D + A'BC'D = A'C'D$,
- $ABC'D' + ABC'D = ABC'$,

- $F = A'C'D + ABC' + ACD + AB'C$.
- This reduced expression is not a unique one.
 - If pairs are formed in different ways, the simplified expression will be different.

| AB \ CD | 00 | 01 | 11 | 10 |
|---------|----|----|----|----|
| 00 | | 1 | | |
| 01 | | 1 | | |
| 11 | 1 | 1 | 1 | |
| 10 | | | 1 | 1 |

简化方式不同, 简化结果也不同

Four-variable K-map

- Example: Simplify the Boolean function
 $F = m_1 + m_5 + m_{10} + m_{11} + m_{12} + m_{13} + m_{15}.$
- $F = A'C'D + ABC' + ABD + AB'C.$

| AB \ CD | 00 | 01 | 11 | 10 |
|---------|----|----|----|----|
| 00 | | 1 | | |
| 01 | | 1 | | |
| 11 | 1 | 1 | 1 | |
| 10 | | | 1 | 1 |

Four-variable K-map



- Example: Simplify the Boolean function
 $F = \sum(7, 9, 10, 11, 12, 13, 14, 15).$
- $F = AB + AC + AD + BCD.$

用最少的Group, 每个 Group 尽可能 big

| AB \ CD | 00 | 01 | 11 | 10 |
|---------|----|----|----|----|
| 00 | | | | |
| 01 | | | 1 | |
| 11 | 1 | 1 | 1 | 1 |
| 10 | | 1 | 1 | 1 |

Four-variable K-map



- Example: Plot the logical expression $F(A, B, C, D) = ABCD + AB'C'D' + AB'C + AB$ on a four-variable Karnaugh map.

$$\begin{aligned} &F(A, B, C, D) \\ &= ABCD + AB'C'D' + AB'C + AB \\ &= ABCD + AB'C'D' + AB'C(D + D') \\ &\quad + AB(C + C')(D + D') \\ &= \dots \\ &= \sum(8, 10, 11, 12, 13, 14, 15) \\ &= AB + AC + AD' \end{aligned}$$

| AB \ CD | 00 | 01 | 11 | 10 |
|---------|----|----|----|----|
| 00 | | | | |
| 01 | | | | |
| 11 | 1 | 1 | 1 | 1 |
| 10 | 1 | | 1 | 1 |

Four-variable K-map



- Simplify the expression $F(A, B, C, D) = \sum(0, 1, 2, 4, 5, 6, 8, 9, 12, 13, 14)$.


| AB \ CD | CD | | | |
|---------|----|----|----|----|
| | 00 | 01 | 11 | 10 |
| 00 | 1 | 1 | | 1 |
| 01 | 1 | 1 | | 1 |
| 11 | 1 | 1 | | 1 |
| 10 | 1 | 1 | | |

Four-variable K-map



- Simplify the expression $F(A, B, C, D) = A'B'C' + B'CD' + A'BCD' + AB'C'$.

$$\begin{aligned} &F(A, B, C, D) \\ &= A'B'C'(D + D') + B'CD'(A + A') \\ &\quad + A'BCD' + AB'C'(D + D') \\ &= A'B'C'D + A'B'C'D' + AB'CD' \\ &\quad + A'B'CD' + A'BCD' + AB'C'D \\ &\quad + AB'C'D' \\ &= \sum(0, 1, 2, 6, 8, 9, 10) \\ &= B'C' + B'D' + A'CD' \end{aligned}$$



简化时先看四个角

Four-variable K-map



- Simplify the expression
 $F(A, B, C, D) = \sum(3, 4, 5, 7, 9, 13, 14, 15)$.
 - It may be noted that one quad can also be formed, but it is redundant as the squares contained by the quad are already covered by the pairs which are essential.
- $F = A'BC' + A'CD + AY'D + ABC$.

| AB \ CD | 00 | 01 | 11 | 10 |
|---------|----|----|----|----|
| 00 | | | 1 | |
| 01 | 1 | 1 | 1 | |
| 11 | | 1 | 1 | 1 |
| 10 | | 1 | | |

Four-variable K-map



- Simplify the expression $F(A, B, C, D) = \prod(0, 1, 4, 5, 6, 8, 9, 12, 13, 14)$.
 - The above expression is given in respect to the maxterms.
 - 0's are to be placed instead of 1's at the corresponding maxterm squares.
- $F = C(B' + D)$.

π 找0 ☆

| AB \ CD | 00 | 01 | 11 | 10 |
|---------|----|----|----|----|
| 00 | 0 | 0 | 1 | 1 |
| 01 | 0 | 0 | 1 | 0 |
| 11 | 0 | 0 | 1 | 0 |
| 10 | 0 | 0 | 1 | 1 |

Four-variable K-map



- Simplify the expression $F(A, B, C, D) = \prod(0, 1, 4, 5, 6, 8, 9, 12, 13, 14)$.
 - The other way to achieve the minimized expression is to consider the 1's of the Karnaugh map.
- $F = CD + B'C = C(B' + D)$.

| AB \ CD | 00 | 01 | 11 | 10 |
|---------|----|----|----|----|
| 00 | 0 | 0 | 1 | 1 |
| 01 | 0 | 0 | 1 | 0 |
| 11 | 0 | 0 | 1 | 0 |
| 10 | 0 | 0 | 1 | 1 |

Five-variable K-map



- Karnaugh maps with more than four variables are not simple to use.
 - The number of cells or squares becomes excessively large and combining the adjacent squares becomes complex.
 - A five-variable Karnaugh map contains 2^5 or 32 cells.

五个以上的变量不再使用卡诺图
单元格太多 不易合并



Prime Implicants

主蕴含项

- A *prime implicant* is a product term obtained by combining the maximum possible number of adjacent squares in the map.
- The prime implicants of a function can be obtained from the map by combining all possible maximum numbers of squares. 组合所有可能的最大方数。
 - If a minterm in a square is covered by only one prime implicant, that prime implicant is said to be *essential*. 如果一个平方中的一个极小项只有一个主蕴含项覆盖, 这个蕴含项被称为最基本的。
- Gate-level minimization:
 - Determine all essential prime implicants.
 - Find other prime implicants that cover remaining minterms.
 - Logical sum all prime implicants.



Don't care conditions

- In practice, Boolean function is not specified for certain combinations of input variables. 没有符号值
 - Input combinations never occur during the process of a normal operation. 不出现输入组合
 - Those input conditions are guaranteed never to occur. 输入条件永不发生
- Such input combinations are called *don't-care conditions*.
- These input combinations can be plotted on the Karnaugh map for further simplification.
 - The don't care conditions are represented by d or X in a K-map.
 - They can be either 1 or 0 upon needed.



Don't care conditions

- Simplify the expression $F(A, B, C, D) = \sum(1, 3, 7, 11, 15), d = \sum(0, 2, 5)$.
- $F = A'B' + CD$.

可以让X变成
任意你需要的值

| AB \ CD | 00 | 01 | 11 | 10 |
|---------|----|----|----|----|
| 00 | X | 1 | 1 | X |
| 01 | | X | 1 | |
| 11 | | | 1 | |
| 10 | | | 1 | |



Don't care conditions

- Simplify the expression $F(A, B, C, D) = \sum(1, 3, 7, 11, 15), d = \sum(0, 2, 5)$.
- $F = A'D + CD$.

| AB \ CD | CD | | | |
|---------|----|----|----|----|
| | 00 | 01 | 11 | 10 |
| 00 | X | 1 | 1 | X |
| 01 | | X | 1 | |
| 11 | | | 1 | |
| 10 | | | 1 | |

More examples

- Using the Karnaugh map method obtain the minimal sum of the products expression for the function $F(A, B, C, D) = \sum(0, 2, 3, 6, 7) + d(8, 10, 11, 15)$.

| AB CD | 00 | 01 | 11 | 10 |
|------------------|----|----|----|----|
| 00 | 1 | | 1 | 1 |
| 01 | | | 1 | 1 |
| 11 | | | x | |
| 10 | 1 | | x | 1 |

$$B'D' + A'C$$