

Contents

Glen Islay	2
Site 1 -Wilding Risk Calculation	2
Site 2 -Wilding Risk Calculation	3
Site 3 -Wilding Risk Calculation	4
Site 4 -Wilding Risk Calculation	5
Site 5 -Wilding Risk Calculation	6
Site 6 -Wilding Risk Calculation	7
Site 7 -Wilding Risk Calculation	8
Site 8 -Wilding Risk Calculation	9
Site 9 -Wilding Risk Calculation	10
Site 10 -Wilding Risk Calculation	11
Site 11 -Wilding Risk Calculation	12
Site 12 -Wilding Risk Calculation	13
Site 13 -Wilding Risk Calculation	14
Site 14 -Wilding Risk Calculation	15
Site 15 -Wilding Risk Calculation	16
Site 16 -Wilding Risk Calculation	17
Site 17 -Wilding Risk Calculation	18
Site 18 -Wilding Risk Calculation	19
Site 19 -Wilding Risk Calculation	20
Site 20 -Wilding Risk Calculation	21
Site 21 -Wilding Risk Calculation	22
Site 22 -Wilding Risk Calculation	23
Site 23 -Wilding Risk Calculation	24
Site 24 -Wilding Risk Calculation	25
Site 25 -Wilding Risk Calculation	26

25 sites were assessed for Glen Islay to provide a comprehensive wilding risk assessment, for simplification a table has been included to summarize the results (Table 1). The highest value received during the wilding risk calculator assessment was 0 since Glen Islay is surrounded by intensively grazed farmland or plantation forest.

Section 11 (2) NES CF

Mort owns Venture Forestry, and has been involved with forest establishment since forming the company in 2011.

Mort is a registered forestry consultant with the New Zealand Institute of Forestry, Mort holds a Bachelor of Engineering Degree from the University of Canterbury and has 20 years industry experience in New Zealand, Canada and Southeast Asia in all aspects of forestry.

Glen Islay

Site 1 -Wilding Risk Calculation

1. SPECIES – GROWTH (score for one species only)

Spreading vigour varies with species

- Redwoods, Leyland cypresses, cedars and spruces (very low risk – no need to proceed further) 0
- Radiata (*P. radiata*) and ponderosa (*P. ponderosa*) pine, Lawsons cypress (*C. lawsoniana*) 1
- Muricata (*P. muricata*) and maritime (*P. pinaster*) pine and larches (*Larix* spp) 2
- Corsican (*P. nigra*) and mountain/dwarf mountain (*P. uncinata/mugo*) pine 3
- Douglas-fir ^g (*Ps. menziesii*), Scots pine (*P. sylvestris*) 4
- Lodgepole/contorta pine (*P. contorta*) 5

1

Enter score (0, 1, 2, 3, 4 or 5) here

2. SPECIES – PALATABILITY

Palatability varies with species

- Radiata, maritime and ponderosa pine 1
- Lodgepole and muricata pine and European larch 2
- Scots and mountain/dwarf mountain pine and Douglas-fir 3
- Corsican pine 4

1

Enter score (0, 1, 2, 3 or 4) here

3. SITING OF NEW PLANTING ^{c, d}

Trees are located on

- Sites well sheltered from prevalent and strong winds 0
- Flat sites (<10°), partially exposed to strong/prevalent winds 1
- Lea slopes where strong eddy gusts are likely 2
- Flat sites (<10°), fully exposed to strong/prevalent winds 3
- Either elevated 'take-off' sites, (ridge-tops, or base of exposed slopes >10°) or sloping land, fully exposed to strong/prevalent winds 4

3

Enter score (0, 1, 2, 3 or 4) here

4. DOWNWIND LANDUSE – GRAZING

Wilding establishment influenced by grazing (particularly with sheep)

- Intensive grazing on developed pasture 0^d
- Regular mob stocking with sheep ^e 1^d
- Semi-improved grazing (sheep/cattle)/ occasional mob stocking with sheep 2^d
- Extensive grazing only ^e 3^d
- No grazing 4^d

0

Enter score (0, 1, 2, 3 or 4) here

5. DOWNWIND VEGETATION COVER (if Douglas-fir involved see ^g in Notes)

Wilding establishment influenced by competition from existing vegetation

- Plantation forest, developed pasture (intensive grazing) 0^d
- Native forest^h, shrubland/tussock/grassland with a continuous and dense vegetation cover 1^d
- Forest/shrubland/tussock/grassland with few gaps 2^d
- Open forest and/or scattered patches of dense shrubland/tussock/grassland with many gaps 3^d
- Open slips/rockland and/or light, low-stature shrubland/tussock/grassland 4^d

0

TOTAL SCORE 0

Site 2 -Wilding Risk Calculation

1. SPECIES – GROWTH (score for one species only)

Spreading vigour varies with species

- Redwoods, Leyland cypresses, cedars and spruces (very low risk – no need to proceed further) 0
- Radiata (*P. radiata*) and ponderosa (*P. ponderosa*) pine, Lawsons cypress (*C. lawsoniana*) 1
- Muricata (*P. muricata*) and maritime (*P. pinaster*) pine and larches (*Larix* spp) 2
- Corsican (*P. nigra*) and mountain/dwarf mountain (*P. uncinata/mugo*) pine 3
- Douglas-fir ^g (*Ps. menziesii*), Scots pine (*P. sylvestris*) 4
- Lodgepole/contorta pine (*P. contorta*) 5

1

Enter score (0, 1, 2, 3, 4 or 5) here

2. SPECIES – PALATABILITY

Palatability varies with species

- Radiata, maritime and ponderosa pine 1
- Lodgepole and muricata pine and European larch 2
- Scots and mountain/dwarf mountain pine and Douglas-fir 3
- Corsican pine 4

1

Enter score (0, 1, 2, 3 or 4) here

3. SITING OF NEW PLANTING ^{c, d}

Trees are located on

- Sites well sheltered from prevalent and strong winds 0
- Flat sites (<10°), partially exposed to strong/prevalent winds 1
- Lea slopes where strong eddy gusts are likely 2
- Flat sites (<10°), fully exposed to strong/prevalent winds 3
- Either elevated 'take-off' sites, (ridge-tops, or base of exposed slopes >10°) or sloping land, fully exposed to strong/prevalent winds 4

3

Enter score (0, 1, 2, 3 or 4) here

4. DOWNWIND LANDUSE – GRAZING

Wilding establishment influenced by grazing (particularly with sheep)

- Intensive grazing on developed pasture 0^d
- Regular mob stocking with sheep ^e 1^d
- Semi-improved grazing (sheep/cattle)/ occasional mob stocking with sheep 2^d
- Extensive grazing only ^e 3^d
- No grazing 4^d

0

Enter score (0, 1, 2, 3 or 4) here

5. DOWNWIND VEGETATION COVER (if Douglas-fir involved see ^g in Notes)

Wilding establishment influenced by competition from existing vegetation

- Plantation forest, developed pasture (intensive grazing) 0^d
- Native forest^h, shrubland/tussock/grassland with a continuous and dense vegetation cover 1^d
- Forest/shrubland/tussock/grassland with few gaps 2^d
- Open forest and/or scattered patches of dense shrubland/tussock/grassland with many gaps 3^d
- Open slips/rockland and/or light, low-stature shrubland/tussock/grassland 4^d

0

TOTAL SCORE 0

Site 3 -Wilding Risk Calculation

1. SPECIES – GROWTH (score for one species only)

Spreading vigour varies with species

- Redwoods, Leyland cypresses, cedars and spruces (very low risk – no need to proceed further) 0
- Radiata (*P. radiata*) and ponderosa (*P. ponderosa*) pine, Lawsons cypress (*C. lawsoniana*) 1
- Muricata (*P. muricata*) and maritime (*P. pinaster*) pine and larches (*Larix* spp) 2
- Corsican (*P. nigra*) and mountain/dwarf mountain (*P. uncinata/mugo*) pine 3
- Douglas-fir ^g (*Ps. menziesii*), Scots pine (*P. sylvestris*) 4
- Lodgepole/contorta pine (*P. contorta*) 5

1

Enter score (0, 1, 2, 3, 4 or 5) here

2. SPECIES – PALATABILITY

Palatability varies with species

- Radiata, maritime and ponderosa pine 1
- Lodgepole and muricata pine and European larch 2
- Scots and mountain/dwarf mountain pine and Douglas-fir 3
- Corsican pine 4

1

Enter score (0, 1, 2, 3 or 4) here

3. SITING OF NEW PLANTING ^{c, d}

Trees are located on

- Sites well sheltered from prevalent and strong winds 0
- Flat sites (<10°), partially exposed to strong/prevalent winds 1
- Lea slopes where strong eddy gusts are likely 2
- Flat sites (<10°), fully exposed to strong/prevalent winds 3
- Either elevated ‘take-off’ sites, (ridge-tops, or base of exposed slopes >10°) or sloping land, fully exposed to strong/prevalent winds 4

3

Enter score (0, 1, 2, 3 or 4) here

4. DOWNWIND LANDUSE – GRAZING

Wilding establishment influenced by grazing (particularly with sheep)

- Intensive grazing on developed pasture 0^d
- Regular mob stocking with sheep ^e 1^d
- Semi-improved grazing (sheep/cattle)/ occasional mob stocking with sheep 2^d
- Extensive grazing only ^e 3^d
- No grazing 4^d

0

Enter score (0, 1, 2, 3 or 4) here

5. DOWNWIND VEGETATION COVER (if Douglas-fir involved see ^g in Notes)

Wilding establishment influenced by competition from existing vegetation

- Plantation forest, developed pasture (intensive grazing) 0^d
- Native forest^h, shrubland/tussock/grassland with a continuous and dense vegetation cover 1^d
- Forest/shrubland/tussock/grassland with few gaps 2^d
- Open forest and/or scattered patches of dense shrubland/tussock/grassland with many gaps 3^d
- Open slips/rockland and/or light, low-stature shrubland/tussock/grassland 4^d

0

TOTAL SCORE 0

Site 4 -Wilding Risk Calculation

1. SPECIES – GROWTH (score for one species only)

Spreading vigour varies with species

- Redwoods, Leyland cypresses, cedars and spruces (very low risk – no need to proceed further) 0
- Radiata (*P. radiata*) and ponderosa (*P. ponderosa*) pine, Lawsons cypress (*C. lawsoniana*) 1
- Muricata (*P. muricata*) and maritime (*P. pinaster*) pine and larches (*Larix* spp) 2
- Corsican (*P. nigra*) and mountain/dwarf mountain (*P. uncinata/mugo*) pine 3
- Douglas-fir ^g (*Ps. menziesii*), Scots pine (*P. sylvestris*) 4
- Lodgepole/contorta pine (*P. contorta*) 5

1

Enter score (0, 1, 2, 3, 4 or 5) here

2. SPECIES – PALATABILITY

Palatability varies with species

- Radiata, maritime and ponderosa pine 1
- Lodgepole and muricata pine and European larch 2
- Scots and mountain/dwarf mountain pine and Douglas-fir 3
- Corsican pine 4

1

Enter score (0, 1, 2, 3 or 4) here

3. SITING OF NEW PLANTING ^{c, d}

Trees are located on

- Sites well sheltered from prevalent and strong winds 0
- Flat sites (<10°), partially exposed to strong/prevalent winds 1
- Lea slopes where strong eddy gusts are likely 2
- Flat sites (<10°), fully exposed to strong/prevalent winds 3
- Either elevated 'take-off' sites, (ridge-tops, or base of exposed slopes >10°) or sloping land, fully exposed to strong/prevalent winds 4

2

Enter score (0, 1, 2, 3 or 4) here

4. DOWNWIND LANDUSE – GRAZING

Wilding establishment influenced by grazing (particularly with sheep)

- Intensive grazing on developed pasture 0^d
- Regular mob stocking with sheep ^e 1^d
- Semi-improved grazing (sheep/cattle)/ occasional mob stocking with sheep 2^d
- Extensive grazing only ^e 3^d
- No grazing 4^d

0

Enter score (0, 1, 2, 3 or 4) here

5. DOWNWIND VEGETATION COVER (if Douglas-fir involved see ^g in Notes)

Wilding establishment influenced by competition from existing vegetation

- Plantation forest, developed pasture (intensive grazing) 0^d
- Native forest^h, shrubland/tussock/grassland with a continuous and dense vegetation cover 1^d
- Forest/shrubland/tussock/grassland with few gaps 2^d
- Open forest and/or scattered patches of dense shrubland/tussock/grassland with many gaps 3^d
- Open slips/rockland and/or light, low-stature shrubland/tussock/grassland 4^d

0

TOTAL SCORE 0

Site 5 -Wilding Risk Calculation

1. SPECIES – GROWTH (score for one species only)

Spreading vigour varies with species

- Redwoods, Leyland cypresses, cedars and spruces (very low risk – no need to proceed further) 0
- Radiata (*P. radiata*) and ponderosa (*P. ponderosa*) pine, Lawsons cypress (*C. lawsoniana*) 1
- Muricata (*P. muricata*) and maritime (*P. pinaster*) pine and larches (*Larix* spp) 2
- Corsican (*P. nigra*) and mountain/dwarf mountain (*P. uncinata/mugo*) pine 3
- Douglas-fir ^g (*Ps. menziesii*), Scots pine (*P. sylvestris*) 4
- Lodgepole/contorta pine (*P. contorta*) 5

1

Enter score (0, 1, 2, 3, 4 or 5) here

2. SPECIES – PALATABILITY

Palatability varies with species

- Radiata, maritime and ponderosa pine 1
- Lodgepole and muricata pine and European larch 2
- Scots and mountain/dwarf mountain pine and Douglas-fir 3
- Corsican pine 4

1

Enter score (0, 1, 2, 3 or 4) here

3. SITING OF NEW PLANTING ^{c, d}

Trees are located on

- Sites well sheltered from prevalent and strong winds 0
- Flat sites (<10°), partially exposed to strong/prevalent winds 1
- Lea slopes where strong eddy gusts are likely 2
- Flat sites (<10°), fully exposed to strong/prevalent winds 3
- Either elevated 'take-off' sites, (ridge-tops, or base of exposed slopes >10°) or sloping land, fully exposed to strong/prevalent winds 4

2

Enter score (0, 1, 2, 3 or 4) here

4. DOWNWIND LANDUSE – GRAZING

Wilding establishment influenced by grazing (particularly with sheep)

- Intensive grazing on developed pasture 0^d
- Regular mob stocking with sheep ^e 1^d
- Semi-improved grazing (sheep/cattle)/ occasional mob stocking with sheep 2^d
- Extensive grazing only ^e 3^d
- No grazing 4^d

0

Enter score (0, 1, 2, 3 or 4) here

5. DOWNWIND VEGETATION COVER (if Douglas-fir involved see ^g in Notes)

Wilding establishment influenced by competition from existing vegetation

- Plantation forest, developed pasture (intensive grazing) 0^d
- Native forest^h, shrubland/tussock/grassland with a continuous and dense vegetation cover 1^d
- Forest/shrubland/tussock/grassland with few gaps 2^d
- Open forest and/or scattered patches of dense shrubland/tussock/grassland with many gaps 3^d
- Open slips/rockland and/or light, low-stature shrubland/tussock/grassland 4^d

0

TOTAL SCORE 0

Site 6 -Wilding Risk Calculation

1. SPECIES – GROWTH (score for one species only)

Spreading vigour varies with species

- Redwoods, Leyland cypresses, cedars and spruces (very low risk – no need to proceed further) 0
- Radiata (*P. radiata*) and ponderosa (*P. ponderosa*) pine, Lawsons cypress (*C. lawsoniana*) 1
- Muricata (*P. muricata*) and maritime (*P. pinaster*) pine and larches (*Larix* spp) 2
- Corsican (*P. nigra*) and mountain/dwarf mountain (*P. uncinata/mugo*) pine 3
- Douglas-fir ^g (*Ps. menziesii*), Scots pine (*P. sylvestris*) 4
- Lodgepole/contorta pine (*P. contorta*) 5

1

Enter score (0, 1, 2, 3, 4 or 5) here

2. SPECIES – PALATABILITY

Palatability varies with species

- Radiata, maritime and ponderosa pine 1
- Lodgepole and muricata pine and European larch 2
- Scots and mountain/dwarf mountain pine and Douglas-fir 3
- Corsican pine 4

1

Enter score (0, 1, 2, 3 or 4) here

3. SITING OF NEW PLANTING ^{c, d}

Trees are located on

- Sites well sheltered from prevalent and strong winds 0
- Flat sites (<10°), partially exposed to strong/prevalent winds 1
- Lea slopes where strong eddy gusts are likely 2
- Flat sites (<10°), fully exposed to strong/prevalent winds 3
- Either elevated 'take-off' sites, (ridge-tops, or base of exposed slopes >10°) or sloping land, fully exposed to strong/prevalent winds 4

1

Enter score (0, 1, 2, 3 or 4) here

4. DOWNWIND LANDUSE – GRAZING

Wilding establishment influenced by grazing (particularly with sheep)

- Intensive grazing on developed pasture 0^d
- Regular mob stocking with sheep ^e 1^d
- Semi-improved grazing (sheep/cattle)/ occasional mob stocking with sheep 2^d
- Extensive grazing only ^e 3^d
- No grazing 4^d

0

Enter score (0, 1, 2, 3 or 4) here

5. DOWNWIND VEGETATION COVER (if Douglas-fir involved see ^g in Notes)

Wilding establishment influenced by competition from existing vegetation

- Plantation forest, developed pasture (intensive grazing) 0^d
- Native forest^h, shrubland/tussock/grassland with a continuous and dense vegetation cover 1^d
- Forest/shrubland/tussock/grassland with few gaps 2^d
- Open forest and/or scattered patches of dense shrubland/tussock/grassland with many gaps 3^d
- Open slips/rockland and/or light, low-stature shrubland/tussock/grassland 4^d

0

TOTAL SCORE 0

Site 7 -Wilding Risk Calculation

1. SPECIES – GROWTH (score for one species only)

Spreading vigour varies with species

- Redwoods, Leyland cypresses, cedars and spruces (very low risk – no need to proceed further) 0
- Radiata (*P. radiata*) and ponderosa (*P. ponderosa*) pine, Lawsons cypress (*C. lawsoniana*) 1
- Muricata (*P. muricata*) and maritime (*P. pinaster*) pine and larches (*Larix* spp) 2
- Corsican (*P. nigra*) and mountain/dwarf mountain (*P. uncinata/mugo*) pine 3
- Douglas-fir ^g (*Ps. menziesii*), Scots pine (*P. sylvestris*) 4
- Lodgepole/contorta pine (*P. contorta*) 5

1

Enter score (0, 1, 2, 3, 4 or 5) here

2. SPECIES – PALATABILITY

Palatability varies with species

- Radiata, maritime and ponderosa pine 1
- Lodgepole and muricata pine and European larch 2
- Scots and mountain/dwarf mountain pine and Douglas-fir 3
- Corsican pine 4

1

Enter score (0, 1, 2, 3 or 4) here

3. SITING OF NEW PLANTING ^{c, d}

Trees are located on

- Sites well sheltered from prevalent and strong winds 0
- Flat sites (<10°), partially exposed to strong/prevalent winds 1
- Lea slopes where strong eddy gusts are likely 2
- Flat sites (<10°), fully exposed to strong/prevalent winds 3
- Either elevated 'take-off' sites, (ridge-tops, or base of exposed slopes >10°) or sloping land, fully exposed to strong/prevalent winds 4

4

Enter score (0, 1, 2, 3 or 4) here

4. DOWNWIND LANDUSE – GRAZING

Wilding establishment influenced by grazing (particularly with sheep)

- Intensive grazing on developed pasture 0^d
- Regular mob stocking with sheep ^e 1^d
- Semi-improved grazing (sheep/cattle)/ occasional mob stocking with sheep 2^d
- Extensive grazing only ^e 3^d
- No grazing 4^d

4

Enter score (0, 1, 2, 3 or 4) here

5. DOWNWIND VEGETATION COVER (if Douglas-fir involved see ^g in Notes)

Wilding establishment influenced by competition from existing vegetation

- Plantation forest, developed pasture (intensive grazing) 0^d
- Native forest^h, shrubland/tussock/grassland with a continuous and dense vegetation cover 1^d
- Forest/shrubland/tussock/grassland with few gaps 2^d
- Open forest and/or scattered patches of dense shrubland/tussock/grassland with many gaps 3^d
- Open slips/rockland and/or light, low-stature shrubland/tussock/grassland 4^d

0

TOTAL SCORE 0

Site 8 -Wilding Risk Calculation

1. SPECIES – GROWTH (score for one species only)

Spreading vigour varies with species

- Redwoods, Leyland cypresses, cedars and spruces (very low risk – no need to proceed further) 0
- Radiata (*P. radiata*) and ponderosa (*P. ponderosa*) pine, Lawsons cypress (*C. lawsoniana*) 1
- Muricata (*P. muricata*) and maritime (*P. pinaster*) pine and larches (*Larix* spp) 2
- Corsican (*P. nigra*) and mountain/dwarf mountain (*P. uncinata/mugo*) pine 3
- Douglas-fir ^g (*Ps. menziesii*), Scots pine (*P. sylvestris*) 4
- Lodgepole/contorta pine (*P. contorta*) 5

1

Enter score (0, 1, 2, 3, 4 or 5) here

2. SPECIES – PALATABILITY

Palatability varies with species

- Radiata, maritime and ponderosa pine 1
- Lodgepole and muricata pine and European larch 2
- Scots and mountain/dwarf mountain pine and Douglas-fir 3
- Corsican pine 4

1

Enter score (0, 1, 2, 3 or 4) here

3. SITING OF NEW PLANTING ^{c, d}

Trees are located on

- Sites well sheltered from prevalent and strong winds 0
- Flat sites (<10°), partially exposed to strong/prevalent winds 1
- Lea slopes where strong eddy gusts are likely 2
- Flat sites (<10°), fully exposed to strong/prevalent winds 3
- Either elevated 'take-off' sites, (ridge-tops, or base of exposed slopes >10°) or sloping land, fully exposed to strong/prevalent winds 4

4

Enter score (0, 1, 2, 3 or 4) here

4. DOWNWIND LANDUSE – GRAZING

Wilding establishment influenced by grazing (particularly with sheep)

- Intensive grazing on developed pasture 0^d
- Regular mob stocking with sheep ^e 1^d
- Semi-improved grazing (sheep/cattle)/ occasional mob stocking with sheep 2^d
- Extensive grazing only ^e 3^d
- No grazing 4^d

4

Enter score (0, 1, 2, 3 or 4) here

5. DOWNWIND VEGETATION COVER (if Douglas-fir involved see ^g in Notes)

Wilding establishment influenced by competition from existing vegetation

- Plantation forest, developed pasture (intensive grazing) 0^d
- Native forest^h, shrubland/tussock/grassland with a continuous and dense vegetation cover 1^d
- Forest/shrubland/tussock/grassland with few gaps 2^d
- Open forest and/or scattered patches of dense shrubland/tussock/grassland with many gaps 3^d
- Open slips/rockland and/or light, low-stature shrubland/tussock/grassland 4^d

0

TOTAL SCORE 0

Site 9 -Wilding Risk Calculation

1. SPECIES – GROWTH (score for one species only)

Spreading vigour varies with species

- Redwoods, Leyland cypresses, cedars and spruces (very low risk – no need to proceed further) 0
- Radiata (*P. radiata*) and ponderosa (*P. ponderosa*) pine, Lawsons cypress (*C. lawsoniana*) 1
- Muricata (*P. muricata*) and maritime (*P. pinaster*) pine and larches (*Larix* spp) 2
- Corsican (*P. nigra*) and mountain/dwarf mountain (*P. uncinata/mugo*) pine 3
- Douglas-fir ^g (*Ps. menziesii*), Scots pine (*P. sylvestris*) 4
- Lodgepole/contorta pine (*P. contorta*) 5

1

Enter score (0, 1, 2, 3, 4 or 5) here

2. SPECIES – PALATABILITY

Palatability varies with species

- Radiata, maritime and ponderosa pine 1
- Lodgepole and muricata pine and European larch 2
- Scots and mountain/dwarf mountain pine and Douglas-fir 3
- Corsican pine 4

1

Enter score (0, 1, 2, 3 or 4) here

3. SITING OF NEW PLANTING ^{c, d}

Trees are located on

- Sites well sheltered from prevalent and strong winds 0
- Flat sites (<10°), partially exposed to strong/prevalent winds 1
- Lea slopes where strong eddy gusts are likely 2
- Flat sites (<10°), fully exposed to strong/prevalent winds 3
- Either elevated 'take-off' sites, (ridge-tops, or base of exposed slopes >10°) or sloping land, fully exposed to strong/prevalent winds 4

4

Enter score (0, 1, 2, 3 or 4) here

4. DOWNWIND LANDUSE – GRAZING

Wilding establishment influenced by grazing (particularly with sheep)

- Intensive grazing on developed pasture 0^d
- Regular mob stocking with sheep ^e 1^d
- Semi-improved grazing (sheep/cattle)/ occasional mob stocking with sheep 2^d
- Extensive grazing only ^e 3^d
- No grazing 4^d

4

Enter score (0, 1, 2, 3 or 4) here

5. DOWNWIND VEGETATION COVER (if Douglas-fir involved see ^g in Notes)

Wilding establishment influenced by competition from existing vegetation

- Plantation forest, developed pasture (intensive grazing) 0^d
- Native forest^h, shrubland/tussock/grassland with a continuous and dense vegetation cover 1^d
- Forest/shrubland/tussock/grassland with few gaps 2^d
- Open forest and/or scattered patches of dense shrubland/tussock/grassland with many gaps 3^d
- Open slips/rockland and/or light, low-stature shrubland/tussock/grassland 4^d

0

TOTAL SCORE 0

Site 10 -Wilding Risk Calculation

1. SPECIES – GROWTH (score for one species only)

Spreading vigour varies with species

- Redwoods, Leyland cypresses, cedars and spruces (very low risk – no need to proceed further) 0
- Radiata (*P. radiata*) and ponderosa (*P. ponderosa*) pine, Lawsons cypress (*C. lawsoniana*) 1
- Muricata (*P. muricata*) and maritime (*P. pinaster*) pine and larches (*Larix* spp) 2
- Corsican (*P. nigra*) and mountain/dwarf mountain (*P. uncinata/mugo*) pine 3
- Douglas-fir ^g (*Ps. menziesii*), Scots pine (*P. sylvestris*) 4
- Lodgepole/contorta pine (*P. contorta*) 5

1

Enter score (0, 1, 2, 3, 4 or 5) here

2. SPECIES – PALATABILITY

Palatability varies with species

- Radiata, maritime and ponderosa pine 1
- Lodgepole and muricata pine and European larch 2
- Scots and mountain/dwarf mountain pine and Douglas-fir 3
- Corsican pine 4

1

Enter score (0, 1, 2, 3 or 4) here

3. SITING OF NEW PLANTING ^{c, d}

Trees are located on

- Sites well sheltered from prevalent and strong winds 0
- Flat sites (<10°), partially exposed to strong/prevalent winds 1
- Lea slopes where strong eddy gusts are likely 2
- Flat sites (<10°), fully exposed to strong/prevalent winds 3
- Either elevated 'take-off' sites, (ridge-tops, or base of exposed slopes >10°) or sloping land, fully exposed to strong/prevalent winds 4

4

Enter score (0, 1, 2, 3 or 4) here

4. DOWNWIND LANDUSE – GRAZING

Wilding establishment influenced by grazing (particularly with sheep)

- Intensive grazing on developed pasture 0^d
- Regular mob stocking with sheep ^e 1^d
- Semi-improved grazing (sheep/cattle)/ occasional mob stocking with sheep 2^d
- Extensive grazing only ^e 3^d
- No grazing 4^d

4

Enter score (0, 1, 2, 3 or 4) here

5. DOWNWIND VEGETATION COVER (if Douglas-fir involved see ^g in Notes)

Wilding establishment influenced by competition from existing vegetation

- Plantation forest, developed pasture (intensive grazing) 0^d
- Native forest^h, shrubland/tussock/grassland with a continuous and dense vegetation cover 1^d
- Forest/shrubland/tussock/grassland with few gaps 2^d
- Open forest and/or scattered patches of dense shrubland/tussock/grassland with many gaps 3^d
- Open slips/rockland and/or light, low-stature shrubland/tussock/grassland 4^d

0

TOTAL SCORE 0

Site 11 -Wilding Risk Calculation

1. SPECIES – GROWTH (score for one species only)

Spreading vigour varies with species

- Redwoods, Leyland cypresses, cedars and spruces (very low risk – no need to proceed further) 0
- Radiata (*P. radiata*) and ponderosa (*P. ponderosa*) pine, Lawsons cypress (*C. lawsoniana*) 1
- Muricata (*P. muricata*) and maritime (*P. pinaster*) pine and larches (*Larix* spp) 2
- Corsican (*P. nigra*) and mountain/dwarf mountain (*P. uncinata/mugo*) pine 3
- Douglas-fir ^g (*Ps. menziesii*), Scots pine (*P. sylvestris*) 4
- Lodgepole/contorta pine (*P. contorta*) 5

1

Enter score (0, 1, 2, 3, 4 or 5) here

2. SPECIES – PALATABILITY

Palatability varies with species

- Radiata, maritime and ponderosa pine 1
- Lodgepole and muricata pine and European larch 2
- Scots and mountain/dwarf mountain pine and Douglas-fir 3
- Corsican pine 4

1

Enter score (0, 1, 2, 3 or 4) here

3. SITING OF NEW PLANTING ^{c, d}

Trees are located on

- Sites well sheltered from prevalent and strong winds 0
- Flat sites (<10°), partially exposed to strong/prevalent winds 1
- Lea slopes where strong eddy gusts are likely 2
- Flat sites (<10°), fully exposed to strong/prevalent winds 3
- Either elevated 'take-off' sites, (ridge-tops, or base of exposed slopes >10°) or sloping land, fully exposed to strong/prevalent winds 4

4

Enter score (0, 1, 2, 3 or 4) here

4. DOWNWIND LANDUSE – GRAZING

Wilding establishment influenced by grazing (particularly with sheep)

- Intensive grazing on developed pasture 0^d
- Regular mob stocking with sheep ^e 1^d
- Semi-improved grazing (sheep/cattle)/ occasional mob stocking with sheep 2^d
- Extensive grazing only ^e 3^d
- No grazing 4^d

4

Enter score (0, 1, 2, 3 or 4) here

5. DOWNWIND VEGETATION COVER (if Douglas-fir involved see ^g in Notes)

Wilding establishment influenced by competition from existing vegetation

- Plantation forest, developed pasture (intensive grazing) 0^d
- Native forest^h, shrubland/tussock/grassland with a continuous and dense vegetation cover 1^d
- Forest/shrubland/tussock/grassland with few gaps 2^d
- Open forest and/or scattered patches of dense shrubland/tussock/grassland with many gaps 3^d
- Open slips/rockland and/or light, low-stature shrubland/tussock/grassland 4^d

0

TOTAL SCORE 0

Site 12 -Wilding Risk Calculation

1. SPECIES – GROWTH (score for one species only)

Spreading vigour varies with species

- Redwoods, Leyland cypresses, cedars and spruces (very low risk – no need to proceed further) 0
- Radiata (*P. radiata*) and ponderosa (*P. ponderosa*) pine, Lawsons cypress (*C. lawsoniana*) 1
- Muricata (*P. muricata*) and maritime (*P. pinaster*) pine and larches (*Larix* spp) 2
- Corsican (*P. nigra*) and mountain/dwarf mountain (*P. uncinata/mugo*) pine 3
- Douglas-fir ^g (*Ps. menziesii*), Scots pine (*P. sylvestris*) 4
- Lodgepole/contorta pine (*P. contorta*) 5

1

Enter score (0, 1, 2, 3, 4 or 5) here

2. SPECIES – PALATABILITY

Palatability varies with species

- Radiata, maritime and ponderosa pine 1
- Lodgepole and muricata pine and European larch 2
- Scots and mountain/dwarf mountain pine and Douglas-fir 3
- Corsican pine 4

1

Enter score (0, 1, 2, 3 or 4) here

3. SITING OF NEW PLANTING ^{c, d}

Trees are located on

- Sites well sheltered from prevalent and strong winds 0
- Flat sites (<10°), partially exposed to strong/prevalent winds 1
- Lea slopes where strong eddy gusts are likely 2
- Flat sites (<10°), fully exposed to strong/prevalent winds 3
- Either elevated 'take-off' sites, (ridge-tops, or base of exposed slopes >10°) or sloping land, fully exposed to strong/prevalent winds 4

4

Enter score (0, 1, 2, 3 or 4) here

4. DOWNWIND LANDUSE – GRAZING

Wilding establishment influenced by grazing (particularly with sheep)

- Intensive grazing on developed pasture 0^d
- Regular mob stocking with sheep ^e 1^d
- Semi-improved grazing (sheep/cattle)/ occasional mob stocking with sheep 2^d
- Extensive grazing only ^e 3^d
- No grazing 4^d

0

Enter score (0, 1, 2, 3 or 4) here

5. DOWNWIND VEGETATION COVER (if Douglas-fir involved see ^g in Notes)

Wilding establishment influenced by competition from existing vegetation

- Plantation forest, developed pasture (intensive grazing) 0^d
- Native forest^h, shrubland/tussock/grassland with a continuous and dense vegetation cover 1^d
- Forest/shrubland/tussock/grassland with few gaps 2^d
- Open forest and/or scattered patches of dense shrubland/tussock/grassland with many gaps 3^d
- Open slips/rockland and/or light, low-stature shrubland/tussock/grassland 4^d

0

TOTAL SCORE 0

Site 13 -Wilding Risk Calculation

1. SPECIES – GROWTH (score for one species only)

Spreading vigour varies with species

- Redwoods, Leyland cypresses, cedars and spruces (very low risk – no need to proceed further) 0
- Radiata (*P. radiata*) and ponderosa (*P. ponderosa*) pine, Lawsons cypress (*C. lawsoniana*) 1
- Muricata (*P. muricata*) and maritime (*P. pinaster*) pine and larches (*Larix* spp) 2
- Corsican (*P. nigra*) and mountain/dwarf mountain (*P. uncinata/mugo*) pine 3
- Douglas-fir ^g (*Ps. menziesii*), Scots pine (*P. sylvestris*) 4
- Lodgepole/contorta pine (*P. contorta*) 5

1

Enter score (0, 1, 2, 3, 4 or 5) here

2. SPECIES – PALATABILITY

Palatability varies with species

- Radiata, maritime and ponderosa pine 1
- Lodgepole and muricata pine and European larch 2
- Scots and mountain/dwarf mountain pine and Douglas-fir 3
- Corsican pine 4

1

Enter score (0, 1, 2, 3 or 4) here

3. SITING OF NEW PLANTING ^{c, d}

Trees are located on

- Sites well sheltered from prevalent and strong winds 0
- Flat sites (<10°), partially exposed to strong/prevalent winds 1
- Lea slopes where strong eddy gusts are likely 2
- Flat sites (<10°), fully exposed to strong/prevalent winds 3
- Either elevated 'take-off' sites, (ridge-tops, or base of exposed slopes >10°) or sloping land, fully exposed to strong/prevalent winds 4

4

Enter score (0, 1, 2, 3 or 4) here

4. DOWNWIND LANDUSE – GRAZING

Wilding establishment influenced by grazing (particularly with sheep)

- Intensive grazing on developed pasture 0^d
- Regular mob stocking with sheep ^e 1^d
- Semi-improved grazing (sheep/cattle)/ occasional mob stocking with sheep 2^d
- Extensive grazing only ^e 3^d
- No grazing 4^d

0

Enter score (0, 1, 2, 3 or 4) here

5. DOWNWIND VEGETATION COVER (if Douglas-fir involved see ^g in Notes)

Wilding establishment influenced by competition from existing vegetation

- Plantation forest, developed pasture (intensive grazing) 0^d
- Native forest^h, shrubland/tussock/grassland with a continuous and dense vegetation cover 1^d
- Forest/shrubland/tussock/grassland with few gaps 2^d
- Open forest and/or scattered patches of dense shrubland/tussock/grassland with many gaps 3^d
- Open slips/rockland and/or light, low-stature shrubland/tussock/grassland 4^d

0

TOTAL SCORE 0

Site 14 -Wilding Risk Calculation

1. SPECIES – GROWTH (score for one species only)

Spreading vigour varies with species

- Redwoods, Leyland cypresses, cedars and spruces (very low risk – no need to proceed further) 0
- Radiata (*P. radiata*) and ponderosa (*P. ponderosa*) pine, Lawsons cypress (*C. lawsoniana*) 1
- Muricata (*P. muricata*) and maritime (*P. pinaster*) pine and larches (*Larix* spp) 2
- Corsican (*P. nigra*) and mountain/dwarf mountain (*P. uncinata/mugo*) pine 3
- Douglas-fir ^g (*Ps. menziesii*), Scots pine (*P. sylvestris*) 4
- Lodgepole/contorta pine (*P. contorta*) 5

1

Enter score (0, 1, 2, 3, 4 or 5) here

2. SPECIES – PALATABILITY

Palatability varies with species

- Radiata, maritime and ponderosa pine 1
- Lodgepole and muricata pine and European larch 2
- Scots and mountain/dwarf mountain pine and Douglas-fir 3
- Corsican pine 4

1

Enter score (0, 1, 2, 3 or 4) here

3. SITING OF NEW PLANTING ^{c, d}

Trees are located on

- Sites well sheltered from prevalent and strong winds 0
- Flat sites (<10°), partially exposed to strong/prevalent winds 1
- Lea slopes where strong eddy gusts are likely 2
- Flat sites (<10°), fully exposed to strong/prevalent winds 3
- Either elevated 'take-off' sites, (ridge-tops, or base of exposed slopes >10°) or sloping land, fully exposed to strong/prevalent winds 4

1

Enter score (0, 1, 2, 3 or 4) here

4. DOWNWIND LANDUSE – GRAZING

Wilding establishment influenced by grazing (particularly with sheep)

- Intensive grazing on developed pasture 0^d
- Regular mob stocking with sheep ^e 1^d
- Semi-improved grazing (sheep/cattle)/ occasional mob stocking with sheep 2^d
- Extensive grazing only ^e 3^d
- No grazing 4^d

4

Enter score (0, 1, 2, 3 or 4) here

5. DOWNWIND VEGETATION COVER (if Douglas-fir involved see ^g in Notes)

Wilding establishment influenced by competition from existing vegetation

- Plantation forest, developed pasture (intensive grazing) 0^d
- Native forest^h, shrubland/tussock/grassland with a continuous and dense vegetation cover 1^d
- Forest/shrubland/tussock/grassland with few gaps 2^d
- Open forest and/or scattered patches of dense shrubland/tussock/grassland with many gaps 3^d
- Open slips/rockland and/or light, low-stature shrubland/tussock/grassland 4^d

0

TOTAL SCORE 0

Site 15 -Wilding Risk Calculation

1. SPECIES – GROWTH (score for one species only)

Spreading vigour varies with species

- Redwoods, Leyland cypresses, cedars and spruces (very low risk – no need to proceed further) 0
- Radiata (*P. radiata*) and ponderosa (*P. ponderosa*) pine, Lawsons cypress (*C. lawsoniana*) 1
- Muricata (*P. muricata*) and maritime (*P. pinaster*) pine and larches (*Larix* spp) 2
- Corsican (*P. nigra*) and mountain/dwarf mountain (*P. uncinata/mugo*) pine 3
- Douglas-fir ^g (*Ps. menziesii*), Scots pine (*P. sylvestris*) 4
- Lodgepole/contorta pine (*P. contorta*) 5

1

Enter score (0, 1, 2, 3, 4 or 5) here

2. SPECIES – PALATABILITY

Palatability varies with species

- Radiata, maritime and ponderosa pine 1
- Lodgepole and muricata pine and European larch 2
- Scots and mountain/dwarf mountain pine and Douglas-fir 3
- Corsican pine 4

1

Enter score (0, 1, 2, 3 or 4) here

3. SITING OF NEW PLANTING ^{c, d}

Trees are located on

- Sites well sheltered from prevalent and strong winds 0
- Flat sites (<10°), partially exposed to strong/prevalent winds 1
- Lea slopes where strong eddy gusts are likely 2
- Flat sites (<10°), fully exposed to strong/prevalent winds 3
- Either elevated 'take-off' sites, (ridge-tops, or base of exposed slopes >10°) or sloping land, fully exposed to strong/prevalent winds 4

4

Enter score (0, 1, 2, 3 or 4) here

4. DOWNWIND LANDUSE – GRAZING

Wilding establishment influenced by grazing (particularly with sheep)

- Intensive grazing on developed pasture 0^d
- Regular mob stocking with sheep ^e 1^d
- Semi-improved grazing (sheep/cattle)/ occasional mob stocking with sheep 2^d
- Extensive grazing only ^e 3^d
- No grazing 4^d

4

Enter score (0, 1, 2, 3 or 4) here

5. DOWNWIND VEGETATION COVER (if Douglas-fir involved see ^g in Notes)

Wilding establishment influenced by competition from existing vegetation

Plantation forest, developed pasture (intensive grazing) 0^d

- Native forest^h, shrubland/tussock/grassland with a continuous and dense vegetation cover 1^d
- Forest/shrubland/tussock/grassland with few gaps 2^d
- Open forest and/or scattered patches of dense shrubland/tussock/grassland with many gaps 3^d
- Open slips/rockland and/or light, low-stature shrubland/tussock/grassland 4^d

0

TOTAL SCORE 0

Site 16 -Wilding Risk Calculation

1. SPECIES – GROWTH (score for one species only)

Spreading vigour varies with species

- Redwoods, Leyland cypresses, cedars and spruces (very low risk – no need to proceed further) 0
- Radiata (*P. radiata*) and ponderosa (*P. ponderosa*) pine, Lawsons cypress (*C. lawsoniana*) 1
- Muricata (*P. muricata*) and maritime (*P. pinaster*) pine and larches (*Larix* spp) 2
- Corsican (*P. nigra*) and mountain/dwarf mountain (*P. uncinata/mugo*) pine 3
- Douglas-fir ^g (*Ps. menziesii*), Scots pine (*P. sylvestris*) 4
- Lodgepole/contorta pine (*P. contorta*) 5

1

Enter score (0, 1, 2, 3, 4 or 5) here

2. SPECIES – PALATABILITY

Palatability varies with species

- Radiata, maritime and ponderosa pine 1
- Lodgepole and muricata pine and European larch 2
- Scots and mountain/dwarf mountain pine and Douglas-fir 3
- Corsican pine 4

1

Enter score (0, 1, 2, 3 or 4) here

3. SITING OF NEW PLANTING ^{c, d}

Trees are located on

- Sites well sheltered from prevalent and strong winds 0
- Flat sites (<10°), partially exposed to strong/prevalent winds 1
- Lea slopes where strong eddy gusts are likely 2
- Flat sites (<10°), fully exposed to strong/prevalent winds 3
- Either elevated 'take-off' sites, (ridge-tops, or base of exposed slopes >10°) or sloping land, fully exposed to strong/prevalent winds 4

0

Enter score (0, 1, 2, 3 or 4) here

4. DOWNWIND LANDUSE – GRAZING

Wilding establishment influenced by grazing (particularly with sheep)

- Intensive grazing on developed pasture 0^d
- Regular mob stocking with sheep ^e 1^d
- Semi-improved grazing (sheep/cattle)/ occasional mob stocking with sheep 2^d
- Extensive grazing only ^e 3^d
- No grazing 4^d

4

Enter score (0, 1, 2, 3 or 4) here

5. DOWNWIND VEGETATION COVER (if Douglas-fir involved see ^g in Notes)

Wilding establishment influenced by competition from existing vegetation

- Plantation forest, developed pasture (intensive grazing) 0^d
- Native forest^h, shrubland/tussock/grassland with a continuous and dense vegetation cover 1^d
- Forest/shrubland/tussock/grassland with few gaps 2^d
- Open forest and/or scattered patches of dense shrubland/tussock/grassland with many gaps 3^d
- Open slips/rockland and/or light, low-stature shrubland/tussock/grassland 4^d

0

TOTAL SCORE 0

Site 17 -Wilding Risk Calculation

1. SPECIES – GROWTH (score for one species only)

Spreading vigour varies with species

- Redwoods, Leyland cypresses, cedars and spruces (very low risk – no need to proceed further) 0
- Radiata (*P. radiata*) and ponderosa (*P. ponderosa*) pine, Lawsons cypress (*C. lawsoniana*) 1
- Muricata (*P. muricata*) and maritime (*P. pinaster*) pine and larches (*Larix* spp) 2
- Corsican (*P. nigra*) and mountain/dwarf mountain (*P. uncinata/mugo*) pine 3
- Douglas-fir ^g (*Ps. menziesii*), Scots pine (*P. sylvestris*) 4
- Lodgepole/contorta pine (*P. contorta*) 5

1

Enter score (0, 1, 2, 3, 4 or 5) here

2. SPECIES – PALATABILITY

Palatability varies with species

- Radiata, maritime and ponderosa pine 1
- Lodgepole and muricata pine and European larch 2
- Scots and mountain/dwarf mountain pine and Douglas-fir 3
- Corsican pine 4

1

Enter score (0, 1, 2, 3 or 4) here

3. SITING OF NEW PLANTING ^{c, d}

Trees are located on

- Sites well sheltered from prevalent and strong winds 0
- Flat sites (<10°), partially exposed to strong/prevalent winds 1
- Lea slopes where strong eddy gusts are likely 2
- Flat sites (<10°), fully exposed to strong/prevalent winds 3
- Either elevated 'take-off' sites, (ridge-tops, or base of exposed slopes >10°) or sloping land, fully exposed to strong/prevalent winds 4

1

Enter score (0, 1, 2, 3 or 4) here

4. DOWNWIND LANDUSE – GRAZING

Wilding establishment influenced by grazing (particularly with sheep)

- Intensive grazing on developed pasture 0^d
- Regular mob stocking with sheep ^e 1^d
- Semi-improved grazing (sheep/cattle)/ occasional mob stocking with sheep 2^d
- Extensive grazing only ^e 3^d
- No grazing 4^d

4

Enter score (0, 1, 2, 3 or 4) here

5. DOWNWIND VEGETATION COVER (if Douglas-fir involved see ^g in Notes)

Wilding establishment influenced by competition from existing vegetation

- Plantation forest, developed pasture (intensive grazing) 0^d
- Native forest^h, shrubland/tussock/grassland with a continuous and dense vegetation cover 1^d
- Forest/shrubland/tussock/grassland with few gaps 2^d
- Open forest and/or scattered patches of dense shrubland/tussock/grassland with many gaps 3^d
- Open slips/rockland and/or light, low-stature shrubland/tussock/grassland 4^d

0

TOTAL SCORE 0

Site 18 -Wilding Risk Calculation

1. SPECIES – GROWTH (score for one species only)

Spreading vigour varies with species

- Redwoods, Leyland cypresses, cedars and spruces (very low risk – no need to proceed further) 0
- Radiata (*P. radiata*) and ponderosa (*P. ponderosa*) pine, Lawsons cypress (*C. lawsoniana*) 1
- Muricata (*P. muricata*) and maritime (*P. pinaster*) pine and larches (*Larix* spp) 2
- Corsican (*P. nigra*) and mountain/dwarf mountain (*P. uncinata/mugo*) pine 3
- Douglas-fir ^g (*Ps. menziesii*), Scots pine (*P. sylvestris*) 4
- Lodgepole/contorta pine (*P. contorta*) 5

1

Enter score (0, 1, 2, 3, 4 or 5) here

2. SPECIES – PALATABILITY

Palatability varies with species

- Radiata, maritime and ponderosa pine 1
- Lodgepole and muricata pine and European larch 2
- Scots and mountain/dwarf mountain pine and Douglas-fir 3
- Corsican pine 4

1

Enter score (0, 1, 2, 3 or 4) here

3. SITING OF NEW PLANTING ^{c, d}

Trees are located on

- Sites well sheltered from prevalent and strong winds 0
- Flat sites (<10°), partially exposed to strong/prevalent winds 1
- Lea slopes where strong eddy gusts are likely 2
- Flat sites (<10°), fully exposed to strong/prevalent winds 3
- Either elevated 'take-off' sites, (ridge-tops, or base of exposed slopes >10°) or sloping land, fully exposed to strong/prevalent winds 4

1

Enter score (0, 1, 2, 3 or 4) here

4. DOWNWIND LANDUSE – GRAZING

Wilding establishment influenced by grazing (particularly with sheep)

- Intensive grazing on developed pasture 0^d
- Regular mob stocking with sheep ^e 1^d
- Semi-improved grazing (sheep/cattle)/ occasional mob stocking with sheep 2^d
- Extensive grazing only ^e 3^d
- No grazing 4^d

4

Enter score (0, 1, 2, 3 or 4) here

5. DOWNWIND VEGETATION COVER (if Douglas-fir involved see ^g in Notes)

Wilding establishment influenced by competition from existing vegetation

- Plantation forest, developed pasture (intensive grazing) 0^d
- Native forest^h, shrubland/tussock/grassland with a continuous and dense vegetation cover 1^d
- Forest/shrubland/tussock/grassland with few gaps 2^d
- Open forest and/or scattered patches of dense shrubland/tussock/grassland with many gaps 3^d
- Open slips/rockland and/or light, low-stature shrubland/tussock/grassland 4^d

0

TOTAL SCORE 0

Site 19 -Wilding Risk Calculation

1. SPECIES – GROWTH (score for one species only)

Spreading vigour varies with species

- Redwoods, Leyland cypresses, cedars and spruces (very low risk – no need to proceed further) 0
- Radiata (*P. radiata*) and ponderosa (*P. ponderosa*) pine, Lawsons cypress (*C. lawsoniana*) 1
- Muricata (*P. muricata*) and maritime (*P. pinaster*) pine and larches (*Larix* spp) 2
- Corsican (*P. nigra*) and mountain/dwarf mountain (*P. uncinata/mugo*) pine 3
- Douglas-fir ^g (*Ps. menziesii*), Scots pine (*P. sylvestris*) 4
- Lodgepole/contorta pine (*P. contorta*) 5

1

Enter score (0, 1, 2, 3, 4 or 5) here

2. SPECIES – PALATABILITY

Palatability varies with species

- Radiata, maritime and ponderosa pine 1
- Lodgepole and muricata pine and European larch 2
- Scots and mountain/dwarf mountain pine and Douglas-fir 3
- Corsican pine 4

1

Enter score (0, 1, 2, 3 or 4) here

3. SITING OF NEW PLANTING ^{c, d}

Trees are located on

- Sites well sheltered from prevalent and strong winds 0
- Flat sites (<10°), partially exposed to strong/prevalent winds 1
- Lea slopes where strong eddy gusts are likely 2
- Flat sites (<10°), fully exposed to strong/prevalent winds 3
- Either elevated 'take-off' sites, (ridge-tops, or base of exposed slopes >10°) or sloping land, fully exposed to strong/prevalent winds 4

4

Enter score (0, 1, 2, 3 or 4) here

4. DOWNWIND LANDUSE – GRAZING

Wilding establishment influenced by grazing (particularly with sheep)

- Intensive grazing on developed pasture 0^d
- Regular mob stocking with sheep ^e 1^d
- Semi-improved grazing (sheep/cattle)/ occasional mob stocking with sheep 2^d
- Extensive grazing only ^e 3^d
- No grazing 4^d

4

Enter score (0, 1, 2, 3 or 4) here

5. DOWNWIND VEGETATION COVER (if Douglas-fir involved see ^g in Notes)

Wilding establishment influenced by competition from existing vegetation

- Plantation forest, developed pasture (intensive grazing) 0^d
- Native forest^h, shrubland/tussock/grassland with a continuous and dense vegetation cover 1^d
- Forest/shrubland/tussock/grassland with few gaps 2^d
- Open forest and/or scattered patches of dense shrubland/tussock/grassland with many gaps 3^d
- Open slips/rockland and/or light, low-stature shrubland/tussock/grassland 4^d

0

TOTAL SCORE 0

Site 20-Wilding Risk Calculation

1. SPECIES – GROWTH (score for one species only)

Spreading vigour varies with species

- Redwoods, Leyland cypresses, cedars and spruces (very low risk – no need to proceed further) 0
- Radiata (*P. radiata*) and ponderosa (*P. ponderosa*) pine, Lawsons cypress (*C. lawsoniana*) 1
- Muricata (*P. muricata*) and maritime (*P. pinaster*) pine and larches (*Larix* spp) 2
- Corsican (*P. nigra*) and mountain/dwarf mountain (*P. uncinata/mugo*) pine 3
- Douglas-fir ^g (*Ps. menziesii*), Scots pine (*P. sylvestris*) 4
- Lodgepole/contorta pine (*P. contorta*) 5

1

Enter score (0, 1, 2, 3, 4 or 5) here

2. SPECIES – PALATABILITY

Palatability varies with species

- Radiata, maritime and ponderosa pine 1
- Lodgepole and muricata pine and European larch 2
- Scots and mountain/dwarf mountain pine and Douglas-fir 3
- Corsican pine 4

1

Enter score (0, 1, 2, 3 or 4) here

3. SITING OF NEW PLANTING ^{c, d}

Trees are located on

- Sites well sheltered from prevalent and strong winds 0
- Flat sites (<10°), partially exposed to strong/prevalent winds 1
- Lea slopes where strong eddy gusts are likely 2
- Flat sites (<10°), fully exposed to strong/prevalent winds 3
- Either elevated 'take-off' sites, (ridge-tops, or base of exposed slopes >10°) or sloping land, fully exposed to strong/prevalent winds 4

4

Enter score (0, 1, 2, 3 or 4) here

4. DOWNWIND LANDUSE – GRAZING

Wilding establishment influenced by grazing (particularly with sheep)

- Intensive grazing on developed pasture 0^d
- Regular mob stocking with sheep ^e 1^d
- Semi-improved grazing (sheep/cattle)/ occasional mob stocking with sheep 2^d
- Extensive grazing only ^e 3^d
- No grazing 4^d

4

Enter score (0, 1, 2, 3 or 4) here

5. DOWNWIND VEGETATION COVER (if Douglas-fir involved see ^g in Notes)

Wilding establishment influenced by competition from existing vegetation

- Plantation forest, developed pasture (intensive grazing) 0^d
- Native forest^h, shrubland/tussock/grassland with a continuous and dense vegetation cover 1^d
- Forest/shrubland/tussock/grassland with few gaps 2^d
- Open forest and/or scattered patches of dense shrubland/tussock/grassland with many gaps 3^d
- Open slips/rockland and/or light, low-stature shrubland/tussock/grassland 4^d

0

TOTAL SCORE 0

Site 21 -Wilding Risk Calculation

1. SPECIES – GROWTH (score for one species only)

Spreading vigour varies with species

- Redwoods, Leyland cypresses, cedars and spruces (very low risk – no need to proceed further) 0
- Radiata (*P. radiata*) and ponderosa (*P. ponderosa*) pine, Lawsons cypress (*C. lawsoniana*) 1
- Muricata (*P. muricata*) and maritime (*P. pinaster*) pine and larches (*Larix* spp) 2
- Corsican (*P. nigra*) and mountain/dwarf mountain (*P. uncinata/mugo*) pine 3
- Douglas-fir ^g (*Ps. menziesii*), Scots pine (*P. sylvestris*) 4
- Lodgepole/contorta pine (*P. contorta*) 5

1

Enter score (0, 1, 2, 3, 4 or 5) here

2. SPECIES – PALATABILITY

Palatability varies with species

- Radiata, maritime and ponderosa pine 1
- Lodgepole and muricata pine and European larch 2
- Scots and mountain/dwarf mountain pine and Douglas-fir 3
- Corsican pine 4

1

Enter score (0, 1, 2, 3 or 4) here

3. SITING OF NEW PLANTING ^{c, d}

Trees are located on

- Sites well sheltered from prevalent and strong winds 0
- Flat sites (<10°), partially exposed to strong/prevalent winds 1
- Lea slopes where strong eddy gusts are likely 2
- Flat sites (<10°), fully exposed to strong/prevalent winds 3
- Either elevated ‘take-off’ sites, (ridge-tops, or base of exposed slopes >10°) or sloping land, fully exposed to strong/prevalent winds 4

4

Enter score (0, 1, 2, 3 or 4) here

4. DOWNWIND LANDUSE – GRAZING

Wilding establishment influenced by grazing (particularly with sheep)

- Intensive grazing on developed pasture 0^d
- Regular mob stocking with sheep ^e 1^d
- Semi-improved grazing (sheep/cattle)/ occasional mob stocking with sheep 2^d
- Extensive grazing only ^e 3^d
- No grazing 4^d

4

Enter score (0, 1, 2, 3 or 4) here

5. DOWNWIND VEGETATION COVER (if Douglas-fir involved see ^g in Notes)

Wilding establishment influenced by competition from existing vegetation

- Plantation forest, developed pasture (intensive grazing) 0^d
- Native forest^h, shrubland/tussock/grassland with a continuous and dense vegetation cover 1^d
- Forest/shrubland/tussock/grassland with few gaps 2^d
- Open forest and/or scattered patches of dense shrubland/tussock/grassland with many gaps 3^d
- Open slips/rockland and/or light, low-stature shrubland/tussock/grassland 4^d

0

TOTAL SCORE 0

Site 22 -Wilding Risk Calculation

1. SPECIES – GROWTH (score for one species only)

Spreading vigour varies with species

- Redwoods, Leyland cypresses, cedars and spruces (very low risk – no need to proceed further) 0
- Radiata (*P. radiata*) and ponderosa (*P. ponderosa*) pine, Lawsons cypress (*C. lawsoniana*) 1
- Muricata (*P. muricata*) and maritime (*P. pinaster*) pine and larches (*Larix* spp) 2
- Corsican (*P. nigra*) and mountain/dwarf mountain (*P. uncinata/mugo*) pine 3
- Douglas-fir ^g (*Ps. menziesii*), Scots pine (*P. sylvestris*) 4
- Lodgepole/contorta pine (*P. contorta*) 5

1

Enter score (0, 1, 2, 3, 4 or 5) here

2. SPECIES – PALATABILITY

Palatability varies with species

- Radiata, maritime and ponderosa pine 1
- Lodgepole and muricata pine and European larch 2
- Scots and mountain/dwarf mountain pine and Douglas-fir 3
- Corsican pine 4

1

Enter score (0, 1, 2, 3 or 4) here

3. SITING OF NEW PLANTING ^{c, d}

Trees are located on

- Sites well sheltered from prevalent and strong winds 0
- Flat sites (<10°), partially exposed to strong/prevalent winds 1
- Lea slopes where strong eddy gusts are likely 2
- Flat sites (<10°), fully exposed to strong/prevalent winds 3
- Either elevated 'take-off' sites, (ridge-tops, or base of exposed slopes >10°) or sloping land, fully exposed to strong/prevalent winds 4

4

Enter score (0, 1, 2, 3 or 4) here

4. DOWNWIND LANDUSE – GRAZING

Wilding establishment influenced by grazing (particularly with sheep)

- Intensive grazing on developed pasture 0^d
- Regular mob stocking with sheep ^e 1^d
- Semi-improved grazing (sheep/cattle)/ occasional mob stocking with sheep 2^d
- Extensive grazing only ^e 3^d
- No grazing 4^d

4

Enter score (0, 1, 2, 3 or 4) here

5. DOWNWIND VEGETATION COVER (if Douglas-fir involved see ^g in Notes)

Wilding establishment influenced by competition from existing vegetation

- Plantation forest, developed pasture (intensive grazing) 0^d
- Native forest^h, shrubland/tussock/grassland with a continuous and dense vegetation cover 1^d
- Forest/shrubland/tussock/grassland with few gaps 2^d
- Open forest and/or scattered patches of dense shrubland/tussock/grassland with many gaps 3^d
- Open slips/rockland and/or light, low-stature shrubland/tussock/grassland 4^d

0

TOTAL SCORE 0

Site 23 -Wilding Risk Calculation

1. SPECIES – GROWTH (score for one species only)

Spreading vigour varies with species

- Redwoods, Leyland cypresses, cedars and spruces (very low risk – no need to proceed further) 0
- Radiata (*P. radiata*) and ponderosa (*P. ponderosa*) pine, Lawsons cypress (*C. lawsoniana*) 1
- Muricata (*P. muricata*) and maritime (*P. pinaster*) pine and larches (*Larix* spp) 2
- Corsican (*P. nigra*) and mountain/dwarf mountain (*P. uncinata/mugo*) pine 3
- Douglas-fir ^g (*Ps. menziesii*), Scots pine (*P. sylvestris*) 4
- Lodgepole/contorta pine (*P. contorta*) 5

1

Enter score (0, 1, 2, 3, 4 or 5) here

2. SPECIES – PALATABILITY

Palatability varies with species

- Radiata, maritime and ponderosa pine 1
- Lodgepole and muricata pine and European larch 2
- Scots and mountain/dwarf mountain pine and Douglas-fir 3
- Corsican pine 4

1

Enter score (0, 1, 2, 3 or 4) here

3. SITING OF NEW PLANTING ^{c, d}

Trees are located on

- Sites well sheltered from prevalent and strong winds 0
- Flat sites (<10°), partially exposed to strong/prevalent winds 1
- Lea slopes where strong eddy gusts are likely 2
- Flat sites (<10°), fully exposed to strong/prevalent winds 3
- Either elevated 'take-off' sites, (ridge-tops, or base of exposed slopes >10°) or sloping land, fully exposed to strong/prevalent winds 4

1

Enter score (0, 1, 2, 3 or 4) here

4. DOWNWIND LANDUSE – GRAZING

Wilding establishment influenced by grazing (particularly with sheep)

- Intensive grazing on developed pasture 0^d
- Regular mob stocking with sheep ^e 1^d
- Semi-improved grazing (sheep/cattle)/ occasional mob stocking with sheep 2^d
- Extensive grazing only ^e 3^d
- No grazing 4^d

4

Enter score (0, 1, 2, 3 or 4) here

5. DOWNWIND VEGETATION COVER (if Douglas-fir involved see ^g in Notes)

Wilding establishment influenced by competition from existing vegetation

- Plantation forest, developed pasture (intensive grazing) 0^d
- Native forest^h, shrubland/tussock/grassland with a continuous and dense vegetation cover 1^d
- Forest/shrubland/tussock/grassland with few gaps 2^d
- Open forest and/or scattered patches of dense shrubland/tussock/grassland with many gaps 3^d
- Open slips/rockland and/or light, low-stature shrubland/tussock/grassland 4^d

0

TOTAL SCORE 0

Site 24 -Wilding Risk Calculation

1. SPECIES – GROWTH (score for one species only)

Spreading vigour varies with species

- Redwoods, Leyland cypresses, cedars and spruces (very low risk – no need to proceed further) 0
- Radiata (*P. radiata*) and ponderosa (*P. ponderosa*) pine, Lawsons cypress (*C. lawsoniana*) 1
- Muricata (*P. muricata*) and maritime (*P. pinaster*) pine and larches (*Larix* spp) 2
- Corsican (*P. nigra*) and mountain/dwarf mountain (*P. uncinata/mugo*) pine 3
- Douglas-fir ^g (*Ps. menziesii*), Scots pine (*P. sylvestris*) 4
- Lodgepole/contorta pine (*P. contorta*) 5

1

Enter score (0, 1, 2, 3, 4 or 5) here

2. SPECIES – PALATABILITY

Palatability varies with species

- Radiata, maritime and ponderosa pine 1
- Lodgepole and muricata pine and European larch 2
- Scots and mountain/dwarf mountain pine and Douglas-fir 3
- Corsican pine 4

1

Enter score (0, 1, 2, 3 or 4) here

3. SITING OF NEW PLANTING ^{c, d}

Trees are located on

- Sites well sheltered from prevalent and strong winds 0
- Flat sites (<10°), partially exposed to strong/prevalent winds 1
- Lea slopes where strong eddy gusts are likely 2
- Flat sites (<10°), fully exposed to strong/prevalent winds 3
- Either elevated 'take-off' sites, (ridge-tops, or base of exposed slopes >10°) or sloping land, fully exposed to strong/prevalent winds 4

4

Enter score (0, 1, 2, 3 or 4) here

4. DOWNWIND LANDUSE – GRAZING

Wilding establishment influenced by grazing (particularly with sheep)

- Intensive grazing on developed pasture 0^d
- Regular mob stocking with sheep ^e 1^d
- Semi-improved grazing (sheep/cattle)/ occasional mob stocking with sheep 2^d
- Extensive grazing only ^e 3^d
- No grazing 4^d

4

Enter score (0, 1, 2, 3 or 4) here

5. DOWNWIND VEGETATION COVER (if Douglas-fir involved see ^g in Notes)

Wilding establishment influenced by competition from existing vegetation

- Plantation forest, developed pasture (intensive grazing) 0^d
- Native forest^h, shrubland/tussock/grassland with a continuous and dense vegetation cover 1^d
- Forest/shrubland/tussock/grassland with few gaps 2^d
- Open forest and/or scattered patches of dense shrubland/tussock/grassland with many gaps 3^d
- Open slips/rockland and/or light, low-stature shrubland/tussock/grassland 4^d

0

TOTAL SCORE 0

Site 25 -Wilding Risk Calculation

1. SPECIES – GROWTH (score for one species only)

Spreading vigour varies with species

- Redwoods, Leyland cypresses, cedars and spruces (very low risk – no need to proceed further) 0
- Radiata (*P. radiata*) and ponderosa (*P. ponderosa*) pine, Lawsons cypress (*C. lawsoniana*) 1
- Muricata (*P. muricata*) and maritime (*P. pinaster*) pine and larches (*Larix* spp) 2
- Corsican (*P. nigra*) and mountain/dwarf mountain (*P. uncinata/mugo*) pine 3
- Douglas-fir ^g (*Ps. menziesii*), Scots pine (*P. sylvestris*) 4
- Lodgepole/contorta pine (*P. contorta*) 5

1

Enter score (0, 1, 2, 3, 4 or 5) here

2. SPECIES – PALATABILITY

Palatability varies with species

- Radiata, maritime and ponderosa pine 1
- Lodgepole and muricata pine and European larch 2
- Scots and mountain/dwarf mountain pine and Douglas-fir 3
- Corsican pine 4

1

Enter score (0, 1, 2, 3 or 4) here

3. SITING OF NEW PLANTING ^{c, d}

Trees are located on

- Sites well sheltered from prevalent and strong winds 0
- Flat sites (<10°), partially exposed to strong/prevalent winds 1
- Lea slopes where strong eddy gusts are likely 2
- Flat sites (<10°), fully exposed to strong/prevalent winds 3
- Either elevated 'take-off' sites, (ridge-tops, or base of exposed slopes >10°) or sloping land, fully exposed to strong/prevalent winds 4

2

Enter score (0, 1, 2, 3 or 4) here

4. DOWNWIND LANDUSE – GRAZING

Wilding establishment influenced by grazing (particularly with sheep)

- Intensive grazing on developed pasture 0^d
- Regular mob stocking with sheep ^e 1^d
- Semi-improved grazing (sheep/cattle)/ occasional mob stocking with sheep 2^d
- Extensive grazing only ^e 3^d
- No grazing 4^d

0

Enter score (0, 1, 2, 3 or 4) here

5. DOWNWIND VEGETATION COVER (if Douglas-fir involved see ^g in Notes)

Wilding establishment influenced by competition from existing vegetation

- Plantation forest, developed pasture (intensive grazing) 0^d
- Native forest^h, shrubland/tussock/grassland with a continuous and dense vegetation cover 1^d
- Forest/shrubland/tussock/grassland with few gaps 2^d
- Open forest and/or scattered patches of dense shrubland/tussock/grassland with many gaps 3^d
- Open slips/rockland and/or light, low-stature shrubland/tussock/grassland 4^d

0

TOTAL SCORE 0

Mort Shepherd



1/04/2025