

Wilding Risk Calculation – Site 1

SPECIES – GROWTH (score for one species only)

Spreading vigour varies with species

- Redwoods, Leyland cypresses, cedars and spruces (very low risk – no need to proceed further) 0
- Radiata (*P. radiata*) and ponderosa (*P. ponderosa*) pine, Lawsons cypress (*C. lawsoniana*) 1
- Muricata (*P. muricata*) and maritime (*P. pinaster*) pine and larches (*Larix* spp) 2
- Corsican (*P. nigra*) and mountain/dwarf mountain (*P. uncinata/mugo*) pine 3
- Douglas-fir ^g (*Ps. menziesii*), Scots pine (*P. sylvestris*) 4g
- Lodgepole/contorta pine (*P. contorta*) 5

1

Enter score (0, 1, 2, 3, 4 or 5) here

SPECIES – PALATABILITY

Palatability varies with species

- Radiata, maritime and ponderosa pine 1
- Lodgepole and muricata pine and European larch 2
- Scots and mountain/dwarf mountain pine and Douglas-fir 3
- Corsican pine 4

1

Enter score (0, 1, 2, 3 or 4) here

SITING OF NEW PLANTING ^{c, d}

Trees are located on

- Sites well sheltered from prevalent and strong winds 0
- Flat sites (<10°), partially exposed to strong/prevalent winds 1
- Lea slopes where strong eddy gusts are likely 2
- Flat sites (<10°), fully exposed to strong/prevalent winds 3
- Either elevated 'take-off' sites, (ridge-tops, or base of exposed slopes >10°) or sloping land, fully exposed to strong/prevalent winds 4

2

Enter score (0, 1, 2, 3 or 4) here

DOWNWIND LANDUSE – GRAZING

Wilding establishment influenced by grazing (particularly with sheep)

- Intensive grazing on developed pasture 0^d
- Regular mob stocking with sheep ^e 1^d
- Semi-improved grazing (sheep/cattle)/ occasional mob stocking with sheep 2^d
- Extensive grazing only ^e 3^d
- No grazing 4^d

1

Enter score (0, 1, 2, 3 or 4) here

DOWNWIND VEGETATION COVER (if Douglas-fir involved see ^g in Notes)

Wilding establishment influenced by competition from existing vegetation

- Plantation forest, developed pasture (intensive grazing) 0^d
- Native forest^h, shrubland/tussock/grassland with a continuous and dense vegetation cover 1^d
- Forest/shrubland/tussock/grassland with few gaps 2^d
- Open forest and/or scattered patches of dense shrubland/tussock/grassland with many gaps 3^d
- Open slips/rockland and/or light, low-stature shrubland/tussock/grassland 4^d

0

TOTAL SCORE 0

Wilding Risk Calculation – Site 2

SPECIES – GROWTH (score for one species only)

Spreading vigour varies with species

- Redwoods, Leyland cypresses, cedars and spruces (very low risk – no need to proceed further) 0
- Radiata (*P. radiata*) and ponderosa (*P. ponderosa*) pine, Lawsons cypress (*C. lawsoniana*) 1
- Muricata (*P. muricata*) and maritime (*P. pinaster*) pine and larches (*Larix* spp) 2
- Corsican (*P. nigra*) and mountain/dwarf mountain (*P. uncinata/mugo*) pine 3
- Douglas-fir ^g (*Ps. menziesii*), Scots pine (*P. sylvestris*) 4g
- Lodgepole/contorta pine (*P. contorta*) 5

1

Enter score (0, 1, 2, 3, 4 or 5) here

SPECIES – PALATABILITY

Palatability varies with species

- Radiata, maritime and ponderosa pine 1
- Lodgepole and muricata pine and European larch 2
- Scots and mountain/dwarf mountain pine and Douglas-fir 3
- Corsican pine 4

1

Enter score (0, 1, 2, 3 or 4) here

SITING OF NEW PLANTING ^{c, d}

Trees are located on

- Sites well sheltered from prevalent and strong winds 0
- Flat sites (<10°), partially exposed to strong/prevalent winds 1
- Lea slopes where strong eddy gusts are likely 2
- Flat sites (<10°), fully exposed to strong/prevalent winds 3
- Either elevated ‘take-off’ sites, (ridge-tops, or base of exposed slopes >10°) or sloping land, fully exposed to strong/prevalent winds 4

4

Enter score (0, 1, 2, 3 or 4) here

DOWNWIND LANDUSE – GRAZING

Wilding establishment influenced by grazing (particularly with sheep)

- Intensive grazing on developed pasture 0^d
- Regular mob stocking with sheep ^e 1^d
- Semi-improved grazing (sheep/cattle)/ occasional mob stocking with sheep 2^d
- Extensive grazing only ^e 3^d
- No grazing 4^d

0

Enter score (0, 1, 2, 3 or 4) here

DOWNWIND VEGETATION COVER (if Douglas-fir involved see ^g in Notes)

Wilding establishment influenced by competition from existing vegetation

- Plantation forest, developed pasture (intensive grazing) 0^d
- Native forest^h, shrubland/tussock/grassland with a continuous and dense vegetation cover 1^d
- Forest/shrubland/tussock/grassland with few gaps 2^d
- Open forest and/or scattered patches of dense shrubland/tussock/grassland with many gaps 3^d
- Open slips/rockland and/or light, low-stature shrubland/tussock/grassland 4^d

0

TOTAL SCORE 0

Mort Shepherd



27//2024