

Examining the Jelly Ocean Hypothesis in the Northwest North Atlantic

Nicholas R. Record, Benjamin Tupper, Omi Johnson

Abstract

The Jelly Ocean Hypothesis posits that gelatinous zooplankton populations are increasing worldwide as a result of changing ocean conditions. This hypothesis has been the subject of debate in ocean science. We examined the Northeast US Continental Shelf for increases in the occurrence of gelatinous zooplankton taxa using a 50-year survey and a species distribution model. Time series analysis showed statistically significant increases in salps, siphonophores, and coelenterates (scyphozoans) in late summer and early autumn across the entire study region. Species distribution models showed that the drivers of the occurrence of gelatinous taxa were XYZ (oxygen?) ETCECT The results form one piece of evidence in favor of a long-term rise in gelatinous species. The roughly synchronized increases in these widely diverse taxa suggest that changes in environmental or ecological conditions favor a gelatinous body plan generally.

Introduction

In the study of zooplankton ecology, there has often been a “crustaceocentric” bias (Haddock 2004). The study of gelatinous species, collectively termed gelata, has been underemphasized as compared to crustaceans. In the early 2000s, gelata moved into the spotlight, following conspicuous blooms that damaged fisheries (Kideys et al. 2005), aquaculture (Clinton et al. 2021), energy production (Wang et al. 2023), and recreation (Nunes et al. 2015). In addition to motivating new research on gelata, these events raised the question of whether there was a larger pattern of change.

Emerging from a series of studies was the Jelly Ocean Hypothesis, which posits that gelatinous zooplankton populations are increasing worldwide as a result of changing ocean conditions, primarily due to anthropogenic activities. These studies argued that due a multiplicity of causes, abundance of gelatinous species is growing. Proposed causes are primarily human-driven, including warming (ref), eutrophication, translocation (Kideys et al. 2005), overfishing, bottom trawling (Qui 2014), pollution, aquaculture, and acidification (Richardson and Gibbons 2008). There is also evidence that shifts toward dominance of gelata are difficult to reverse because of dynamics like hysteresis (Richardson et al. 2009). The hypothesis echos a concern that has been around for centuries; the 19th century novelist Jules Verne once predicted that, “crowded with jellyfish, squid, and other devilfish, the oceans will have become huge centers of infection” (REF). While the term “infection” might not be fair to jellyfish, who simply inhabit the ocean, the potential influence on human industry and recreation make the question worthy of examination.

Testing the Jelly Ocean Hypothesis requires long time series (>30 years) of gelata abundances in order to compare to historical levels. The shortage of such datasets has left the Jelly Ocean Hypothesis largely unsupported (Condon et al. 2012). In lieu of time series, studies have used qualitative data such as news reports to argue that gelata are increasing globally (Brotz et al. 2012). However, it is difficult to distinguish these perceived increases from multi-annual or multi-decadal fluctuations. A recent meta-analysis of time series showed that most perceived increases appear to reflect a ~20 year cyclicity, with a worldwide upswing in the 1990s, coinciding with the increase in concern over gelata (Condon et al. 2013), although some notable long-term increases in gelata were not included in the analysis (Atkinson et al. 2004). A metaanalysis of a corpus of research on this topic found flawed citations and misinterpreted conclusions (Sanz-Martin et al. 2016), and at this point, the Jelly Ocean Hypothesis remains very much an open question.

In some ways, the Jelly Ocean Hypothesis is loosely posed. Gelata are taxonomically very diverse, encompassing multiple phyla. This diversity confounds attempts to quantify increasing trends. The list of species classified under the gelatinous umbrella varies, and includes taxa with drastic and important differences in their life histories, trophic positions, and ecological roles. For example, pelagic ctenophores are primarily copepod predators, where as pelagic tunicates graze some of the smallest microbes. Cnidarians and ctenophores are the two most common taxa to group together in analysis, and yet they are phylogenetically very distant (Moroz et al. 2014). Perhaps the only characteristic common to the many gelatinous phyla is a body with a high water content, alternatively expressed as a low carbon:volume ratio. There is no *a priori* reason to expect that phyla with such striking differences and perhaps only a superficial similarity should be grouped together with respect to their temporal trends. Some have suggested that the Jelly Ocean Hypothesis is ill-posed—i.e. which species are increasing, where, and for what reason (Gibbons & Richardson 2013)?

On the other hand, despite their major differences, there is some evidence that a diverse collection of gelata can vary together (Condon et al. 2013). This suggests that certain conditions can occur that favor the gelatinous body form in general, over other strategies that utilize a high carbon:volume ratio (e.g. crustaceans). This reframes the Jelly Ocean Hypothesis in trait-based terms: conditions in the ocean ecosystem can shift to benefit the gelatinous body form. This would be inclusive of gelata in general, despite differences in phylogeny, trophic position, or other life history traits.

We revisited the Jelly Ocean Hypothesis using a 50-year spatial survey on the northwest North Atlantic Shelf. We addressed two main questions: 1) what are the temporal changes in gelata presence; and 2) what are the underlying conditions associated with changes. We examined three taxonomic groups well-represented in the dataset—siphonophores, salps, and coelenterates—to explore the idea that certain conditions can favor the gelatinous body plan in general.

Methods

Survey Data

The Northeast Fisheries Science Center Ecosystem Monitoring (ECOMON) program collects seasonal zooplankton measurements as part of its mission to monitor and assess the Northeast Continental Shelf ecosystem (Figure 1). Sampling includes oblique tows using a 61 cm bongo net fitted with a 333 μm mesh at stratified locations across the shelf (Figure 1). Tows are a minimum of five minutes in duration and sample from the surface to within 5 m of the seabed or to a maximum depth of 200 m. A mechanical flowmeter fitted in the mouth of each net is used to quantify volume sampled. Gelatinous zooplankton are identified into broad taxonomic groups, including siphonophores, ctenophores, salps, and coelenterates (i.e. scyphozoans). The broad grouping, while missing taxonomic detail has allowed for internal consistency over the duration of the survey. The analysis conducted here uses presence or absence of each taxonomic group.

Time series analysis

The ECOMON survey has varied in frequency over its duration. To correct for biases, we followed the common approach of grouping by large spatial polygons that represent different habitats, and analyzed the time series by month. Polygons include: Gulf of Maine, Georges Bank, New York Bight, and Mid Atlantic Bight (Walsh et al. 2015). Because of the low taxonomic resolution of gelatinous taxa and the bulk biomass quantification, we viewed measurements in a presence-absence context, tracking the proportion of samples in a region and in a time period that contain presences. For each month and within each region, we tested for statistically significant increases over time using a bivariate correlation test against year.

Spatial analysis

To test for environmental associations with high or low occurrence of gelatinous taxa across the study domain, we developed a species distribution model for each taxonomic group. We used physical covariates from




Figure 1: Map of the study region. Points indicate sampling locations colored by the four polygons: Gulf of Maine, Georges Bank, Southern New England, and Mid-Atlantic Bight.

the Global Ocean Physics Reanalysis (GLORYS) and biogeochemical covariates the Operational Mercator Ocean biogeochemical global ocean analysis (NEMO) obtained from the Copernicus repository (copernicus.eu), which provide daily temporal resolution and 1/12 and quarter-degree resolution respectively. Physical covariates included: sea surface temperature, sea bottom temperature, sea surface salinity, mixed layer thickness, and surface velocity magnitude. Biogeochemical covariates included surface chlorophyll, surface net primary productivity, surface oxygen, surface nitrate, surface phosphate, and surface silicate. We also included day length and the time derivative of day length to represent seasonal dependencies, as well as bathymetric depth and bathymetric slope.

We used a boosted regression tree algorithm to fit the observations to environmental covariates. Boosted regression trees are well suited to variables that have sharp jumps or thresholds (Johnson et al. 2024), as is the case for the presence of gelatinous zooplankton, and is less prone to overfitting in species distribution models (Elith et al., 2008). We modeled each taxon using tidymodels workflows, optimizing the inbuilt area-under-curve (AUC) metric (Fielding and Bell 1997, Kuhn and Wickham, 2020). We used a set of hyperparameters optimized in a previous study that used the same dataset and methodology (Johnson et al. 2024). We used a testing split of 25%, and ran an ensemble of 25 model runs to quantify algorithmic uncertainty around the median output. Cross validation between environmental covariates was used in the interpretation of output, but not in covariate selection because the goal of the model was to explore covariate associations. The resulting response curves were used to understand associations with environmental covariates.

Results

Time series analysis

Over almost five decades of measurements, there were consistent patterns of statistically significant increases in the recorded presence of gelatinous taxa (Figure 2). While the survey does not cover every month in every year, there is enough coverage across seasons to see consistent patterns. Out of 144 month-region combinations, 56 (39%) indicate a statistically significant increase ($p < 0.05$) and zero (0%) indicate a statistically significant decrease. In the months August–November, 40 of the 48 time series (83%) show a statistically significant increase. In other words, in late summer and early fall, all taxa show an increase across the entire study region.

The results during other times of year were more mixed. On Georges Bank, there were statistically significant increases across the spring months for siphonophores and coelenterates. For the other taxa, regions, and months, there were some scattered increases but no strong pattern.

Spatial analysis

Discussion

References

- Atkinson, Angus, Volker Siegel, Evgeny Pakhomov, and Peter Rothery. 2004. “Long-Term Decline in Krill Stock and Increase in Salps Within the Southern Ocean.” *Nature* 432 (7013): 100–103.
- Brotz, Lucas, William WL Cheung, Kristin Kleisner, Evgeny Pakhomov, and Daniel Pauly. 2012. “Increasing Jellyfish Populations: Trends in Large Marine Ecosystems.” In *Jellyfish Blooms IV: Interactions with Humans and Fisheries*, 3–20. Springer.
- Clinton, Morag, David EK Ferrier, Samuel AM Martin, and Andrew S Brierley. 2021. “Impacts of Jellyfish on Marine Cage Aquaculture: An Overview of Existing Knowledge and the Challenges to Finfish Health.” *ICES Journal of Marine Science* 78 (5): 1557–73.
- Condon, Robert H, Carlos M Duarte, Kylie A Pitt, Kelly L Robinson, Cathy H Lucas, Kelly R Sutherland, Hermes W Mianzan, et al. 2013. “Recurrent Jellyfish Blooms Are a Consequence of Global Oscillations.” *Proceedings of the National Academy of Sciences* 110 (3): 1000–1005.




Figure 2: Time series of the proportion of presences of the indicated taxon broken down by month and regional polygon. Stars indicate a statistically significant ($p < 0.05$) increase over the duration of the time series.

- Condon, Robert H, William M Graham, Carlos M Duarte, Kylie A Pitt, Cathy H Lucas, Steven HD Haddock, Kelly R Sutherland, et al. 2012. "Questioning the Rise of Gelatinous Zooplankton in the World's Oceans." *Bioscience* 62 (2): 160–69.
- Haddock, Steven HD. 2004. "A Golden Age of Gelata: Past and Future Research on Planktonic Ctenophores and Cnidarians." *Hydrobiologia* 530 (1): 549–56.
- Kideys, Ahmet Erkan, Abolchaseem Roohi, Siamak Bagheri, Galina Finenko, and Lyudmila Kamburska. 2005. "Impacts of Invasive Ctenophores on the Fisheries of the Black Sea and Caspian Sea." *Oceanography* 18 (2): 76–85.
- Nunes, Paulo ALD, Maria L Loureiro, Laia Piñol, Sergio Sastre, Louinord Voltaire, and Antonio Canepa. 2015. "Analyzing Beach Recreationists' Preferences for the Reduction of Jellyfish Blooms: Economic Results from a Stated-Choice Experiment in Catalonia, Spain." *PloS One* 10 (6): e0126681.
- Richardson, Anthony J, Andrew Bakun, Graeme C Hays, and Mark J Gibbons. 2009. "The Jellyfish Joyride: Causes, Consequences and Management Responses to a More Gelatinous Future." *Trends in Ecology & Evolution* 24 (6): 312–22.
- Richardson, Anthony J, and Mark J Gibbons. 2008. "Are Jellyfish Increasing in Response to Ocean Acidification?" *Limnology and Oceanography* 53 (5): 2040–45.
- Sanz-Martin, Marina, Kylie A Pitt, Robert H Condon, Cathy H Lucas, Charles Novaes de Santana, and Carlos M Duarte. 2016. "Flawed Citation Practices Facilitate the Unsubstantiated Perception of a Global Trend Toward Increased Jellyfish Blooms." *Global Ecology and Biogeography* 25 (9): 1039–49.
- Walsh, Harvey J, David E Richardson, Katrin E Marancik, and Jonathan A Hare. 2015. "Long-Term Changes in the Distributions of Larval and Adult Fish in the Northeast US Shelf Ecosystem." *PLoS One* 10 (9): e0137382.
- Wang, Xiaocheng, Qingqing Jin, Lu Yang, Chuan Jia, Chunjiang Guan, Haining Wang, and Hao Guo. 2023. "Aggregation Process of Two Disaster-Causing Jellyfish Species, *Nemopilema Nomurai* and *Aurelia Coerulea*, at the Intake Area of a Nuclear Power Cooling-Water System in Eastern Liaodong Bay, China." *Frontiers in Marine Science* 9: 1098232.