DMMR Tutorial sheet 3

Functions, Sequences

October 7, 2015

Some of the exercises for this tutorial are taken from Chapter 2 and 8 of the book: Kenneth Rosen, Discrete Mathematics and its Applications, 7th Edition, McGraw-Hill, 2012.

1. Determine whether the function $f: \mathbb{Z} \times \mathbb{Z} \to \mathbb{Z}$ is surjective if

(a)
$$f(m,n) = m^2 + n^2$$

(c)
$$f(m,n) = |n|$$

(b)
$$f(m, n) = m$$

(d)
$$f(m,n) = m - n$$

- 2. (a) Prove that a strictly decreasing function from \mathbb{R} to itself is one-to-one.
 - (b) Give an example of a decreasing function from \mathbb{R} to itself that is not one-to-one.
- 3. Determine (and prove) whether each of these sets is countably infinite or uncountable. For those that are countably infinite, exhibit a one-to-one correspondence (i.e., bijection) between the set of positive integers and that set.
 - (a) the odd negative integers
 - (b) the real numbers in the open interval (0,2)
 - (c) the irrational numbers in the open interval (0, 2)
 - (d) the set $A \times \mathbb{Z}^+$ where $A = \{2, 3\}$
- 4. A vending machine dispensing books of stamps accepts only \$1 coins, \$1 bills and \$5 bills.
 - (a) Find a recurrence relation for the number of ways to deposit n dollars in the vending machine, where the order in which the coins and bills are deposited matters.
 - (b) What are the initial conditions?
 - (c) How many ways are there to deposit \$10 for a book of stamps?
- 5. Show that the set of functions from positive integers to the set $\{0, 1, 2, 3, 4, 5, 6, 7, 8, 9\}$ is uncountable.

Solutions (to the last question ONLY on the sheet) must be handed in on paper to the ITO by Wednesday, 14 October, 4:00pm. Please post it into the grey metal box on the wall outside the ITO.