

# VIP Cheatsheet: Second-order ODE

Shervine AMIDI

May 27, 2018

## General case

□ **General form** – The general form of a second-order ODE can be written as a function  $F$  of  $x, y, y'$  and  $y''$  as follows:

$$F(x, y, y', y'') = 0$$

□ **Methods of resolution** – The table below summarizes the general tricks to apply when the ODE has the following classic forms:

Old form	Trick	New form
$F(x, y', y'') = 0$	$y' \triangleq u, \quad y'' = \frac{du}{dx}$	$G(x, u, \frac{du}{dx}) = 0$
$F(y, y', y'') = 0$	$y' \triangleq u, \quad y'' = u \frac{du}{dy}$	$G(y, u, \frac{du}{dy}) = 0$
$F(y', y'') = 0$	$y' \triangleq u, \quad y'' = \frac{du}{dx}$ $y' \triangleq u, \quad y'' = u \frac{du}{dy}$	Missing- $y$ approach $G(u, \frac{du}{dx}) = 0$ Missing- $x$ approach $G(u, \frac{du}{dy}) = 0$

□ **Standard form of a linear ODE** – The standard form of a second-order linear ODE is expressed with  $p, q$  and  $r$  known functions of  $x$  such that:

$$y'' + p(x)y' + q(x)y = r(x)$$

for which the total solution  $y$  is the sum of a homogeneous solution  $y_h$  and a particular solution  $y_p$ :

$$y = y_h + y_p$$

*Remark: if  $r = 0$ , then the ODE is homogeneous (and we have  $y_p = 0$ ). If  $r \neq 0$ , then the ODE is said to be inhomogeneous.*

□ **Linear dependency** – Two functions  $y_1, y_2$  are said to be *linearly dependent* if  $\frac{y_2}{y_1} = C$  constant. Conversely, they are *linearly independent* if  $\frac{y_2}{y_1} \neq C$ .

## Linear homogeneous – Variable coefficients

□ **Method of reduction of order** – Let  $y_1$  be a solution to the equation  $y'' + p(x)y' + q(x)y = 0$ . By noting  $C_1, C_2$  constants, the global solution  $y_h$  is written as:

$$y_h = C_1 y_1 + C_2 y_1 \int \frac{e^{-\int p dx}}{y_1^2} dx$$

*Remark: Here, for any function  $p$ , the notation  $\int p dx$  denotes the primitive of  $p$  without additive constant.*

## Linear homogeneous – Constant coefficients

□ **General form** – The general form of a linear homogeneous second-order ODE with  $a, b, c$  constant coefficients is:

$$ay'' + by' + cy = 0$$

□ **Resolution** – Based on the types of solution of the characteristic equation  $a\lambda^2 + b\lambda + c = 0$ , and by noting  $\Delta = b^2 - 4ac$  its discriminant, we distinguish the following cases:

Name	Case	Roots	Solution
Two distinct real roots	$\Delta > 0$	$\lambda_1 = \frac{-b + \sqrt{\Delta}}{2a}$ $\lambda_2 = \frac{-b - \sqrt{\Delta}}{2a}$	$y_h = C_1 e^{\lambda_1 x} + C_2 e^{\lambda_2 x}$
Double real root	$\Delta = 0$	$\lambda = -\frac{b}{2a}$	$y_h = [C_1 + C_2 x] e^{\lambda x}$
Complex conjugate roots	$\Delta < 0$	$\lambda_1 = \alpha + i\beta$ $\lambda_2 = \alpha - i\beta$ where $\alpha = -\frac{b}{2a}$ and $\beta = \frac{\sqrt{ \Delta }}{2a}$	$y_h = [C_1 \cos(\beta x) + C_2 \sin(\beta x)] e^{\alpha x}$

## A special case: the Euler-Cauchy equation

□ **General form** – The Euler-Cauchy equation is a special case of linear homogeneous ODEs and has the following general form, where each  $a_i \in \mathbb{R}$  is a constant coefficient:

$$a_n x^n y^{(n)} + a_{n-1} x^{n-1} y^{(n-1)} + \dots + a_1 x y' + a_0 y = 0$$

□ **Second-order case** – For  $n = 2$ , by noting  $y = x^m$ , the ODE provides the indicial equation:

$$am^2 + (b-a)m + c = 0$$

with discriminant  $\Delta = (b-a)^2 - 4ac$  and where the resolution of the ODE depends on the cases summarized in the table below.

Name	Case	Roots	Solution
Two distinct real roots	$\Delta > 0$	$m_1 = \frac{-b + a + \sqrt{\Delta}}{2a}$ $m_2 = \frac{-b + a - \sqrt{\Delta}}{2a}$	$y_h = C_1 x^{m_1} + C_2 x^{m_2}$
Double real root	$\Delta = 0$	$m = -\frac{b-a}{2a}$	$y_h = [C_1 + C_2 \ln  x ] x^m$
Complex conjugate roots	$\Delta < 0$	$m_1 = \alpha + i\beta$ $m_2 = \alpha - i\beta$ <p>where <math>\alpha = -\frac{b-a}{2a}</math> and <math>\beta = \frac{\sqrt{ \Delta }}{2a}</math></p>	$y_h = [C_1 \cos(\beta \ln  x )$ $+ C_2 \sin(\beta \ln  x )] x^\alpha$

### Linear inhomogeneous – Variable coefficients

□ **Wronskian** – Given  $y_1$  and  $y_2$  the two solutions of the homogeneous equation, we define the Wronskian  $W$  as follows:

$$W = y_1 y_2' - y_2 y_1'$$

□ **Method of Variation of Parameters** – The particular solution  $y_p$  of the inhomogeneous ODE is given by:

$$y_p = -y_1 \int \frac{y_2 r}{W} dx + y_2 \int \frac{y_1 r}{W} dx$$

### Linear inhomogeneous – Constant coefficients

□ **Undetermined coefficients method** – The particular solution  $y_p$  of the inhomogeneous ODE  $ay'' + by' + cy = r(x)$  is determined from the correspondence table below:

Form of $r$	Form of $y_p$
$C$	$A$
$x^n, n \in \mathbb{N}^*$	$A_0 + A_1 x + \dots + A_n x^n$
$e^{\gamma(x)}$	$A e^{\gamma x}$
$\cos(\omega x)$ or $\sin(\omega x)$	$A \cos(\omega x) + B \sin(\omega x)$
$x^n e^{\gamma x} \cos(\omega x)$ or $x^n e^{\gamma x} \sin(\omega x)$	$(A_0 + A_1 x + \dots + A_n x^n) \cos(\omega x) e^{\gamma x} +$ $(B_0 + B_1 x + \dots + B_n x^n) \sin(\omega x) e^{\gamma x}$

*Remark: all new constants are determined after plugging back  $y_p$  into the ODE.*

□ **Modification rule** – If the particular solution  $y_p$  picked from the above table matches either  $y_1$  or  $y_2$ , then has to be multiplied by the lowest power of  $x$  such that it is no more the case.

□ **Sum rule** – If  $r(x)$  is a sum of functions of the first column of the above table, then  $y_p$  is the sum of its associated particular solutions.