

On how to compute $\int_a^b f$?

1st attempt: (Sequential approach)

Thm: Let $f \in \mathcal{B}[a, b]$. Then $f \in \mathcal{R}[a, b] \iff \exists$ a seqn. $\{P_n\} \subseteq \mathcal{P}[a, b]$

s.t.
$$\lim_{n \rightarrow \infty} [U(f, P_n) - L(f, P_n)] = 0.$$

Moreover, in this case:

$$\int_a^b f = \lim_{n \rightarrow \infty} U(f, P_n) = \lim_{n \rightarrow \infty} L(f, P_n).$$

Proof: [Note: $\because U(f, P) - L(f, P) \geq 0 \forall P \in \mathcal{P}[a, b]$,
 $\{U(f, P_n) - L(f, P_n)\} \subseteq \mathbb{R}_{\geq 0}$.]

" \Rightarrow " Let $f \in \mathcal{R}[a, b]$. We know: "for $\varepsilon > 0$, $\exists P_\varepsilon \in \mathcal{P}[a, b]$
 $\Rightarrow U(f, P_\varepsilon) - L(f, P_\varepsilon) < \varepsilon$ ".

\leftarrow Call it "Cauchy Criterion"

\therefore For each $n \in \mathbb{N}$, $\exists P_n \in \mathcal{P}[a, b]$ s.t.

$$U(f, P_n) - L(f, P_n) < \frac{1}{n}.$$

By \otimes above: $U(f, P_n) - L(f, P_n) \longrightarrow 0$
as $n \longrightarrow \infty$.

" \Leftarrow " If $\lim_{n \rightarrow \infty} [U(f, P_n) - L(f, P_n)] = 0$, then for $\varepsilon > 0$

$$\exists N \in \mathbb{N} \text{ s.t. } U(f, P_N) - L(f, P_N) < \varepsilon.$$

$\Rightarrow f \in \mathcal{R}[a, b]$. By Cauchy Criterion.

Finally, if $f \in \mathcal{R}[a, b]$, then for $\{P_n\} \subseteq \mathcal{P}[a, b]$
as above:

We have:

$$0 \leq U(f, P_n) - \overline{\int_a^b f} \quad \leftarrow (\because \overline{\int f} = \inf U(f, P))$$

$$= U(f, P_n) - \underline{\int_a^b f} \quad \leftarrow (\because f \in \mathcal{R}[a, b])$$

$$(\text{As } \underline{\int f} = \sup_P L(f, P)) \rightarrow \leq U(f, P_n) - L(f, P_n) \rightarrow 0 \text{ as } n \rightarrow \infty$$

$$\Rightarrow \lim_{n \rightarrow \infty} U(f, P_n) = \overline{\int_a^b f} = \underline{\int_a^b f}.$$

Finally, $L(f, P_n) = U(f, P_n) - (U(f, P_n) - L(f, P_n))$

$$\downarrow$$

$$\int_a^b f - 0 \quad \text{as } n \rightarrow \infty$$

$$\Rightarrow \lim_{n \rightarrow \infty} L(f, P_n) = \int_a^b f. \quad \square$$

DANGER: $a_n - b_n \rightarrow 0 \not\Rightarrow$ Even $\lim a_n$ &/or $\lim b_n$ exist !!

Remark: Evidently, if $\exists \{P_n\} \subseteq \mathcal{P}[a, b]$ s.t.

$L(f, P_n) \rightarrow c$ & $U(f, P_n) \rightarrow c$, then

$$f \in \mathcal{R}[a, b] \quad \& \quad \int_a^b f = c.$$

A nice way to prove existence of \mathcal{R} -integrability
(& evaluating too) of bdd fns. [Reminding us
Newton].

Eg: $f(x) = x^2$ $x \in [0, 1]$.

Fix $n \in \mathbb{N}$.

Consider "the classical" partition:

↑ Why call it classical?

[BTW: $f \in C[0, 1]$
 $\Rightarrow f \in R[0, 1]$.
But we prove it directly.]

$$P_n: 0 = x_0 < x_1 = \frac{1}{n} < x_2 = \frac{2}{n} < \dots < x_{n-1} = \frac{n-1}{n} < x_n = 1$$

$$\therefore \underline{I_j} = \left[\frac{j-1}{n}, \frac{j}{n} \right] \quad \forall j = 1, \dots, n$$

$\therefore f$ is \uparrow , it follows that:

$$m_j = \left(\frac{j-1}{n} \right)^2 \quad \& \quad M_j = \left(\frac{j}{n} \right)^2$$

$\forall j = 1, \dots, n$

$$\therefore U(f, P_n) = \sum_{j=1}^n M_j \times \frac{1}{n} = \sum_{j=1}^n \frac{j^2}{n^2} \times \frac{1}{n}$$

$$= \frac{1}{n^3} \sum_{j=1}^n j^2$$

$$= \frac{1}{n^3} (1^2 + 2^2 + \dots + n^2) = \frac{1}{n^3} \times \frac{1}{6} \times n \times (n+1)(2n+1)$$

$$= \frac{1}{6} \left(1 + \frac{1}{n} \right) \left(2 + \frac{1}{n} \right)$$

Also, $L(f, P_n) = \sum_{j=1}^n m_j \frac{1}{n} = \frac{1}{n^3} \sum_{j=1}^n (j-1)^2$

$$= \frac{1}{n^3} (1^2 + 2^2 + \dots + (n-1)^2)$$

$$= \frac{1}{n^3} \times \frac{1}{6} \times (n-1) \times n \times (2(n-1)+1)$$

$$= \frac{1}{6} \times \left(1 - \frac{1}{n} \right) \times \left(2 - \frac{1}{n} \right)$$

$$\therefore U(f, P_n) \rightarrow \frac{1}{3} \quad \& \quad L(f, P_n) \rightarrow \frac{1}{3}$$

$$\Rightarrow U(f, P_n) - L(f, P_n) \longrightarrow 0.$$


$$\Rightarrow \underline{f \in \mathcal{R}[0,1]} \quad \& \quad \underline{\int_0^1 f = \lim_{n \rightarrow \infty} U(f, P_n) = \frac{1}{3}}.$$

Q: $\mathcal{R}[a,b] \setminus \mathcal{C}[a,b] \neq \varnothing$?

i.e. $\exists f \in \mathcal{R}[a,b]$ s.t. $f \in \mathcal{R}[a,b]$ but $f \notin \mathcal{C}[a,b]$??

Ans: Yes.

Eg:



$$f(x) = \begin{cases} 1 & 0 \leq x < \frac{1}{2} \\ \frac{1}{2} & x = \frac{1}{2} \\ 0 & \frac{1}{2} < x \leq 1. \end{cases}$$

Claim: $f \in \mathcal{R}[0,1]$, clearly, $f \notin \mathcal{C}[0,1]$.


Proof

Let $\varepsilon > 0$ (small).

Consider the partition: $P_\varepsilon: 0 = x_0 < x_1 = \frac{1}{2} - \varepsilon < x_2 = \frac{1}{2} + \varepsilon < x_3 = 1$

$$\text{i.e. } 0 < \frac{1}{2} - \varepsilon < \frac{1}{2} + \varepsilon < 1.$$

A 2-node \rightsquigarrow partition.



$$\begin{aligned} I_1 &= [0, \frac{1}{2} - \varepsilon] \\ I_2 &= [\frac{1}{2} - \varepsilon, \frac{1}{2} + \varepsilon] \\ I_3 &= [\frac{1}{2} + \varepsilon, 1] \end{aligned}$$

∴

$$\begin{aligned} \therefore m_1 &= 1, m_2 = 0, m_3 = 0 \\ M_1 &= 1, M_2 = 1, M_3 = 0 \end{aligned}$$

$$\begin{aligned}\therefore \underline{L(f, P_\varepsilon)} &= \sum_{j=1}^3 m_j |I_j| \\ &= 1 \times \left(\frac{1}{2} - \varepsilon\right) + 0 + 0 \\ &= \frac{1}{2} - \varepsilon.\end{aligned}$$

$$\begin{aligned}\nabla \underline{U(f, P_\varepsilon)} &= \sum_{j=1}^3 M_j |I_j| \\ &= 1 \times \left(\frac{1}{2} - \varepsilon\right) + 1 \times 2\varepsilon + 0 \\ &= \frac{1}{2} + \varepsilon.\end{aligned}$$

$$\therefore \underline{U(f, P_\varepsilon) - L(f, P_\varepsilon) = 2\varepsilon.}$$

$$\text{i.e. } U(f, P_\varepsilon) - L(f, P_\varepsilon) < 3\varepsilon.$$

why? $\Rightarrow f \in \mathcal{R}[0, 1].$

$$\underline{Q:} \quad \int_0^1 f = ?$$

$$\begin{aligned}\therefore \text{For } \varepsilon = \frac{1}{2n}, \text{ we have,} \\ U(f, P_n) - L(f, P_n) &= \frac{1}{n} \\ &\downarrow 0 \\ \Rightarrow \lim [U(f, P_n) - L(f, P_n)] &= 0. \\ \Rightarrow \int_0^1 f &= \frac{1}{2}.\end{aligned}$$

Remark: In fact, if $f \in \mathcal{B}[0, 1]$ with finitely many discontinuity, then $f \in \mathcal{R}[0, 1].$

— WAIT —

Let's make it more refined!!

The theory

Def: Let $P: a = x_0 < x_1 < \dots < x_{n-1} < x_n = b$. ~~For~~ A fr.

$T_P: \{I_j\}_{j=1}^n \longrightarrow [a, b]$ is called a tag ^{of P} if

$$T_P(I_j) \in I_j \quad \forall j = 1, \dots, n.$$

Simply


Given a partition $P: a = x_0 < x_1 < \dots < x_{n-1} < x_n = b$, a

tag or tag set ^{of P} is a collection of points $\{g_j\}_{j=1}^n$

s.t. $g_j \in I_j \quad \forall j = 1, \dots, n.$

Even Simpler.

A tag set $T_P = \{g_j\}_{j=1}^n$ where $g_j \in I_j \quad \forall j = 1, \dots, n.$



$$P: 0 < \frac{1}{2} < 1 < 2 < 2\frac{1}{2} < 4$$

$$T_P = \left\{ \frac{1}{6}, \frac{5}{8}, \frac{4}{3}, \frac{2}{5}, \frac{3}{3} \right\}$$

a tag set.


Note: If P has n -nodes [nodes excludes end points $\Rightarrow n+1$ subintervals], then $\#T_P = n+1 \quad \forall$ tag set T_P .

Def: Let $f \in \mathcal{B}[a, b]$, $P \in \mathcal{P}[a, b]$. & Suppose T_P a tag set of P .

The Riemann sum of f w.r.t. (P, T_P) is defined by:

$$\underline{S(f, P)} := \sum_{j=1}^n f(y_j) |I_j|, \quad \text{--- } (*)$$

where: $T_P = \{y_j\}_{j=1}^n$.



Note: ① $S(f, P)$ depends on $T_P = \{y_j\}_{j=1}^n$.

② LHS of $(*)$ doesn't involve T_P but it is there!!

③ \exists infinitely many tag sets for a given partition P .

④ In fact a question; What is the meaning of Riemann sum?

Fact: (Answering ④)

Suppose $P: a = x_0 < \dots < x_n = b$ be a partition &

$T_P = \{y_j\}_{j=1}^n$ be a tag of P .

$$\therefore m_j \leq f(y_j) \leq M_j \quad \forall j = 1, \dots, n$$

$$\Rightarrow m_j |I_j| \leq f(y_j) |I_j| \leq M_j |I_j|$$

$$\forall j = 1, \dots, n.$$

$$\Rightarrow L(f, P) \leq S(f, P) \leq U(f, P).$$



\therefore it is a better approximation.

HOWEVER, $S(f, P)$ depends on T_P
but $L(f, P)$ & $U(f, P)$ does not!!

