


u.c. & differentiability:

Recall:  $f_n \rightarrow f$  &  $f'_n(x_0)$  exists  $\forall n \not\Rightarrow f'(x_0)$  exists.

Another extreme example:

eg:



eg: Suppose  $f_n(x) = \frac{x}{1+nx^2}$ ,  $x \in \mathbb{R}$ ,  $n \geq 1$ .

$\therefore \{f_n\} \subseteq C(\mathbb{R})$ ,

$$|f_n(x)| = \left| \frac{x}{1+nx^2} \right|$$

$$\leq \frac{|x|}{2\sqrt{n}|x|}$$

$$= \frac{1}{2\sqrt{n}}$$

$$\forall x \neq 0.$$

dec.

$$\begin{aligned} & \because 1+nx^2 \geq 2\sqrt{n}|x| \\ & \forall x \in \mathbb{R} \\ & \Rightarrow \frac{1}{1+nx^2} \leq \frac{1}{2} \frac{1}{\sqrt{n}|x|} \\ & \forall x \in \mathbb{R}, n \geq 1, x \neq 0. \end{aligned}$$

$$\therefore |f_n(x)| \leq \frac{1}{2\sqrt{n}} \quad \forall n \geq 1, x \in \mathbb{R}.$$

$\uparrow (\because f_n(0) = 0 \forall n)$ .

$$\Rightarrow f_n \xrightarrow{u} f, \text{ where } f \equiv 0 \text{ on } \mathbb{R}.$$

M-test.

[Note that  $f_n(0) = 0 \forall n \geq 1$ ]

$$\text{Now: } f'_n(x) = \frac{-x \times 2nx + (1+nx^2)}{(1+nx^2)^2} = \frac{-n x^2 + 1}{(1+nx^2)^2}$$

$$= \frac{1-nx^2}{(1+nx^2)^2}$$

$\forall x \neq 0$

$$\therefore \frac{1-n}{(1+n)^2} = \frac{1/n-1}{(1/n^2+1/n)^2} \rightarrow 0 \text{ as } n \rightarrow \infty,$$

we have  $f'_n(x) \rightarrow F(x) \forall x \in \mathbb{R}$ ,

$$\text{where } F(x) = \begin{cases} 0 & x \neq 0 \\ 1 & x = 0. \end{cases}$$

[Note:  $f'_n(0) = 1 \forall n$ ]

In particular,  $f'_n(0) = 1 \quad \forall n$

$$\Rightarrow \lim_{n \rightarrow \infty} f'_n(0) = 1 \neq f'(0) = 0.$$

i.e. <sup>although</sup> ~~even~~  $\{f_n\}$   $f_n \xrightarrow{u} f$  &  $\{f_n\}$  &  $\{f\}$  diff. at

$x_0$ , still  $\frac{d}{dx} \left( \lim_{n \rightarrow \infty} f_n \right) \neq \lim_{n \rightarrow \infty} \left( \frac{d}{dx} f_n \right) . !!$

Failure of interchange of limits.

Still, we have some news:

Thm: Let  $\{f_n\}$  be a seqn of diff.  $f_n$  on  $[a, b]$  s.t.

(i)  $\exists x_0 \in [a, b]$  s.t.  $\{f_n(x_0)\}$  converges.

(ii)  $\{f'_n\}$  converges uniformly on  $[a, b]$ .

~~(iii)~~ Then  $\{f_n\}$  converges uniformly on  $[a, b]$  to a diff.  $f_n \rightarrow f$  &

$$f'(x) = \lim_{n \rightarrow \infty} f'_n(x) \quad \forall x \in [a, b].$$

Proof: Let  $\varepsilon > 0$ ,  $\exists N \in \mathbb{N}$  s.t.

$$\left. \begin{aligned} \|f'_n - f'_m\| &< \frac{\varepsilon}{2(b-a)} \\ \& \quad |f_n(x_0) - f_m(x_0)| &< \frac{\varepsilon}{2} \end{aligned} \right\} \quad \forall m, n \geq N, \quad \text{--- } (*)$$

Fix  $m, n \geq N$ ,

& fix  $x \in [a, b] \setminus \{x_0\}$ . By MVT ( $\because f_n - f_m$  is diff.)

$\exists \xi$  between  $x$  &  $x_0$  s.t. (applied to  $f_n - f_m$ )

$$(f_n - f_m)(x) - (f_n - f_m)(x_0) = (f_n - f_m)'(\xi) * (x - x_0).$$

$$\Rightarrow f_n(x) - f_m(x) = (f_n(x_0) - f_m(x_0)) + (f'_n(\xi) - f'_m(\xi))(x - x_0).$$

$$\Rightarrow |f_n(x) - f_m(x)| \leq |f_n(x_0) - f_m(x_0)| + |x - x_0| |f'_n(\xi) - f'_m(\xi)|$$

$\Delta$ -ineq.

$$\stackrel{\text{by } (*)}{<} \frac{\varepsilon}{2} + \frac{\varepsilon}{2(b-a)} \times |x - x_0|$$

$$\leq \frac{\varepsilon}{2} + \frac{\varepsilon}{2} \quad \left[ \because |x - x_0| \leq b - a \right]$$

$$\Rightarrow |f_n(x) - f_m(x)| < \varepsilon \quad \forall n, m \geq N \text{ \& } x \in [a, b].$$

(xx)

For  $x = x_0$  it is true anyway.

$$\text{i.e. } \|f_n - f_m\| < \varepsilon \quad \forall n, m \geq N.$$

$$\Rightarrow \{f_n\} \text{ is u.c. on } [a, b].$$

$$\text{Let } \lim_{n \rightarrow \infty} f_n = f \text{ on } [a, b].$$

Claim:  $f$  is diff. on  $[a, b]$  &  $f'(x) = \lim_{n \rightarrow \infty} f'_n(x)$   
 $\forall x \in [a, b].$

Fix  $\tilde{x} \in [a, b]$ . Set  $\xrightarrow{\text{Next page}}$

$$\tilde{f}_n(x) := \begin{cases} \frac{f_n(x) - f_n(\tilde{x})}{x - \tilde{x}} & \forall x \neq \tilde{x} \\ f'_n(\tilde{x}) & \text{if } x = \tilde{x} \end{cases}$$

$$\tilde{f}(x) = \begin{cases} \frac{f(x) - f(\tilde{x})}{x - \tilde{x}} & \forall x \neq \tilde{x} \\ f' & \end{cases}$$

(25)

Fix  $\tilde{x} \in [a, b]$ .

$$\text{Set } F_n(x) := \frac{f_n(x) - f_n(\tilde{x})}{x - \tilde{x}} \quad \left. \vphantom{\frac{f_n(x) - f_n(\tilde{x})}{x - \tilde{x}}} \right\} \quad \forall n \quad \forall \underline{x \neq \tilde{x}}.$$

$$\nexists F(x) := \frac{f(x) - f(\tilde{x})}{x - \tilde{x}}$$

 $\therefore f_n$  is diff. on  $[a, b] \quad \forall n$ , it follows that

$$\lim_{x \rightarrow \tilde{x}} F_n(x) = f'_n(\tilde{x}) \quad \forall n.$$

Also, for each  $x \in [a, b] \setminus \{\tilde{x}\}$ , by MVT,  $\exists \xi$  between  $x$  &  $\tilde{x}$ 

$$\text{s.t.} \quad (f_n - f_m)(x) - (f_n - f_m)(\tilde{x}) = (x - \tilde{x}) \times (f_n - f_m)'(\xi).$$

$$\Rightarrow \quad \left( F_n(x) - F_m(x) \right) = f'_n(\xi) - f'_m(\xi).$$

$$(\text{Recall: } \|f'_n - f'_m\| < \varepsilon / 2(b-a)).$$

$$\therefore (*) \text{ implies, } |F_n(x) - F_m(x)| < \frac{\varepsilon}{2(b-a)} \quad \forall m, n \geq N \quad \forall \underline{x \neq \tilde{x}}.$$

$$\Rightarrow \{F_n\} \text{ is } \underline{\text{u.c.}} \text{ on } \underline{[a, b] \setminus \{\tilde{x}\}}.$$

$\Rightarrow$   
interchanging  
limits.

$$\lim_{n \rightarrow \infty} \underbrace{\lim_{x \rightarrow \tilde{x}} F_n(x)}_{f'_n(\tilde{x})} = \lim_{x \rightarrow \tilde{x}} \underbrace{\lim_{n \rightarrow \infty} F_n(x)}_{\text{?}}$$

$$\text{But } f_n \xrightarrow{u} f \quad \Rightarrow \quad F_n(x) \rightarrow \frac{f(x) - f(\tilde{x})}{x - \tilde{x}} = F(x) \quad \text{as } n \rightarrow \infty$$

$$\text{i.e. } \lim_{n \rightarrow \infty} F_n(x) = F(x), \quad \forall x \neq \tilde{x}.$$



$$\therefore \lim_{n \rightarrow \infty} \lim_{x \rightarrow \tilde{x}} F_n(x) = \lim_{x \rightarrow \tilde{x}} F(x).$$

$$\therefore (\star) \Rightarrow \lim_{n \rightarrow \infty} f'_n(\tilde{x}) = \lim_{x \rightarrow \tilde{x}} F(x). \quad \text{--- } \textcircled{+}$$

$\therefore F(x) = \frac{f(x) - f(\tilde{x})}{x - \tilde{x}}$  & since the LHS <sup>of  $\textcircled{+}$</sup>  ~~limit~~ exists,

it follows that  $\lim_{x \rightarrow \tilde{x}} F(x) = f'(\tilde{x})$  exists.

$$\therefore f'(\tilde{x}) = \lim_{n \rightarrow \infty} f'_n(\tilde{x}). \quad \square$$

Another exotic example:

$\{f_n\}$  be a diff. fn. &  $f_n \rightarrow f$  unif.

Suppose  $f$  is also diff. Still,  $f'_n \not\rightarrow f'$  in general!!

Eg:  $f_n(x) = \frac{\sin nx}{n}$   $x \in [0, 1], n \geq 1$

$\therefore \left| \frac{\sin nx}{n} \right| \leq \frac{1}{n}, \forall n$ , it follows that  $f_n \rightarrow f$  unif. on  $[0, 1]$ , where

$$\underline{f \equiv 0.}$$

$\therefore \underline{f' \equiv 0.}$  on  $[0, 1]$ . Also,  $f'_n(x) = \cos nx$ .  
 $\forall n, \& x \in [0, 1]$

But  $f'_n(x)$  does not converge for  $x \in (0, 1]$ .

$\therefore \underline{f'_n \not\rightarrow f'}$ . (~~even point~~)

$\square$

## Series of Functions.

Def. Let  $\{f_n\} \subseteq \mathcal{F}(S)$ . The formal sum of functions

$$f_1 + f_2 + \dots := \sum_{n \geq 1} f_n$$

is called a series of  $f_n$ 's.

#  $\forall x \in S$ , by  $\sum f_n(x)$ , we understand the formal sum  $f_1(x) + f_2(x) + \dots$

# Given  $\sum_{n=1}^{\infty} f_n$ ,  $\forall n \in \mathbb{N}$ , define the  $n$ -th partial sum of the series as:

$$S_n(x) = \sum_{k=1}^n f_k(x). \quad (\forall x \in S). \quad n \in \mathbb{N}.$$

↑ (Finite sum: So good to go.)

Def: Let  $f: S \rightarrow \mathbb{R}$  &  $\{f_n\}_{n \geq 1} \subseteq \mathcal{F}(S)$ . We say that

$f$  is the pointwise limit of  $\sum f_n$  if  $S_n \rightarrow f$  pointwise on  $S$ .

$\sum f_n = f$ , pointwise.  ~~$S_n \rightarrow f$  on  $S$~~  In this case, we write

If  $S_n \xrightarrow{u} f$  on  $S$ , then we say that  $\sum f_n = f$  uniformly on  $S$ . □

eg:  $S = [0, 1)$ .  $f_n(x) = x^n \quad \forall n, x \in [0, 1)$ .

$$\therefore S_n(x) = \sum_{m=0}^{n-1} x^m \Rightarrow S_n(x) = \frac{1-x^n}{1-x} \quad \forall x \in [0, 1).$$

$$\therefore \lim_{n \rightarrow \infty} S_n(x) = \frac{1}{1-x} \quad \forall x \in [0, 1).$$

$$\text{i.e.} \quad \sum_{n=0}^{\infty} x^n = \frac{1}{1-x} \quad \text{pointwise on } [0, 1).$$

However,  $\sum_{n=0}^{\infty} x^n = \frac{1}{1-x}$  is not uniform on  $[0,1)$ .

Indeed:  $\frac{1}{1-x} \notin \mathcal{B}[0,1)$ . However:

$$S_n \in \mathcal{B}[0,1) \quad \& \quad \lim_{n \rightarrow \infty} S_n(x) = \frac{1}{1-x}.$$



Thm: (Cauchy criterion) Let  $\{f_n\} \subseteq \mathcal{F}(S)$ . Then  $\sum f_n$  converges uniformly on  $S \iff$  for  $\varepsilon > 0 \exists N \in \mathbb{N}$  s.t.

$$\left\| \sum_{k=m+1}^n f_k \right\| < \varepsilon \quad \forall n > m \geq N.$$

Proof:  $\because S_n = \sum_{k=1}^n f_k$ , it follows that

$$S_n - S_m = \sum_{k=m+1}^n f_k \quad \forall n > m.$$

Then  $\sum f_n$  is u.c.  $\iff \{S_n\}$  is u.c.  $\iff$  Cauchy criterion for seq. of f.n.s. □

Cor: If  $\sum f_n$  converges uniformly on  $S$ , then  $\|f_n\| \rightarrow 0$  as  $n \rightarrow \infty$ .

Proof:  $\because f_n = S_{n+1} - S_n \quad \forall n$ , this follows from the above theorem. □

eg: Consider  $0 < \varepsilon < 1$  &  $\sum_{n=0}^{\infty} x^n$ . Let  $n > m, x \in [-\varepsilon, \varepsilon]$ .

$$\begin{aligned} \text{Now } |S_n(x) - S_m(x)| &= \frac{1}{|1-x|} \times |(1-x^n) - (1-x^m)| \\ &= \frac{|x^n - x^m|}{|1-x|} \leq \frac{2|x|^m}{|1-x|} \quad \left( \because n > m \text{ \& } |x| < 1 \right) \end{aligned}$$