

# LINEAR ALGEBRA -II

B V Rajarama Bhat

Indian Statistical Institute, Bangalore

# Lecture 15: Algebraic description of projections

- ▶ We recall: [Definition 12.1](#). Let  $S$  be a non-empty subset of an inner product space  $V$ . Then the [orthogonal complement](#) of  $S$  is defined as:

$$S^\perp = \{v \in V : \langle x, v \rangle = 0, \quad \forall x \in S\}.$$

# Lecture 15: Algebraic description of projections

- ▶ We recall: **Definition 12.1.** Let  $S$  be a non-empty subset of an inner product space  $V$ . Then the **orthogonal complement** of  $S$  is defined as:

$$S^\perp = \{v \in V : \langle x, v \rangle = 0, \quad \forall x \in S\}.$$

- ▶ **Example 12.2:** Consider  $S \subset \mathbb{R}^3$  where

$$S = \left\{ \begin{pmatrix} 1 \\ -1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix} \right\}$$

## Lecture 15: Algebraic description of projections

- ▶ We recall: **Definition 12.1.** Let  $S$  be a non-empty subset of an inner product space  $V$ . Then the **orthogonal complement** of  $S$  is defined as:

$$S^\perp = \{v \in V : \langle x, v \rangle = 0, \quad \forall x \in S\}.$$

- ▶ **Example 12.2:** Consider  $S \subset \mathbb{R}^3$  where

$$S = \left\{ \begin{pmatrix} 1 \\ -1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix} \right\}$$

- ▶ Then

$$S^\perp = \left\{ \begin{pmatrix} c \\ c \\ c \end{pmatrix} : c \in \mathbb{R} \right\}.$$

## Continuation

- **Proposition 12.2:** Let  $S$  be a non-empty subset of an inner product space  $V$ . Then  $S^\perp$  is a subspace of  $V$ . Further,  $(S^\perp)^\perp$  is a subspace containing  $S$ .

# Continuation

- ▶ **Proposition 12.2:** Let  $S$  be a non-empty subset of an inner product space  $V$ . Then  $S^\perp$  is a subspace of  $V$ . Further,  $(S^\perp)^\perp$  is a subspace containing  $S$ .
- ▶ **Proof:** We recall the definition of  $S^\perp$ :

$$S^\perp = \{v \in V : \langle x, v \rangle = 0, \quad \forall x \in S\}.$$

# Continuation

- ▶ **Proposition 12.2:** Let  $S$  be a non-empty subset of an inner product space  $V$ . Then  $S^\perp$  is a subspace of  $V$ . Further,  $(S^\perp)^\perp$  is a subspace containing  $S$ .
- ▶ **Proof:** We recall the definition of  $S^\perp$ :

$$S^\perp = \{v \in V : \langle x, v \rangle = 0, \quad \forall x \in S\}.$$

- ▶ Now if  $v, w \in S^\perp$  and  $c, d \in \mathbb{F}$ : For  $x \in S$ ,

$$\langle x, cv + dw \rangle = c\langle x, v \rangle + d\langle x, w \rangle = c \cdot 0 + d \cdot 0 = 0.$$

# Continuation

- ▶ **Proposition 12.2:** Let  $S$  be a non-empty subset of an inner product space  $V$ . Then  $S^\perp$  is a subspace of  $V$ . Further,  $(S^\perp)^\perp$  is a subspace containing  $S$ .
- ▶ **Proof:** We recall the definition of  $S^\perp$ :

$$S^\perp = \{v \in V : \langle x, v \rangle = 0, \quad \forall x \in S\}.$$

- ▶ Now if  $v, w \in S^\perp$  and  $c, d \in \mathbb{F}$ : For  $x \in S$ ,

$$\langle x, cv + dw \rangle = c\langle x, v \rangle + d\langle x, w \rangle = c \cdot 0 + d \cdot 0 = 0.$$

- ▶ Hence  $cv + dw \in S^\perp$ . This proves that  $S^\perp$  is a subspace of  $V$ .



# Continuation

- ▶ **Proposition 12.2:** Let  $S$  be a non-empty subset of an inner product space  $V$ . Then  $S^\perp$  is a subspace of  $V$ . Further,  $(S^\perp)^\perp$  is a subspace containing  $S$ .
- ▶ **Proof:** We recall the definition of  $S^\perp$ :

$$S^\perp = \{v \in V : \langle x, v \rangle = 0, \quad \forall x \in S\}.$$

- ▶ Now if  $v, w \in S^\perp$  and  $c, d \in \mathbb{F}$ : For  $x \in S$ ,

$$\langle x, cv + dw \rangle = c\langle x, v \rangle + d\langle x, w \rangle = c \cdot 0 + d \cdot 0 = 0.$$

- ▶ Hence  $cv + dw \in S^\perp$ . This proves that  $S^\perp$  is a subspace of  $V$ .
- ▶ It is easy to see that if  $x \in S$  then  $x \in (S^\perp)^\perp$ . Therefore  $S \subseteq (S^\perp)^\perp$ .

# Continuation

- ▶ **Proposition 12.2:** Let  $S$  be a non-empty subset of an inner product space  $V$ . Then  $S^\perp$  is a subspace of  $V$ . Further,  $(S^\perp)^\perp$  is a subspace containing  $S$ .
- ▶ **Proof:** We recall the definition of  $S^\perp$ :

$$S^\perp = \{v \in V : \langle x, v \rangle = 0, \quad \forall x \in S\}.$$

- ▶ Now if  $v, w \in S^\perp$  and  $c, d \in \mathbb{F}$ : For  $x \in S$ ,

$$\langle x, cv + dw \rangle = c\langle x, v \rangle + d\langle x, w \rangle = c \cdot 0 + d \cdot 0 = 0.$$

- ▶ Hence  $cv + dw \in S^\perp$ . This proves that  $S^\perp$  is a subspace of  $V$ .
- ▶ It is easy to see that if  $x \in S$  then  $x \in (S^\perp)^\perp$ . Therefore  $S \subseteq (S^\perp)^\perp$ .
- ▶ We have already seen that orthogonal complement of any non-empty subset is a subspace. In particular,  $(S^\perp)^\perp$  is a subspace.

- ▶ Consider  $V = \mathbb{R}^3$  with standard inner product.

## $\mathbb{R}^2$ in $\mathbb{R}^3$

- ▶ Consider  $V = \mathbb{R}^3$  with standard inner product.
- ▶ Consider the subspace

$$V_0 = \left\{ \begin{pmatrix} x_1 \\ x_2 \\ 0 \end{pmatrix} : x_1, x_2 \in \mathbb{R} \right\}$$

## $\mathbb{R}^2$ in $\mathbb{R}^3$

- ▶ Consider  $V = \mathbb{R}^3$  with standard inner product.
- ▶ Consider the subspace

$$V_0 = \left\{ \begin{pmatrix} x_1 \\ x_2 \\ 0 \end{pmatrix} : x_1, x_2 \in \mathbb{R} \right\}$$

- ▶ Take  $V_1 = (V_0)^\perp$ .

## $\mathbb{R}^2$ in $\mathbb{R}^3$

- ▶ Consider  $V = \mathbb{R}^3$  with standard inner product.
- ▶ Consider the subspace

$$V_0 = \left\{ \begin{pmatrix} x_1 \\ x_2 \\ 0 \end{pmatrix} : x_1, x_2 \in \mathbb{R} \right\}$$

- ▶ Take  $V_1 = (V_0)^\perp$ .
- ▶ Clearly,

$$V_1 = \left\{ \begin{pmatrix} 0 \\ 0 \\ x_3 \end{pmatrix} : x_3 \in \mathbb{R} \right\}.$$

## Continuation

- ▶ We see that any vector  $x \in V$  decomposes uniquely as  $x = y + z$  with  $y \in V_0$  and  $z \in V_1$ .

# Continuation

- ▶ We see that any vector  $x \in V$  decomposes uniquely as  $x = y + z$  with  $y \in V_0$  and  $z \in V_1$ .
- ▶ Indeed for

$$x = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}$$

the only choice is:

$$y = \begin{pmatrix} x_1 \\ x_2 \\ 0 \end{pmatrix}; z = \begin{pmatrix} 0 \\ 0 \\ x_3 \end{pmatrix}.$$



# Continuation

- ▶ We see that any vector  $x \in V$  decomposes uniquely as  $x = y + z$  with  $y \in V_0$  and  $z \in V_1$ .
- ▶ Indeed for

$$x = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}$$

the only choice is:

$$y = \begin{pmatrix} x_1 \\ x_2 \\ 0 \end{pmatrix}; z = \begin{pmatrix} 0 \\ 0 \\ x_3 \end{pmatrix}.$$

- ▶ We want to show that this is a general phenomenon.

# Extending bases and orthonormal bases

- **Theorem 12.4:** Let  $V_0$  be a non-trivial subspace of a finite dimensional vector space  $V$ . Then any basis of  $V_0$  extends to a basis of  $V$ , that is, if  $\{v_1, v_2, \dots, v_k\}$  is a basis of  $V_0$  then there exists  $\{v_{k+1}, \dots, v_n\}$  such that  $\{v_1, \dots, v_n\}$  is a basis of  $V$ .

# Extending bases and orthonormal bases

- ▶ **Theorem 12.4:** Let  $V_0$  be a non-trivial subspace of a finite dimensional vector space  $V$ . Then any basis of  $V_0$  extends to a basis of  $V$ , that is, if  $\{v_1, v_2, \dots, v_k\}$  is a basis of  $V_0$  then there exists  $\{v_{k+1}, \dots, v_n\}$  such that  $\{v_1, \dots, v_n\}$  is a basis of  $V$ .
- ▶ **Proof:** Take

$$M_k := \text{span} \{v_1, v_2, \dots, v_k\}$$

# Extending bases and orthonormal bases

- ▶ **Theorem 12.4:** Let  $V_0$  be a non-trivial subspace of a finite dimensional vector space  $V$ . Then any basis of  $V_0$  extends to a basis of  $V$ , that is, if  $\{v_1, v_2, \dots, v_k\}$  is a basis of  $V_0$  then there exists  $\{v_{k+1}, \dots, v_n\}$  such that  $\{v_1, \dots, v_n\}$  is a basis of  $V$ .

- ▶ **Proof:** Take

$$M_k := \text{span} \{v_1, v_2, \dots, v_k\}$$

- ▶ If  $M_k = V$  then  $V_0 = V$ ,  $\{v_1, \dots, v_k\}$  is a basis for  $V$  and so no extension is required.

# Extending bases and orthonormal bases

- ▶ **Theorem 12.4:** Let  $V_0$  be a non-trivial subspace of a finite dimensional vector space  $V$ . Then any basis of  $V_0$  extends to a basis of  $V$ , that is, if  $\{v_1, v_2, \dots, v_k\}$  is a basis of  $V_0$  then there exists  $\{v_{k+1}, \dots, v_n\}$  such that  $\{v_1, \dots, v_n\}$  is a basis of  $V$ .

- ▶ **Proof:** Take

$$M_k := \text{span} \{v_1, v_2, \dots, v_k\}$$

- ▶ If  $M_k = V$  then  $V_0 = V$ ,  $\{v_1, \dots, v_k\}$  is a basis for  $V$  and so no extension is required.
- ▶ If not, choose any  $v_{k+1} \in V \setminus M_k$ . Then  $\{v_1, \dots, v_{k+1}\}$  is a linearly independent set (Why?). Take

$$M_{k+1} := \text{span}\{v_1, \dots, v_{k+1}\}.$$

# Extending bases and orthonormal bases

- ▶ **Theorem 12.4:** Let  $V_0$  be a non-trivial subspace of a finite dimensional vector space  $V$ . Then any basis of  $V_0$  extends to a basis of  $V$ , that is, if  $\{v_1, v_2, \dots, v_k\}$  is a basis of  $V_0$  then there exists  $\{v_{k+1}, \dots, v_n\}$  such that  $\{v_1, \dots, v_n\}$  is a basis of  $V$ .

- ▶ **Proof:** Take

$$M_k := \text{span} \{v_1, v_2, \dots, v_k\}$$

- ▶ If  $M_k = V$  then  $V_0 = V$ ,  $\{v_1, \dots, v_k\}$  is a basis for  $V$  and so no extension is required.
- ▶ If not, choose any  $v_{k+1} \in V \setminus M_k$ . Then  $\{v_1, \dots, v_{k+1}\}$  is a linearly independent set (Why?). Take

$$M_{k+1} := \text{span}\{v_1, \dots, v_{k+1}\}.$$

- ▶ If  $V = M_{k+1}$  then  $\{v_1, \dots, v_{k+1}\}$  is a basis for  $V$  and we are done. If not, take  $v_{k+2} \in V \setminus M_{k+1}$  and continue the induction process.

- ▶ The process terminates after a finite number of steps as  $V$  is finite dimensional and so it can have at most  $\dim(V)$  linearly independent elements.

- ▶ The process terminates after a finite number of steps as  $V$  is finite dimensional and so it can have at most  $\dim(V)$  linearly independent elements.
- ▶ Therefore  $V = M_n$  for some  $n$  and  $\{v_1, \dots, v_n\}$  is a basis for  $V$ .



# Extending orthonormal bases

- **Theorem 12.5:** Let  $V_0$  be a non-trivial subspace of a finite dimensional inner product space  $V$ . Then any orthonormal basis of  $V_0$  extends to an orthonormal basis of  $V$ , that is, if  $\{v_1, v_2, \dots, v_k\}$  is an orthonormal basis of  $V_0$  then there exists  $\{v_{k+1}, \dots, v_n\}$  such that  $\{v_1, \dots, v_n\}$  is an orthonormal basis of  $V$ .

# Extending orthonormal bases

- ▶ **Theorem 12.5:** Let  $V_0$  be a non-trivial subspace of a finite dimensional inner product space  $V$ . Then any orthonormal basis of  $V_0$  extends to an orthonormal basis of  $V$ , that is, if  $\{v_1, v_2, \dots, v_k\}$  is an orthonormal basis of  $V_0$  then there exists  $\{v_{k+1}, \dots, v_n\}$  such that  $\{v_1, \dots, v_n\}$  is an orthonormal basis of  $V$ .
- ▶ **Proof:** By the previous theorem we may extend  $\{v_1, \dots, v_k\}$  to a basis  $\{v_1, \dots, v_k, w_{k+1}, \dots, w_n\}$  of  $V$ .

# Extending orthonormal bases

- ▶ **Theorem 12.5:** Let  $V_0$  be a non-trivial subspace of a finite dimensional inner product space  $V$ . Then any orthonormal basis of  $V_0$  extends to an orthonormal basis of  $V$ , that is, if  $\{v_1, v_2, \dots, v_k\}$  is an orthonormal basis of  $V_0$  then there exists  $\{v_{k+1}, \dots, v_n\}$  such that  $\{v_1, \dots, v_n\}$  is an orthonormal basis of  $V$ .
- ▶ **Proof:** By the previous theorem we may extend  $\{v_1, \dots, v_k\}$  to a basis  $\{v_1, \dots, v_k, w_{k+1}, \dots, w_n\}$  of  $V$ .
- ▶ Now apply the Gram-Schmidt procedure on  $\{v_1, \dots, v_k, w_{k+1}, \dots, w_n\}$  to get an orthonormal basis  $\{e_1, \dots, e_n\}$  of  $V$ .

# Extending orthonormal bases

- ▶ **Theorem 12.5:** Let  $V_0$  be a non-trivial subspace of a finite dimensional inner product space  $V$ . Then any orthonormal basis of  $V_0$  extends to an orthonormal basis of  $V$ , that is, if  $\{v_1, v_2, \dots, v_k\}$  is an orthonormal basis of  $V_0$  then there exists  $\{v_{k+1}, \dots, v_n\}$  such that  $\{v_1, \dots, v_n\}$  is an orthonormal basis of  $V$ .
- ▶ **Proof:** By the previous theorem we may extend  $\{v_1, \dots, v_k\}$  to a basis  $\{v_1, \dots, v_k, w_{k+1}, \dots, w_n\}$  of  $V$ .
- ▶ Now apply the Gram-Schmidt procedure on  $\{v_1, \dots, v_k, w_{k+1}, \dots, w_n\}$  to get an orthonormal basis  $\{e_1, \dots, e_n\}$  of  $V$ .
- ▶ It is an elementary exercise to see that  $e_j = v_j$  for  $1 \leq j \leq k$  as  $v_1, \dots, v_k$  are already orthonormal. ■

# Orthogonal complement of a subspace

- ▶ Consider the set up as above, that is,  $V_0$  is a non-trivial subspace of a finite dimensional inner product space  $V$ . Suppose  $\{v_1, \dots, v_k\}$  is an orthonormal basis of  $V_0$  and  $\{v_1, \dots, v_n\}$  is an orthonormal basis of  $V$ .

# Orthogonal complement of a subspace

- ▶ Consider the set up as above, that is,  $V_0$  is a non-trivial subspace of a finite dimensional inner product space  $V$ . Suppose  $\{v_1, \dots, v_k\}$  is an orthonormal basis of  $V_0$  and  $\{v_1, \dots, v_n\}$  is an orthonormal basis of  $V$ .
- ▶ Take

$$V_1 = \text{span} \{v_{k+1}, \dots, v_n\}.$$

# Orthogonal complement of a subspace

- ▶ Consider the set up as above, that is,  $V_0$  is a non-trivial subspace of a finite dimensional inner product space  $V$ . Suppose  $\{v_1, \dots, v_k\}$  is an orthonormal basis of  $V_0$  and  $\{v_1, \dots, v_n\}$  is an orthonormal basis of  $V$ .
- ▶ Take

$$V_1 = \text{span} \{v_{k+1}, \dots, v_n\}.$$

- ▶ We claim that  $V_1 = (V_0)^\perp$  and  $\{v_{k+1}, \dots, v_n\}$  is an ortho-normal basis of  $V_1$ .

# Orthogonal complement of a subspace

- ▶ Consider the set up as above, that is,  $V_0$  is a non-trivial subspace of a finite dimensional inner product space  $V$ . Suppose  $\{v_1, \dots, v_k\}$  is an orthonormal basis of  $V_0$  and  $\{v_1, \dots, v_n\}$  is an orthonormal basis of  $V$ .

- ▶ Take

$$V_1 = \text{span} \{v_{k+1}, \dots, v_n\}.$$

- ▶ We claim that  $V_1 = (V_0)^\perp$  and  $\{v_{k+1}, \dots, v_n\}$  is an ortho-normal basis of  $V_1$ .
- ▶ The second part is obvious. We only need to prove  $V_1 = (V_0)^\perp$ .



# Orthogonal complement of a subspace

- ▶ Consider the set up as above, that is,  $V_0$  is a non-trivial subspace of a finite dimensional inner product space  $V$ . Suppose  $\{v_1, \dots, v_k\}$  is an orthonormal basis of  $V_0$  and  $\{v_1, \dots, v_n\}$  is an orthonormal basis of  $V$ .

- ▶ Take

$$V_1 = \text{span} \{v_{k+1}, \dots, v_n\}.$$

- ▶ We claim that  $V_1 = (V_0)^\perp$  and  $\{v_{k+1}, \dots, v_n\}$  is an ortho-normal basis of  $V_1$ .
- ▶ The second part is obvious. We only need to prove  $V_1 = (V_0)^\perp$ .
- ▶ Note that  $\langle v_i, v_j \rangle = 0$  for all  $1 \leq i \leq k$  and  $(k+1) \leq j \leq n$

# Orthogonal complement of a subspace

- ▶ Consider the set up as above, that is,  $V_0$  is a non-trivial subspace of a finite dimensional inner product space  $V$ . Suppose  $\{v_1, \dots, v_k\}$  is an orthonormal basis of  $V_0$  and  $\{v_1, \dots, v_n\}$  is an orthonormal basis of  $V$ .

- ▶ Take

$$V_1 = \text{span} \{v_{k+1}, \dots, v_n\}.$$

- ▶ We claim that  $V_1 = (V_0)^\perp$  and  $\{v_{k+1}, \dots, v_n\}$  is an ortho-normal basis of  $V_1$ .
- ▶ The second part is obvious. We only need to prove  $V_1 = (V_0)^\perp$ .
- ▶ Note that  $\langle v_i, v_j \rangle = 0$  for all  $1 \leq i \leq k$  and  $(k+1) \leq j \leq n$
- ▶ Therefore  $\langle \sum_{i=1}^k c_i v_i, \sum_{j=(k+1)}^n c_j v_j \rangle$  for any scalars  $c_1, \dots, c_n$ .

# Orthogonal complement of a subspace

- ▶ Consider the set up as above, that is,  $V_0$  is a non-trivial subspace of a finite dimensional inner product space  $V$ . Suppose  $\{v_1, \dots, v_k\}$  is an orthonormal basis of  $V_0$  and  $\{v_1, \dots, v_n\}$  is an orthonormal basis of  $V$ .

- ▶ Take

$$V_1 = \text{span} \{v_{k+1}, \dots, v_n\}.$$

- ▶ We claim that  $V_1 = (V_0)^\perp$  and  $\{v_{k+1}, \dots, v_n\}$  is an ortho-normal basis of  $V_1$ .
- ▶ The second part is obvious. We only need to prove  $V_1 = (V_0)^\perp$ .
- ▶ Note that  $\langle v_i, v_j \rangle = 0$  for all  $1 \leq i \leq k$  and  $(k+1) \leq j \leq n$
- ▶ Therefore  $\langle \sum_{i=1}^k c_i v_i, \sum_{j=(k+1)}^n c_j v_j \rangle$  for any scalars  $c_1, \dots, c_n$ .

# Orthogonal complement of a subspace

- ▶ Consider the set up as above, that is,  $V_0$  is a non-trivial subspace of a finite dimensional inner product space  $V$ . Suppose  $\{v_1, \dots, v_k\}$  is an orthonormal basis of  $V_0$  and  $\{v_1, \dots, v_n\}$  is an orthonormal basis of  $V$ .

- ▶ Take

$$V_1 = \text{span} \{v_{k+1}, \dots, v_n\}.$$

- ▶ We claim that  $V_1 = (V_0)^\perp$  and  $\{v_{k+1}, \dots, v_n\}$  is an ortho-normal basis of  $V_1$ .
- ▶ The second part is obvious. We only need to prove  $V_1 = (V_0)^\perp$ .
- ▶ Note that  $\langle v_i, v_j \rangle = 0$  for all  $1 \leq i \leq k$  and  $(k+1) \leq j \leq n$
- ▶ Therefore  $\langle \sum_{i=1}^k c_i v_i, \sum_{j=(k+1)}^n c_j v_j \rangle$  for any scalars  $c_1, \dots, c_n$ .
- ▶ This shows  $\langle x, y \rangle = 0$  for all  $x \in V_0$  and  $y \in V_1$ . Hence  $V_1 \subseteq (V_0)^\perp$ .

# Continuation

- ▶ Suppose  $x \in V_0^\perp$ .

## Continuation

- ▶ Suppose  $x \in V_0^\perp$ .
- ▶ As  $\{v_1, \dots, v_n\}$  is an orthonormal basis of  $V$ , we get
$$x = \sum_{j=1}^n \langle v_j, x \rangle v_j.$$

# Continuation

- ▶ Suppose  $x \in V_0^\perp$ .
- ▶ As  $\{v_1, \dots, v_n\}$  is an orthonormal basis of  $V$ , we get  $x = \sum_{j=1}^n \langle v_j, x \rangle v_j$ .
- ▶ As  $x$  is orthogonal to  $V_0$ , we get  $\langle v_j, x \rangle = 0$  for  $1 \leq j \leq k$ .

# Continuation

- ▶ Suppose  $x \in V_0^\perp$ .
- ▶ As  $\{v_1, \dots, v_n\}$  is an orthonormal basis of  $V$ , we get  $x = \sum_{j=1}^n \langle v_j, x \rangle v_j$ .
- ▶ As  $x$  is orthogonal to  $V_0$ , we get  $\langle v_j, x \rangle = 0$  for  $1 \leq j \leq k$ .
- ▶ Hence  $x = \sum_{j=k+1}^n \langle v_j, x \rangle v_j$  and therefore  $x \in V_1$ .



# Continuation

- ▶ Suppose  $x \in V_0^\perp$ .
- ▶ As  $\{v_1, \dots, v_n\}$  is an orthonormal basis of  $V$ , we get  $x = \sum_{j=1}^n \langle v_j, x \rangle v_j$ .
- ▶ As  $x$  is orthogonal to  $V_0$ , we get  $\langle v_j, x \rangle = 0$  for  $1 \leq j \leq k$ .
- ▶ Hence  $x = \sum_{j=k+1}^n \langle v_j, x \rangle v_j$  and therefore  $x \in V_1$ .
- ▶ This proves  $(V_0)^\perp \subseteq V_1$  and completes the proof of our claim.

# Projection theorem

- **Theorem 12.6:** Let  $V_0$  be a subspace of a finite dimensional inner product space  $V$ . Then every  $x \in V$  decomposes uniquely as

$$x = y + z$$

where  $y \in V_0$  and  $z \in V_0^\perp$ .

# Projection theorem

- **Theorem 12.6:** Let  $V_0$  be a subspace of a finite dimensional inner product space  $V$ . Then every  $x \in V$  decomposes uniquely as

$$x = y + z$$

where  $y \in V_0$  and  $z \in V_0^\perp$ .

- **Proof:** Suppose  $V_0 = \{0\}$ . Then  $V_0^\perp = V$  and we can decompose  $x$  as  $x = 0 + x$ , with  $0 \in V_0$  and  $x \in V_0^\perp$ .

# Projection theorem

- ▶ **Theorem 12.6:** Let  $V_0$  be a subspace of a finite dimensional inner product space  $V$ . Then every  $x \in V$  decomposes uniquely as

$$x = y + z$$

where  $y \in V_0$  and  $z \in V_0^\perp$ .

- ▶ **Proof:** Suppose  $V_0 = \{0\}$ . Then  $V_0^\perp = V$  and we can decompose  $x$  as  $x = 0 + x$ , with  $0 \in V_0$  and  $x \in V_0^\perp$ .
- ▶ If  $V_0 \neq \{0\}$ , choose an orthonormal basis  $\{v_1, \dots, v_k\}$  for  $V_0$ . Extend it to an orthonormal basis  $\{v_1, \dots, v_n\}$  of  $V$ .

# Projection theorem

- ▶ **Theorem 12.6:** Let  $V_0$  be a subspace of a finite dimensional inner product space  $V$ . Then every  $x \in V$  decomposes uniquely as

$$x = y + z$$

where  $y \in V_0$  and  $z \in V_0^\perp$ .

- ▶ **Proof:** Suppose  $V_0 = \{0\}$ . Then  $V_0^\perp = V$  and we can decompose  $x$  as  $x = 0 + x$ , with  $0 \in V_0$  and  $x \in V_0^\perp$ .
- ▶ If  $V_0 \neq \{0\}$ , choose an orthonormal basis  $\{v_1, \dots, v_k\}$  for  $V_0$ . Extend it to an orthonormal basis  $\{v_1, \dots, v_n\}$  of  $V$ .
- ▶ Now we know that any  $x \in V$  decomposes as

$$x = \sum_{j=1}^n \langle v_j, x \rangle v_j$$

# Uniqueness

► Take

$$y = \sum_{j=1}^k \langle v_j, x \rangle v_j$$

and

$$z = \sum_{j=(k+1)}^n \langle v_j, x \rangle v_j.$$

# Uniqueness

- ▶ Take

$$y = \sum_{j=1}^k \langle v_j, x \rangle v_j$$

and

$$z = \sum_{j=(k+1)}^n \langle v_j, x \rangle v_j.$$

- ▶ Clearly  $y \in V_0$  and  $z \in V_0^\perp$ . This proves the existence.

# Uniqueness

- ▶ Take

$$y = \sum_{j=1}^k \langle v_j, x \rangle v_j$$

and

$$z = \sum_{j=(k+1)}^n \langle v_j, x \rangle v_j.$$

- ▶ Clearly  $y \in V_0$  and  $z \in V_0^\perp$ . This proves the existence.
- ▶ Suppose  $x = y + z$  and  $x = y' + z'$  are two decompositions of  $x$  with  $y, y' \in V_0$  and  $z, z' \in V_0^\perp$ .



# Uniqueness

- ▶ Take

$$y = \sum_{j=1}^k \langle v_j, x \rangle v_j$$

and

$$z = \sum_{j=(k+1)}^n \langle v_j, x \rangle v_j.$$

- ▶ Clearly  $y \in V_0$  and  $z \in V_0^\perp$ . This proves the existence.
- ▶ Suppose  $x = y + z$  and  $x = y' + z'$  are two decompositions of  $x$  with  $y, y' \in V_0$  and  $z, z' \in V_0^\perp$ .
- ▶ We have,

$$y + z = y' + z'.$$

# Uniqueness

- ▶ Take

$$y = \sum_{j=1}^k \langle v_j, x \rangle v_j$$

and

$$z = \sum_{j=(k+1)}^n \langle v_j, x \rangle v_j.$$

- ▶ Clearly  $y \in V_0$  and  $z \in V_0^\perp$ . This proves the existence.
- ▶ Suppose  $x = y + z$  and  $x = y' + z'$  are two decompositions of  $x$  with  $y, y' \in V_0$  and  $z, z' \in V_0^\perp$ .
- ▶ We have,

$$y + z = y' + z'.$$

- ▶ Therefore  $y - y' = z' - z$ . As  $y, y' \in V_0$ ,  $y - y' \in V_0$ .

# Uniqueness

- ▶ Take

$$y = \sum_{j=1}^k \langle v_j, x \rangle v_j$$

and

$$z = \sum_{j=(k+1)}^n \langle v_j, x \rangle v_j.$$

- ▶ Clearly  $y \in V_0$  and  $z \in V_0^\perp$ . This proves the existence.
- ▶ Suppose  $x = y + z$  and  $x = y' + z'$  are two decompositions of  $x$  with  $y, y' \in V_0$  and  $z, z' \in V_0^\perp$ .
- ▶ We have,

$$y + z = y' + z'.$$

- ▶ Therefore  $y - y' = z' - z$ . As  $y, y' \in V_0$ ,  $y - y' \in V_0$ .
- ▶ Also as  $z, z' \in V_0^\perp$ ,  $y - y' = z' - z \in V_0^\perp$ .

# Uniqueness

- ▶ Take

$$y = \sum_{j=1}^k \langle v_j, x \rangle v_j$$

and

$$z = \sum_{j=(k+1)}^n \langle v_j, x \rangle v_j.$$

- ▶ Clearly  $y \in V_0$  and  $z \in V_0^\perp$ . This proves the existence.
- ▶ Suppose  $x = y + z$  and  $x = y' + z'$  are two decompositions of  $x$  with  $y, y' \in V_0$  and  $z, z' \in V_0^\perp$ .
- ▶ We have,

$$y + z = y' + z'.$$

- ▶ Therefore  $y - y' = z' - z$ . As  $y, y' \in V_0$ ,  $y - y' \in V_0$ .
- ▶ Also as  $z, z' \in V_0^\perp$ ,  $y - y' = z' - z \in V_0^\perp$ .
- ▶ Hence  $\langle y - y', y - y' \rangle = 0$ . Consequently  $y = y'$  and  $z' = z$ .  
This proves the uniqueness.

## A special case

- ▶ Suppose  $V$  is a finite dimensional inner product space and let  $y$  be a non-zero vector in  $V$ .

## A special case

- ▶ Suppose  $V$  is a finite dimensional inner product space and let  $y$  be a non-zero vector in  $V$ .
- ▶ Consider the one dimensional space  $V_0 = \{cy : c \in \mathbb{F}\}$ .

## A special case

- ▶ Suppose  $V$  is a finite dimensional inner product space and let  $y$  be a non-zero vector in  $V$ .
- ▶ Consider the one dimensional space  $V_0 = \{cy : c \in \mathbb{F}\}$ .
- ▶ Now  $\{v\}$  is an ortho-normal basis for  $V_0$  where

$$v = \frac{y}{\|y\|}.$$

## A special case

- ▶ Suppose  $V$  is a finite dimensional inner product space and let  $y$  be a non-zero vector in  $V$ .
- ▶ Consider the one dimensional space  $V_0 = \{cy : c \in \mathbb{F}\}$ .
- ▶ Now  $\{v\}$  is an ortho-normal basis for  $V_0$  where

$$v = \frac{y}{\|y\|}.$$

- ▶ Therefore any  $x \in V$  decomposes as  $x = \langle v, x \rangle v + z$  where  $z$  is orthogonal to  $v$ .



## A special case

- ▶ Suppose  $V$  is a finite dimensional inner product space and let  $y$  be a non-zero vector in  $V$ .
- ▶ Consider the one dimensional space  $V_0 = \{cy : c \in \mathbb{F}\}$ .
- ▶ Now  $\{v\}$  is an ortho-normal basis for  $V_0$  where

$$v = \frac{y}{\|y\|}.$$

- ▶ Therefore any  $x \in V$  decomposes as  $x = \langle v, x \rangle v + z$  where  $z$  is orthogonal to  $v$ .
- ▶ As shown in the previous lecture this is related to Cauchy-Schwarz inequality.

# Example

- **Example 13.1:** Let  $V = \mathbb{R}^n$  with the standard inner product.  
Let  $V_0 = \{x \in \mathbb{R}^n : \sum_{i=1}^n x_i = 0\}$ .

# Example

- ▶ **Example 13.1:** Let  $V = \mathbb{R}^n$  with the standard inner product. Let  $V_0 = \{x \in \mathbb{R}^n : \sum_{i=1}^n x_i = 0\}$ .
- ▶ We first analyze the case when  $n = 3$ . Now  $V = \mathbb{R}^3$  and

$$V_0 = \left\{ \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} : x_1 + x_2 + x_3 = 0 \right\}.$$

# Example

- ▶ **Example 13.1:** Let  $V = \mathbb{R}^n$  with the standard inner product. Let  $V_0 = \{x \in \mathbb{R}^n : \sum_{i=1}^n x_i = 0\}$ .
- ▶ We first analyze the case when  $n = 3$ . Now  $V = \mathbb{R}^3$  and

$$V_0 = \left\{ \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} : x_1 + x_2 + x_3 = 0 \right\}.$$

- ▶ One can see that

$$\left\{ \begin{pmatrix} 1 \\ -1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix} \right\}$$

is a basis for  $V_0$ .

## Example

- ▶ **Example 13.1:** Let  $V = \mathbb{R}^n$  with the standard inner product. Let  $V_0 = \{x \in \mathbb{R}^n : \sum_{i=1}^n x_i = 0\}$ .
- ▶ We first analyze the case when  $n = 3$ . Now  $V = \mathbb{R}^3$  and

$$V_0 = \left\{ \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} : x_1 + x_2 + x_3 = 0 \right\}.$$

- ▶ One can see that

$$\left\{ \begin{pmatrix} 1 \\ -1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix} \right\}$$

is a basis for  $V_0$ .

- ▶ Let us apply Gram-Schmidt on this to get an orthonormal basis for  $V_0$ .

## Continuation

- ▶ We get the first vector as

$$v_1 = \begin{pmatrix} 1/\sqrt{2} \\ -1/\sqrt{2} \\ 0 \end{pmatrix}.$$

# Continuation

- ▶ We get the first vector as

$$v_1 = \begin{pmatrix} 1/\sqrt{2} \\ -1/\sqrt{2} \\ 0 \end{pmatrix}.$$

- ▶ Now take

$$\begin{aligned} w_2 &= \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix} - \left\langle \begin{pmatrix} 1/\sqrt{2} \\ -1/\sqrt{2} \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix} \right\rangle \begin{pmatrix} 1/\sqrt{2} \\ -1/\sqrt{2} \\ 0 \end{pmatrix} \\ &= \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix} - \begin{pmatrix} 1/2 \\ -1/2 \\ 0 \end{pmatrix} \\ &= \begin{pmatrix} 1/2 \\ 1/2 \\ -1 \end{pmatrix}. \end{aligned}$$

# Continuation

► Now

$$v_2 = \frac{w_2}{\|w_2\|} = \begin{pmatrix} 1/\sqrt{6} \\ 1/\sqrt{6} \\ -2/\sqrt{6} \end{pmatrix}$$



## Continuation

- ▶ Now

$$v_2 = \frac{w_2}{\|w_2\|} = \begin{pmatrix} 1/\sqrt{6} \\ 1/\sqrt{6} \\ -2/\sqrt{6} \end{pmatrix}$$

- ▶  $\{v_1, v_2\}$  is an ortho-normal basis for  $V_0$ .

## Continuation

► Now

$$v_2 = \frac{w_2}{\|w_2\|} = \begin{pmatrix} 1/\sqrt{6} \\ 1/\sqrt{6} \\ -2/\sqrt{6} \end{pmatrix}$$

- $\{v_1, v_2\}$  is an ortho-normal basis for  $V_0$ .
- Given  $x \in \mathbb{R}^3$ , it decomposes as  $y + z$ , where  $y \in V_0$ ,  $z \in V_0^\perp$ .

$$\begin{aligned} y &= \langle v_1, x \rangle v_1 + \langle v_2, x \rangle v_2 \\ &= \frac{x_1 - x_2}{\sqrt{2}} \begin{pmatrix} 1/\sqrt{2} \\ -1/\sqrt{2} \\ 0 \end{pmatrix} + \frac{(x_1 + x_2 - 2x_3)}{\sqrt{6}} \begin{pmatrix} 1/\sqrt{6} \\ 1/\sqrt{6} \\ -2/\sqrt{6} \end{pmatrix} \\ &= \frac{1}{3} \begin{pmatrix} 2x_1 - x_2 - x_3 \\ -x_1 + 2x_2 - x_3 \\ -x_1 - x_2 + 2x_3 \end{pmatrix} \end{aligned}$$

## Continuation

$$\blacktriangleright z = \frac{1}{3} \begin{pmatrix} x_1 + x_2 + x_3 \\ x_1 + x_2 + x_3 \\ x_1 + x_2 + x_3 \end{pmatrix}.$$

# Continuation

►  $z = \frac{1}{3} \begin{pmatrix} x_1 + x_2 + x_3 \\ x_1 + x_2 + x_3 \\ x_1 + x_2 + x_3 \end{pmatrix}.$

► For general  $n$ , with  $\bar{x} = \frac{1}{n}(x_1 + x_2 + \cdots + x_n)$ ,

$$y = \begin{pmatrix} x_1 - \bar{x} \\ x_2 - \bar{x} \\ \vdots \\ x_n - \bar{x} \end{pmatrix}, \quad z = \begin{pmatrix} \bar{x} \\ \bar{x} \\ \vdots \\ \bar{x} \end{pmatrix}$$

## Continuation

►  $z = \frac{1}{3} \begin{pmatrix} x_1 + x_2 + x_3 \\ x_1 + x_2 + x_3 \\ x_1 + x_2 + x_3 \end{pmatrix}.$

► For general  $n$ , with  $\bar{x} = \frac{1}{n}(x_1 + x_2 + \cdots + x_n)$ ,

$$y = \begin{pmatrix} x_1 - \bar{x} \\ x_2 - \bar{x} \\ \vdots \\ x_n - \bar{x} \end{pmatrix}, \quad z = \begin{pmatrix} \bar{x} \\ \bar{x} \\ \vdots \\ \bar{x} \end{pmatrix}$$

► It is easy to see that  $y \in V_0$ ,  $z \in (V_0)^\perp$  and  $x = y + z$ .

# Projection as a linear map

- **Definition 13.2:** Let  $V_0$  be a subspace of a finite dimensional inner product space  $V$ . Then **the projection on to  $V_0$** , is the map

$$P : V \rightarrow V$$

defined by

$$P(x) = y$$

where  $x = y + z$ , with  $y \in V_0, z \in (V_0)^\perp$ .

# Projection as a linear map

- **Definition 13.2:** Let  $V_0$  be a subspace of a finite dimensional inner product space  $V$ . Then **the projection on to  $V_0$** , is the map

$$P : V \rightarrow V$$

defined by

$$P(x) = y$$

where  $x = y + z$ , with  $y \in V_0, z \in (V_0)^\perp$ .

- Note that since every  $x \in V$  decomposes uniquely as above,  $P$  is well-defined. If we want to emphasize the dependence of  $P$  on  $V_0$ , we may denote it by  $P_{V_0}$ .

# Projection as a linear map

- **Definition 13.2:** Let  $V_0$  be a subspace of a finite dimensional inner product space  $V$ . Then **the projection on to  $V_0$** , is the map

$$P : V \rightarrow V$$

defined by

$$P(x) = y$$

where  $x = y + z$ , with  $y \in V_0, z \in (V_0)^\perp$ .

- Note that since every  $x \in V$  decomposes uniquely as above,  $P$  is well-defined. If we want to emphasize the dependence of  $P$  on  $V_0$ , we may denote it by  $P_{V_0}$ .
- **Theorem 13.3:** Under the set up as above,



# Projection as a linear map

- **Definition 13.2:** Let  $V_0$  be a subspace of a finite dimensional inner product space  $V$ . Then **the projection on to  $V_0$** , is the map

$$P : V \rightarrow V$$

defined by

$$P(x) = y$$

where  $x = y + z$ , with  $y \in V_0, z \in (V_0)^\perp$ .

- Note that since every  $x \in V$  decomposes uniquely as above,  $P$  is well-defined. If we want to emphasize the dependence of  $P$  on  $V_0$ , we may denote it by  $P_{V_0}$ .
- **Theorem 13.3:** Under the set up as above,
- (i)  $P$  is a linear map. (ii)  $Px = x$  if and only if  $x \in V_0$  and  $Px = 0$  if and only if  $x \in (V_0)^\perp$ .

# Projection as a linear map

- **Definition 13.2:** Let  $V_0$  be a subspace of a finite dimensional inner product space  $V$ . Then **the projection on to  $V_0$** , is the map

$$P : V \rightarrow V$$

defined by

$$P(x) = y$$

where  $x = y + z$ , with  $y \in V_0, z \in (V_0)^\perp$ .

- Note that since every  $x \in V$  decomposes uniquely as above,  $P$  is well-defined. If we want to emphasize the dependence of  $P$  on  $V_0$ , we may denote it by  $P_{V_0}$ .
- **Theorem 13.3:** Under the set up as above,
- (i)  $P$  is a linear map. (ii)  $Px = x$  if and only if  $x \in V_0$  and  $Px = 0$  if and only if  $x \in (V_0)^\perp$ .
- (iii)  $P(V) = V_0$ .

# Projection as a linear map

- ▶ **Definition 13.2:** Let  $V_0$  be a subspace of a finite dimensional inner product space  $V$ . Then **the projection on to  $V_0$** , is the map

$$P : V \rightarrow V$$

defined by

$$P(x) = y$$

where  $x = y + z$ , with  $y \in V_0, z \in (V_0)^\perp$ .

- ▶ Note that since every  $x \in V$  decomposes uniquely as above,  $P$  is well-defined. If we want to emphasize the dependence of  $P$  on  $V_0$ , we may denote it by  $P_{V_0}$ .
- ▶ **Theorem 13.3:** Under the set up as above,
- ▶ (i)  $P$  is a linear map. (ii)  $Px = x$  if and only if  $x \in V_0$  and  $Px = 0$  if and only if  $x \in (V_0)^\perp$ .
- ▶ (iii)  $P(V) = V_0$ .
- ▶ (iv)  $P = P^2 = P^*$ .

# Projection as a linear map

- ▶ **Definition 13.2:** Let  $V_0$  be a subspace of a finite dimensional inner product space  $V$ . Then **the projection on to  $V_0$** , is the map

$$P : V \rightarrow V$$

defined by

$$P(x) = y$$

where  $x = y + z$ , with  $y \in V_0, z \in (V_0)^\perp$ .

- ▶ Note that since every  $x \in V$  decomposes uniquely as above,  $P$  is well-defined. If we want to emphasize the dependence of  $P$  on  $V_0$ , we may denote it by  $P_{V_0}$ .
- ▶ **Theorem 13.3:** Under the set up as above,
- ▶ (i)  $P$  is a linear map. (ii)  $Px = x$  if and only if  $x \in V_0$  and  $Px = 0$  if and only if  $x \in (V_0)^\perp$ .
- ▶ (iii)  $P(V) = V_0$ .
- ▶ (iv)  $P = P^2 = P^*$ .
- ▶ (v)  $P_{V_1} = I - P$  where  $V_1 = (V_0)^\perp$ .

## Continuation

- ▶ **Proof.** If  $V_0 = \{0\}$  then  $P = 0$  and all the properties mentioned above are easy to see.

## Continuation

- ▶ **Proof.** If  $V_0 = \{0\}$  then  $P = 0$  and all the properties mentioned above are easy to see.
- ▶ So assume  $V_0 \neq \{0\}$ .

# Continuation

- ▶ **Proof.** If  $V_0 = \{0\}$  then  $P = 0$  and all the properties mentioned above are easy to see.
- ▶ So assume  $V_0 \neq \{0\}$ .
- ▶ (i). Let  $\{v_1, \dots, v_k\}$  be an orthonormal basis of  $V_0$ . Extend it to an orthonormal basis  $\{v_1, v_2, \dots, v_n\}$  of  $V$ .
- ▶ Then we know that

$$P(x) = \sum_{j=1}^k \langle v_j, x \rangle v_j.$$

(Note that  $P$  does not depend upon the choice of this basis!)

# Continuation

- ▶ **Proof.** If  $V_0 = \{0\}$  then  $P = 0$  and all the properties mentioned above are easy to see.
- ▶ So assume  $V_0 \neq \{0\}$ .
- ▶ (i). Let  $\{v_1, \dots, v_k\}$  be an orthonormal basis of  $V_0$ . Extend it to an orthonormal basis  $\{v_1, v_2, \dots, v_n\}$  of  $V$ .
- ▶ Then we know that

$$P(x) = \sum_{j=1}^k \langle v_j, x \rangle v_j.$$

(Note that  $P$  does not depend upon the choice of this basis!)

- ▶ Since the inner product is linear in the second variable,  $P$  is a linear map. This proves (i).



## Continuation

- ▶ (ii). We know that  $x = \sum_{j=1}^n \langle v_j, x \rangle v_j$ . Therefore  $Px = x$  implies

$$\sum_{j=k+1}^n \langle v_j, x \rangle v_j = 0.$$

Therefore  $x = \sum_{j=1}^k \langle v_j, x \rangle v_j$  and hence  $x \in V_0$ .

# Continuation

- ▶ (ii). We know that  $x = \sum_{j=1}^n \langle v_j, x \rangle v_j$ . Therefore  $Px = x$  implies

$$\sum_{j=k+1}^n \langle v_j, x \rangle v_j = 0.$$

Therefore  $x = \sum_{j=1}^k \langle v_j, x \rangle v_j$  and hence  $x \in V_0$ .

- ▶ The converse is easy to see from the definition of  $P$ .

# Continuation

- ▶ (ii). We know that  $x = \sum_{j=1}^n \langle v_j, x \rangle v_j$ . Therefore  $Px = x$  implies

$$\sum_{j=k+1}^n \langle v_j, x \rangle v_j = 0.$$

Therefore  $x = \sum_{j=1}^k \langle v_j, x \rangle v_j$  and hence  $x \in V_0$ .

- ▶ The converse is easy to see from the definition of  $P$ .
- ▶ Now if  $Px = 0$ , then  $\sum_{j=1}^k \langle v_j, x \rangle v_j = 0$  and hence  $x = \sum_{j=k+1}^n \langle v_j, x \rangle v_j$ , that is,  $x \in (V_0)^\perp$ .

# Continuation

- ▶ (ii). We know that  $x = \sum_{j=1}^n \langle v_j, x \rangle v_j$ . Therefore  $Px = x$  implies

$$\sum_{j=k+1}^n \langle v_j, x \rangle v_j = 0.$$

Therefore  $x = \sum_{j=1}^k \langle v_j, x \rangle v_j$  and hence  $x \in V_0$ .

- ▶ The converse is easy to see from the definition of  $P$ .
- ▶ Now if  $Px = 0$ , then  $\sum_{j=1}^k \langle v_j, x \rangle v_j = 0$  and hence  $x = \sum_{j=k+1}^n \langle v_j, x \rangle v_j$ , that is,  $x \in (V_0)^\perp$ .
- ▶ Conversely if  $x \in (V_0)^\perp$ , then  $x = \sum_{j=k+1}^n \langle v_j, x \rangle v_j$ , and consequently  $Px = 0$ .

## Continuation

- ▶ (ii). We know that  $x = \sum_{j=1}^n \langle v_j, x \rangle v_j$ . Therefore  $Px = x$  implies

$$\sum_{j=k+1}^n \langle v_j, x \rangle v_j = 0.$$

Therefore  $x = \sum_{j=1}^k \langle v_j, x \rangle v_j$  and hence  $x \in V_0$ .

- ▶ The converse is easy to see from the definition of  $P$ .
- ▶ Now if  $Px = 0$ , then  $\sum_{j=1}^k \langle v_j, x \rangle v_j = 0$  and hence  $x = \sum_{j=k+1}^n \langle v_j, x \rangle v_j$ , that is,  $x \in (V_0)^\perp$ .
- ▶ Conversely if  $x \in (V_0)^\perp$ , then  $x = \sum_{j=k+1}^n \langle v_j, x \rangle v_j$ , and consequently  $Px = 0$ .
- ▶ This proves (ii).

- ▶ (iii). We want to show  $P(V) = V_0$ .

## Continuation

- ▶ (iii). We want to show  $P(V) = V_0$ .
- ▶ From the formula given for  $P$ ,  $Px \in V_0$  for every  $x \in V$  and hence  $P(V) \subseteq V_0$ . Since  $Px = x$  for every  $x \in V_0$ , the range of  $P$  includes whole of  $V_0$ . This proves (iii).

## Continuation

- ▶ (iii). We want to show  $P(V) = V_0$ .
- ▶ From the formula given for  $P$ ,  $Px \in V_0$  for every  $x \in V$  and hence  $P(V) \subseteq V_0$ . Since  $Px = x$  for every  $x \in V_0$ , the range of  $P$  includes whole of  $V_0$ . This proves (iii).



## Continuation

- ▶ (iii). We want to show  $P(V) = V_0$ .
- ▶ From the formula given for  $P$ ,  $Px \in V_0$  for every  $x \in V$  and hence  $P(V) \subseteq V_0$ . Since  $Px = x$  for every  $x \in V_0$ , the range of  $P$  includes whole of  $V_0$ . This proves (iii).
- ▶ (iv). If  $x = \sum_{j=1}^n c_j v_j$ , then  $Px = \sum_{j=1}^k c_j v_j$ .

## Continuation

- ▶ (iii). We want to show  $P(V) = V_0$ .
- ▶ From the formula given for  $P$ ,  $Px \in V_0$  for every  $x \in V$  and hence  $P(V) \subseteq V_0$ . Since  $Px = x$  for every  $x \in V_0$ , the range of  $P$  includes whole of  $V_0$ . This proves (iii).
- ▶ (iv). If  $x = \sum_{j=1}^n c_j v_j$ , then  $Px = \sum_{j=1}^k c_j v_j$ .
- ▶ Now  $P(P(x)) = P(\sum_{j=1}^k c_j v_j) = \sum_{j=1}^k c_j v_j = Px$ .

## Continuation

- ▶ (iii). We want to show  $P(V) = V_0$ .
- ▶ From the formula given for  $P$ ,  $Px \in V_0$  for every  $x \in V$  and hence  $P(V) \subseteq V_0$ . Since  $Px = x$  for every  $x \in V_0$ , the range of  $P$  includes whole of  $V_0$ . This proves (iii).
- ▶ (iv). If  $x = \sum_{j=1}^n c_j v_j$ , then  $Px = \sum_{j=1}^k c_j v_j$ .
- ▶ Now  $P(P(x)) = P(\sum_{j=1}^k c_j v_j) = \sum_{j=1}^k c_j v_j = Px$ .
- ▶ Hence  $P^2(x) = P(x)$  for every  $x$ , or  $P^2 = P$ .

## Continuation

- ▶ Suppose  $x_1, x_2$  are in  $V$ . Let  $x_1 = y_1 + z_1$  and  $x_2 = y_2 + z_2$  be the unique decompositions of  $x_1, x_2$  so that

$$y_1, y_2 \in V_0; \quad z_1, z_2 \in V_0^\perp.$$

## Continuation

- ▶ Suppose  $x_1, x_2$  are in  $V$ . Let  $x_1 = y_1 + z_1$  and  $x_2 = y_2 + z_2$  be the unique decompositions of  $x_1, x_2$  so that

$$y_1, y_2 \in V_0; \quad z_1, z_2 \in V_0^\perp.$$

- ▶ Note that  $\langle y_i, z_j \rangle = 0$  for all  $i, j$ .

## Continuation

- ▶ Suppose  $x_1, x_2$  are in  $V$ . Let  $x_1 = y_1 + z_1$  and  $x_2 = y_2 + z_2$  be the unique decompositions of  $x_1, x_2$  so that

$$y_1, y_2 \in V_0; \quad z_1, z_2 \in V_0^\perp.$$

- ▶ Note that  $\langle y_i, z_j \rangle = 0$  for all  $i, j$ .
- ▶ Now

$$\langle Px_1, x_2 \rangle = \langle y_1, y_2 + z_2 \rangle = \langle y_1, y_2 \rangle.$$

# Continuation

- ▶ Suppose  $x_1, x_2$  are in  $V$ . Let  $x_1 = y_1 + z_1$  and  $x_2 = y_2 + z_2$  be the unique decompositions of  $x_1, x_2$  so that

$$y_1, y_2 \in V_0; \quad z_1, z_2 \in V_0^\perp.$$

- ▶ Note that  $\langle y_i, z_j \rangle = 0$  for all  $i, j$ .

- ▶ Now

$$\langle Px_1, x_2 \rangle = \langle y_1, y_2 + z_2 \rangle = \langle y_1, y_2 \rangle.$$

- ▶ Similarly,

$$\langle x_1, Px_2 \rangle = \langle y_1 + z_1, y_2 \rangle = \langle y_1, y_2 \rangle.$$

# Continuation

- ▶ Suppose  $x_1, x_2$  are in  $V$ . Let  $x_1 = y_1 + z_1$  and  $x_2 = y_2 + z_2$  be the unique decompositions of  $x_1, x_2$  so that

$$y_1, y_2 \in V_0; \quad z_1, z_2 \in V_0^\perp.$$

- ▶ Note that  $\langle y_i, z_j \rangle = 0$  for all  $i, j$ .

- ▶ Now

$$\langle Px_1, x_2 \rangle = \langle y_1, y_2 + z_2 \rangle = \langle y_1, y_2 \rangle.$$

- ▶ Similarly,

$$\langle x_1, Px_2 \rangle = \langle y_1 + z_1, y_2 \rangle = \langle y_1, y_2 \rangle.$$

- ▶ Consequently,

$$\langle Px_1, x_2 \rangle = \langle x_1, Px_2 \rangle$$

for all  $x_1, x_2$  in  $V$ .



## Continuation

- ▶ Suppose  $x_1, x_2$  are in  $V$ . Let  $x_1 = y_1 + z_1$  and  $x_2 = y_2 + z_2$  be the unique decompositions of  $x_1, x_2$  so that

$$y_1, y_2 \in V_0; \quad z_1, z_2 \in V_0^\perp.$$

- ▶ Note that  $\langle y_i, z_j \rangle = 0$  for all  $i, j$ .

- ▶ Now

$$\langle Px_1, x_2 \rangle = \langle y_1, y_2 + z_2 \rangle = \langle y_1, y_2 \rangle.$$

- ▶ Similarly,

$$\langle x_1, Px_2 \rangle = \langle y_1 + z_1, y_2 \rangle = \langle y_1, y_2 \rangle.$$

- ▶ Consequently,

$$\langle Px_1, x_2 \rangle = \langle x_1, Px_2 \rangle$$

for all  $x_1, x_2$  in  $V$ .

- ▶ This shows that  $P^* = P$  from the defining property of the adjoint of  $P$ .

## Continuation

- (v). If  $x = \sum_{j=1}^n c_j v_j$ ,

$$P_{V_0}(x) = \sum_{j=1}^k c_j v_j, \quad P_{V_1}(x) = \sum_{j=k+1}^n c_j v_j.$$

## Continuation

- ▶ (v). If  $x = \sum_{j=1}^n c_j v_j$ ,

$$P_{V_0}(x) = \sum_{j=1}^k c_j v_j, \quad P_{V_1}(x) = \sum_{j=k+1}^n c_j v_j.$$

- ▶ From these formulae, it is easy to see that  $P_{V_1} = 1 - P_{V_0}$ .

## Continuation

- ▶ (v). If  $x = \sum_{j=1}^n c_j v_j$ ,

$$P_{V_0}(x) = \sum_{j=1}^k c_j v_j, \quad P_{V_1}(x) = \sum_{j=k+1}^n c_j v_j.$$

- ▶ From these formulae, it is easy to see that  $P_{V_1} = 1 - P_{V_0}$ .
- ▶ This completes the proof Theorem 13.2.

## Continuation

- ▶ (v). If  $x = \sum_{j=1}^n c_j v_j$ ,

$$P_{V_0}(x) = \sum_{j=1}^k c_j v_j, \quad P_{V_1}(x) = \sum_{j=k+1}^n c_j v_j.$$

- ▶ From these formulae, it is easy to see that  $P_{V_1} = 1 - P_{V_0}$ .
- ▶ This completes the proof Theorem 13.2.
- ▶ **Remark 13.4:** Observe that  $P_{\{0\}} = 0$  and  $P_V = I$ . In particular,

$$P_V(x) = x = \sum_{j=1}^n \langle v_j, x \rangle v_j$$

independent of the choice of the basis.

# Continuation

- ▶ (v). If  $x = \sum_{j=1}^n c_j v_j$ ,

$$P_{V_0}(x) = \sum_{j=1}^k c_j v_j, \quad P_{V_1}(x) = \sum_{j=k+1}^n c_j v_j.$$

- ▶ From these formulae, it is easy to see that  $P_{V_1} = 1 - P_{V_0}$ .
- ▶ This completes the proof Theorem 13.2.
- ▶ **Remark 13.4:** Observe that  $P_{\{0\}} = 0$  and  $P_V = I$ . In particular,

$$P_V(x) = x = \sum_{j=1}^n \langle v_j, x \rangle v_j$$

independent of the choice of the basis.

- ▶ We have just revisited our formula for the expansion of  $x$  in terms of an orthonormal basis.

# Distance between sets

- **Notation:** Let  $A, B$  be non-empty subsets of an inner product space  $V$  and let  $a \in V$ . Then

$$d(A, B) := \inf\{d(a, b) : a \in A, b \in B\}$$

and

$$d(a, B) := \inf\{d(a, b) : b \in B\}.$$

# Distance between sets

- **Notation:** Let  $A, B$  be non-empty subsets of an inner product space  $V$  and let  $a \in V$ . Then

$$d(A, B) := \inf\{d(a, b) : a \in A, b \in B\}$$

and

$$d(a, B) := \inf\{d(a, b) : b \in B\}.$$

- We may informally call  $d(A, B)$  as the distance between  $A, B$  and  $d(a, B)$  as the distance between  $a$  and  $B$ . But note that now we may have  $d(A, B) = 0$  without having  $A = B$ .



# Distance between sets

- ▶ **Notation:** Let  $A, B$  be non-empty subsets of an inner product space  $V$  and let  $a \in V$ . Then

$$d(A, B) := \inf\{d(a, b) : a \in A, b \in B\}$$

and

$$d(a, B) := \inf\{d(a, b) : b \in B\}.$$

- ▶ We may informally call  $d(A, B)$  as the distance between  $A, B$  and  $d(a, B)$  as the distance between  $a$  and  $B$ . But note that now we may have  $d(A, B) = 0$  without having  $A = B$ .
- ▶ In general,  $d(a, B)$  may not be attained at any point in  $B$ . Also when it is attained it may not be at some unique point in  $B$ .

# Distance between sets

- **Notation:** Let  $A, B$  be non-empty subsets of an inner product space  $V$  and let  $a \in V$ . Then

$$d(A, B) := \inf\{d(a, b) : a \in A, b \in B\}$$

and

$$d(a, B) := \inf\{d(a, b) : b \in B\}.$$

- We may informally call  $d(A, B)$  as the distance between  $A, B$  and  $d(a, B)$  as the distance between  $a$  and  $B$ . But note that now we may have  $d(A, B) = 0$  without having  $A = B$ .
- In general,  $d(a, B)$  may not be attained at any point in  $B$ . Also when it is attained it may not be at some unique point in  $B$ .
- **Example 14.1:** Take  $V = \mathbb{R}^2$ . Take  $a = (1, 0)$ . Consider  $B_1 = \{(x_1, x_2) : x_1 < 0\}$  and  $B_2 = \{(x_1, x_2) : |x_1 - 1| \geq 1\}$

# Distance between sets

- ▶ **Notation:** Let  $A, B$  be non-empty subsets of an inner product space  $V$  and let  $a \in V$ . Then

$$d(A, B) := \inf\{d(a, b) : a \in A, b \in B\}$$

and

$$d(a, B) := \inf\{d(a, b) : b \in B\}.$$

- ▶ We may informally call  $d(A, B)$  as the distance between  $A, B$  and  $d(a, B)$  as the distance between  $a$  and  $B$ . But note that now we may have  $d(A, B) = 0$  without having  $A = B$ .
- ▶ In general,  $d(a, B)$  may not be attained at any point in  $B$ . Also when it is attained it may not be at some unique point in  $B$ .
- ▶ **Example 14.1:** Take  $V = \mathbb{R}^2$ . Take  $a = (1, 0)$ . Consider  $B_1 = \{(x_1, x_2) : x_1 < 0\}$  and  $B_2 = \{(x_1, x_2) : |x_1 - 1| \geq 1\}$
- ▶ Then  $d(a, B_1) = 1$  is not attained at any point.  $d(a, B_2) = 1$  gets attained at two points.

# Best approximation property

- **Theorem 14.2:** Let  $V_0$  be a subspace of an inner product space  $V$ . Let  $P$  be the projection onto  $V_0$ . Then for  $x \in V$ ,

$$d(x, V_0) = d(x, Px).$$

Moreover,  $Px$  is the unique point  $v$  such that  $d(x, v) = d(x, V_0)$ .

# Best approximation property

- ▶ **Theorem 14.2:** Let  $V_0$  be a subspace of an inner product space  $V$ . Let  $P$  be the projection onto  $V_0$ . Then for  $x \in V$ ,

$$d(x, V_0) = d(x, Px).$$

Moreover,  $Px$  is the unique point  $v$  such that  $d(x, v) = d(x, V_0)$ .

- ▶ This theorem tells us that  $Px$  is the unique 'best approximation' for  $x$  in  $V_0$ .

# Best approximation property

- ▶ **Theorem 14.2:** Let  $V_0$  be a subspace of an inner product space  $V$ . Let  $P$  be the projection onto  $V_0$ . Then for  $x \in V$ ,

$$d(x, V_0) = d(x, Px).$$

Moreover,  $Px$  is the unique point  $v$  such that  $d(x, v) = d(x, V_0)$ .

- ▶ This theorem tells us that  $Px$  is the unique 'best approximation' for  $x$  in  $V_0$ .
- ▶ **Proof:** Suppose  $x = y + z$ , is the unique decomposition of  $x$ , with  $y \in V_0, z \in V_0^\perp$ .

## Continuation

- ▶ We have  $Px = y$ . Now consider any  $v \in V_0$ . Due to orthogonality of  $y - v$  and  $z$ , we get

## Continuation

- ▶ We have  $Px = y$ . Now consider any  $v \in V_0$ . Due to orthogonality of  $y - v$  and  $z$ , we get



$$\begin{aligned}\|x - v\|^2 &= \|(y + z) - v\|^2 \\ &= \langle (y - v) + z, (y - v) + z \rangle \\ &= \|y - v\|^2 + \|z\|^2.\end{aligned}$$



## Continuation

- ▶ We have  $Px = y$ . Now consider any  $v \in V_0$ . Due to orthogonality of  $y - v$  and  $z$ , we get



$$\begin{aligned}\|x - v\|^2 &= \|(y + z) - v\|^2 \\ &= \langle (y - v) + z, (y - v) + z \rangle \\ &= \|(y - v)\|^2 + \|z\|^2.\end{aligned}$$

- ▶ Hence

$$\inf_{v \in V_0} \|x - v\|^2 = \|z\|^2$$

and the infimum is attained only at  $v = y$ .

## Continuation

- ▶ We have  $Px = y$ . Now consider any  $v \in V_0$ . Due to orthogonality of  $y - v$  and  $z$ , we get



$$\begin{aligned}\|x - v\|^2 &= \|(y + z) - v\|^2 \\ &= \langle (y - v) + z, (y - v) + z \rangle \\ &= \|(y - v)\|^2 + \|z\|^2.\end{aligned}$$

- ▶ Hence

$$\inf_{v \in V_0} \|x - v\|^2 = \|z\|^2$$

and the infimum is attained only at  $v = y$ .

- ▶ This proves the theorem.

# Continuation

- ▶ We have  $Px = y$ . Now consider any  $v \in V_0$ . Due to orthogonality of  $y - v$  and  $z$ , we get



$$\begin{aligned}\|x - v\|^2 &= \|(y + z) - v\|^2 \\ &= \langle (y - v) + z, (y - v) + z \rangle \\ &= \|(y - v)\|^2 + \|z\|^2.\end{aligned}$$

- ▶ Hence

$$\inf_{v \in V_0} \|x - v\|^2 = \|z\|^2$$

and the infimum is attained only at  $v = y$ .

- ▶ This proves the theorem.
- ▶ Note that we are using the 'Pythagoras theorem' of inner product spaces.

## Example

- ▶ Consider the Example 13.1, where  $V = \mathbb{R}^n$  and

## Example

- ▶ Consider the Example 13.1, where  $V = \mathbb{R}^n$  and
- ▶  $V_0 = \{x \in V : \sum_{j=1}^n x_j = 0\}$ .

## Example

- ▶ Consider the Example 13.1, where  $V = \mathbb{R}^n$  and
- ▶  $V_0 = \{x \in V : \sum_{j=1}^n x_j = 0\}$ .
- ▶

$$V_1 = (V_0)^\perp = \left\{ \begin{pmatrix} c \\ c \\ \vdots \\ c \end{pmatrix} : c \in \mathbb{R} \right\}.$$

## Example

- ▶ Consider the Example 13.1, where  $V = \mathbb{R}^n$  and
- ▶  $V_0 = \{x \in V : \sum_{j=1}^n x_j = 0\}$ .



$$V_1 = (V_0)^\perp = \left\{ \begin{pmatrix} c \\ c \\ \vdots \\ c \end{pmatrix} : c \in \mathbb{R} \right\}.$$

- ▶ Let  $P_1$  be the projection onto  $V_1$ .

## Example

- ▶ Consider the Example 13.1, where  $V = \mathbb{R}^n$  and
- ▶  $V_0 = \{x \in V : \sum_{j=1}^n x_j = 0\}$ .



$$V_1 = (V_0)^\perp = \left\{ \begin{pmatrix} c \\ c \\ \vdots \\ c \end{pmatrix} : c \in \mathbb{R} \right\}.$$

- ▶ Let  $P_1$  be the projection onto  $V_1$ .
- ▶ Then  $P_1 x = \frac{1}{n}(x_1 + \cdots + x_n) =: \bar{x}$ .



## Continuation

- ▶ So the best approximation for  $x = (x_1, \dots, x_n)$  among constant sequences is  $(\bar{x}, \dots, \bar{x})$ .

## Continuation

- ▶ So the best approximation for  $x = (x_1, \dots, x_n)$  among constant sequences is  $(\bar{x}, \dots, \bar{x})$ .
- ▶ In other words, we have proved the theorem

$$\inf_{c \in \mathbb{R}} \sum_{j=1}^n (x_j - c)^2 = \sum_{j=1}^n (x_j - \bar{x})^2.$$

## Continuation

- ▶ So the best approximation for  $x = (x_1, \dots, x_n)$  among constant sequences is  $(\bar{x}, \dots, \bar{x})$ .
- ▶ In other words, we have proved the theorem

$$\inf_{c \in \mathbb{R}} \sum_{j=1}^n (x_j - c)^2 = \sum_{j=1}^n (x_j - \bar{x})^2.$$

- ▶ This value is  $n$  times the variance of the tuple  $\{x_1, \dots, x_n\}$ . In other words,

$$\inf_{c \in \mathbb{R}} \frac{1}{n} \sum_{j=1}^n (x_j - c)^2 = \text{Var} \{x_1, \dots, x_n\},$$

and the infimum gets attained only at  $\bar{x}$ .

## Continuation

- ▶ So the best approximation for  $x = (x_1, \dots, x_n)$  among constant sequences is  $(\bar{x}, \dots, \bar{x})$ .
- ▶ In other words, we have proved the theorem

$$\inf_{c \in \mathbb{R}} \sum_{j=1}^n (x_j - c)^2 = \sum_{j=1}^n (x_j - \bar{x})^2.$$

- ▶ This value is  $n$  times the variance of the tuple  $\{x_1, \dots, x_n\}$ . In other words,

$$\inf_{c \in \mathbb{R}} \frac{1}{n} \sum_{j=1}^n (x_j - c)^2 = \text{Var} \{x_1, \dots, x_n\},$$

and the infimum gets attained only at  $\bar{x}$ .

- ▶ **Exercise:** Work out more examples.

# Example

► **Example 14.3:** Consider  $V = \mathbb{R}^2$ . Let

$$V_0 = \left\{ c \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} : c \in \mathbb{R} \right\} \text{ where } \theta \text{ is a fixed real number.}$$

Write down the matrix of the projection onto  $V_0$ .

# Example

- **Example 14.3:** Consider  $V = \mathbb{R}^2$ . Let  
 $V_0 = \{c \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} : c \in \mathbb{R}\}$  where  $\theta$  is a fixed real number.  
Write down the matrix of the projection onto  $V_0$ .

- **Solution.** We take

$$P = \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} \cdot \begin{pmatrix} \cos \theta & \sin \theta \end{pmatrix}.$$

# Example

- **Example 14.3:** Consider  $V = \mathbb{R}^2$ . Let  
 $V_0 = \{c \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} : c \in \mathbb{R}\}$  where  $\theta$  is a fixed real number.  
Write down the matrix of the projection onto  $V_0$ .

- **Solution.** We take

$$P = \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} \cdot \begin{pmatrix} \cos \theta & \sin \theta \end{pmatrix}.$$

- Then for any vector  $x \in \mathbb{R}^2$ ,

$$Px = \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} \cdot \begin{pmatrix} \cos \theta & \sin \theta \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = v \langle v, x \rangle = \langle v, x \rangle \cdot v,$$

## Example

- **Example 14.3:** Consider  $V = \mathbb{R}^2$ . Let  
 $V_0 = \{c \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} : c \in \mathbb{R}\}$  where  $\theta$  is a fixed real number.  
Write down the matrix of the projection onto  $V_0$ .

- **Solution.** We take

$$P = \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} \cdot \begin{pmatrix} \cos \theta & \sin \theta \end{pmatrix}.$$

- Then for any vector  $x \in \mathbb{R}^2$ ,

$$Px = \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} \cdot \begin{pmatrix} \cos \theta & \sin \theta \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = v \langle v, x \rangle = \langle v, x \rangle \cdot v,$$

- where

$$v = \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix}.$$



- ▶ Observe that  $\{v\}$  is an orthonormal basis for  $V_0$ .

## Continuation

- ▶ Observe that  $\{v\}$  is an orthonormal basis for  $V_0$ .
- ▶ Therefore  $P$  given as above, that is,

$$P = \begin{bmatrix} \cos^2 \theta & \sin \theta \cos \theta \\ \sin \theta \cos \theta & \sin^2 \theta \end{bmatrix},$$

is the projection on to  $V_0$ .

# Continuation

- ▶ Observe that  $\{v\}$  is an orthonormal basis for  $V_0$ .
- ▶ Therefore  $P$  given as above, that is,

$$P = \begin{bmatrix} \cos^2 \theta & \sin \theta \cos \theta \\ \sin \theta \cos \theta & \sin^2 \theta \end{bmatrix},$$

is the projection on to  $V_0$ .

- ▶ You may verify  $P = P^2 = P^*$  and  $P(\mathbb{R}^2) = V_0$ .

# Algebraic description of projections

- **Theorem 15.1:** Let  $V$  be a finite dimensional inner product space. Then a linear map  $P : V \rightarrow V$  is a projection if and only if

$$P = P^2 = P^*.$$

# Algebraic description of projections

- ▶ **Theorem 15.1:** Let  $V$  be a finite dimensional inner product space. Then a linear map  $P : V \rightarrow V$  is a projection if and only if

$$P = P^2 = P^*.$$

- ▶ **Proof:** We have already seen that if  $P$  is a projection then  $P = P^2 = P^*$ .

# Algebraic description of projections

- ▶ **Theorem 15.1:** Let  $V$  be a finite dimensional inner product space. Then a linear map  $P : V \rightarrow V$  is a projection if and only if

$$P = P^2 = P^*.$$

- ▶ **Proof:** We have already seen that if  $P$  is a projection then  $P = P^2 = P^*$ .
- ▶ Now suppose  $P : V \rightarrow V$  is a linear map such that

$$P = P^2 = P^*.$$

# Algebraic description of projections

- ▶ **Theorem 15.1:** Let  $V$  be a finite dimensional inner product space. Then a linear map  $P : V \rightarrow V$  is a projection if and only if

$$P = P^2 = P^*.$$

- ▶ **Proof:** We have already seen that if  $P$  is a projection then  $P = P^2 = P^*$ .
- ▶ Now suppose  $P : V \rightarrow V$  is a linear map such that

$$P = P^2 = P^*.$$

- ▶ Take  $V_0 = P(V) = \{Px : x \in V\}$ .

# Algebraic description of projections

- ▶ **Theorem 15.1:** Let  $V$  be a finite dimensional inner product space. Then a linear map  $P : V \rightarrow V$  is a projection if and only if

$$P = P^2 = P^*.$$

- ▶ **Proof:** We have already seen that if  $P$  is a projection then  $P = P^2 = P^*$ .
- ▶ Now suppose  $P : V \rightarrow V$  is a linear map such that

$$P = P^2 = P^*.$$

- ▶ Take  $V_0 = P(V) = \{Px : x \in V\}$ .
- ▶ Clearly  $V_0$  is a subspace of  $V$ .



- ▶ We wish to show that  $P$  is the projection onto  $V_0$ .

## Continuation

- ▶ We wish to show that  $P$  is the projection onto  $V_0$ .
- ▶ For  $x \in V$ , take  $y = Px$  and  $z = x - Px$ .

## Continuation

- ▶ We wish to show that  $P$  is the projection onto  $V_0$ .
- ▶ For  $x \in V$ , take  $y = Px$  and  $z = x - Px$ .
- ▶ Clearly  $x = y + z$  and  $y \in V_0$ .

# Continuation

- ▶ We wish to show that  $P$  is the projection onto  $V_0$ .
- ▶ For  $x \in V$ , take  $y = Px$  and  $z = x - Px$ .
- ▶ Clearly  $x = y + z$  and  $y \in V_0$ .
- ▶ It suffices to show that  $z \in V_0^\perp$ .

## Continuation

- ▶ Consider any  $w \in V_0$ . So  $w = Pv$  for some  $v \in V$ .

# Continuation

- ▶ Consider any  $w \in V_0$ . So  $w = Pv$  for some  $v \in V$ .
- ▶ Now

$$\begin{aligned}\langle w, z \rangle &= \langle Pv, x - Px \rangle \\ &= \langle v, P^*(x - Px) \rangle \\ &= \langle v, P(x - Px) \rangle \\ &= \langle v, Px - P^2x \rangle \\ &= \langle v, Px - Px \rangle \\ &= 0.\end{aligned}$$

# Continuation

- ▶ Consider any  $w \in V_0$ . So  $w = Pv$  for some  $v \in V$ .
- ▶ Now

$$\begin{aligned}\langle w, z \rangle &= \langle Pv, x - Px \rangle \\ &= \langle v, P^*(x - Px) \rangle \\ &= \langle v, P(x - Px) \rangle \\ &= \langle v, Px - P^2x \rangle \\ &= \langle v, Px - Px \rangle \\ &= 0.\end{aligned}$$

- ▶ This shows that  $z \in V_0^\perp$ . ■

# Diagonalization of a projection

- **Theorem 15.2:** Let  $V_0$  be a non-zero finite dimensional subspace of a finite dimensional inner product space  $V$  and let  $P$  the projection onto  $V_0$ . Then there exists an orthonormal basis  $\mathcal{B}$  such that on  $\mathcal{B}$ , the matrix of  $P$  is given by

$$A = \begin{bmatrix} I & 0 \\ 0 & 0 \end{bmatrix}$$

where  $I$  is of order  $k \times k$ , with  $k = \dim(V_0)$ .



# Diagonalization of a projection

- ▶ **Theorem 15.2:** Let  $V_0$  be a non-zero finite dimensional subspace of a finite dimensional inner product space  $V$  and let  $P$  the projection onto  $V_0$ . Then there exists an orthonormal basis  $\mathcal{B}$  such that on  $\mathcal{B}$ , the matrix of  $P$  is given by

$$A = \begin{bmatrix} I & 0 \\ 0 & 0 \end{bmatrix}$$

where  $I$  is of order  $k \times k$ , with  $k = \dim(V_0)$ .

- ▶ **Proof:** Let  $\{v_1, \dots, v_k, v_{k+1}, \dots, v_n\}$  be an orthonormal basis for  $V$ , where  $\{v_1, \dots, v_k\}$  is an orthonormal basis for  $V_0$ .

# Diagonalization of a projection

- ▶ **Theorem 15.2:** Let  $V_0$  be a non-zero finite dimensional subspace of a finite dimensional inner product space  $V$  and let  $P$  the projection onto  $V_0$ . Then there exists an orthonormal basis  $\mathcal{B}$  such that on  $\mathcal{B}$ , the matrix of  $P$  is given by

$$A = \begin{bmatrix} I & 0 \\ 0 & 0 \end{bmatrix}$$

where  $I$  is of order  $k \times k$ , with  $k = \dim(V_0)$ .

- ▶ **Proof:** Let  $\{v_1, \dots, v_k, v_{k+1}, \dots, v_n\}$  be an orthonormal basis for  $V$ , where  $\{v_1, \dots, v_k\}$  is an orthonormal basis for  $V_0$ .
- ▶ Take  $\mathcal{B} = \{v_1, \dots, v_n\}$ .

# Diagonalization of a projection

- ▶ **Theorem 15.2:** Let  $V_0$  be a non-zero finite dimensional subspace of a finite dimensional inner product space  $V$  and let  $P$  the projection onto  $V_0$ . Then there exists an orthonormal basis  $\mathcal{B}$  such that on  $\mathcal{B}$ , the matrix of  $P$  is given by

$$A = \begin{bmatrix} I & 0 \\ 0 & 0 \end{bmatrix}$$

where  $I$  is of order  $k \times k$ , with  $k = \dim(V_0)$ .

- ▶ **Proof:** Let  $\{v_1, \dots, v_k, v_{k+1}, \dots, v_n\}$  be an orthonormal basis for  $V$ , where  $\{v_1, \dots, v_k\}$  is an orthonormal basis for  $V_0$ .
- ▶ Take  $\mathcal{B} = \{v_1, \dots, v_n\}$ .
- ▶ We have  $Pv_j = v_j$  for  $1 \leq j \leq k$  and  $Pv_j = 0$  for  $(k+1) \leq j \leq n$ .

# Eigenvalues of projections

- **Theorem 15.3:** Let  $V$  be a finite dimensional inner product space and let  $P : V \rightarrow V$  be a projection. Suppose  $\lambda$  is an eigenvalue of  $P$  then  $\lambda \in \{0, 1\}$ .

# Eigenvalues of projections

- ▶ **Theorem 15.3:** Let  $V$  be a finite dimensional inner product space and let  $P : V \rightarrow V$  be a projection. Suppose  $\lambda$  is an eigenvalue of  $P$  then  $\lambda \in \{0, 1\}$ .
- ▶ **Proof:** This is clear as the characteristic polynomial of the matrix

$$A = \begin{bmatrix} I & 0 \\ 0 & 0 \end{bmatrix}$$

where  $I$  is of order  $k \times k$ , is given by  $p(x) = (x - 1)^k x^{n-k}$ . ■

# Eigenvalues of projections

- ▶ **Theorem 15.3:** Let  $V$  be a finite dimensional inner product space and let  $P : V \rightarrow V$  be a projection. Suppose  $\lambda$  is an eigenvalue of  $P$  then  $\lambda \in \{0, 1\}$ .
- ▶ **Proof:** This is clear as the characteristic polynomial of the matrix

$$A = \begin{bmatrix} I & 0 \\ 0 & 0 \end{bmatrix}$$

where  $I$  is of order  $k \times k$ , is given by  $p(x) = (x - 1)^k x^{n-k}$ . ■

- ▶ Note that 0 is the only eigenvalue of a projection  $P$  iff  $P = 0$ . Similarly 1 is the only eigenvalue of  $P$  if and only if  $P = I$ .

► Consequently

$$\langle v_i, P v_j \rangle = \begin{cases} \delta_{ij} & \text{if } 1 \leq i, j \leq k \\ 0 & \text{otherwise.} \end{cases}$$

- ▶ Consequently

$$\langle v_i, P v_j \rangle = \begin{cases} \delta_{ij} & \text{if } 1 \leq i, j \leq k \\ 0 & \text{otherwise.} \end{cases}$$

- ▶ Hence the matrix of  $P$  on the orthonormal basis  $\mathcal{B}$  is  $A$  as above.



# Rank and trace

- **Corollary 15.4:** Let  $R$  be the matrix of a projection map on some orthonormal basis. Then  $\text{trace}(R) = \text{rank}(R)$ .

# Rank and trace

- ▶ **Corollary 15.4:** Let  $R$  be the matrix of a projection map on some orthonormal basis. Then  $\text{trace}(R) = \text{rank}(R)$ .
- ▶ **Proof:** Clear from the description of the matrix of  $P$ .

# Rank and trace

- ▶ **Corollary 15.4:** Let  $R$  be the matrix of a projection map on some orthonormal basis. Then  $\text{trace}(R) = \text{rank}(R)$ .
- ▶ **Proof:** Clear from the description of the matrix of  $P$ .
- ▶ **Example 15.5:** Show that  $R : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  defined by

$$R \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} \frac{x_1+x_2+x_3}{3} \\ \frac{x_1+x_2+x_3}{3} \\ \frac{x_1+x_2+x_3}{3} \end{pmatrix}$$

is a projection. Write down the matrix of  $R$  on standard basis. Find a suitable orthonormal basis such that the entries of the matrix of  $R$  on that basis are 0's and 1's.

# Rank and trace

- ▶ **Corollary 15.4:** Let  $R$  be the matrix of a projection map on some orthonormal basis. Then  $\text{trace}(R) = \text{rank}(R)$ .
- ▶ **Proof:** Clear from the description of the matrix of  $P$ .
- ▶ **Example 15.5:** Show that  $R : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  defined by

$$R \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} \frac{x_1+x_2+x_3}{3} \\ \frac{x_1+x_2+x_3}{3} \\ \frac{x_1+x_2+x_3}{3} \end{pmatrix}$$

is a projection. Write down the matrix of  $R$  on standard basis. Find a suitable orthonormal basis such that the entries of the matrix of  $R$  on that basis are 0's and 1's.

# Rank and trace

- ▶ **Corollary 15.4:** Let  $R$  be the matrix of a projection map on some orthonormal basis. Then  $\text{trace}(R) = \text{rank}(R)$ .
- ▶ **Proof:** Clear from the description of the matrix of  $P$ .
- ▶ **Example 15.5:** Show that  $R : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  defined by

$$R \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} \frac{x_1+x_2+x_3}{3} \\ \frac{x_1+x_2+x_3}{3} \\ \frac{x_1+x_2+x_3}{3} \end{pmatrix}$$

is a projection. Write down the matrix of  $R$  on standard basis. Find a suitable orthonormal basis such that the entries of the matrix of  $R$  on that basis are 0's and 1's.

- ▶ **END OF LECTURE 15.**