

$$[x_n \rightarrow x : |x_n - x| < \frac{1}{n} \forall n \geq N]$$

5

Jaydeb Sarkar

Def: For  $S \subseteq \mathbb{R}$ ,  $\mathcal{B}(S) := \{f: S \rightarrow \mathbb{R} \text{ bdd.}\}$

Set of all bdd fns on  $S$ .

A vector space over  $\mathbb{R}$ .

$$[\text{Here } (\alpha f + g)(x) = \alpha f(x) + g(x), \forall \alpha \in \mathbb{R}, f, g \in \mathcal{B}(S)]$$

Def:  $\forall f \in \mathcal{B}(S)$ , define  $\|f\|$  (read: norm of  $f$ )

$$\|f\| = \sup_{x \in S} |f(x)|$$

Also known as "the sup norm".

$f(x) = x$   
 $g(x) = 1 - x$   
 $\|f\| = 1 = \|g\|$   
 $\|f + g\| = \frac{1}{4}$   
 $S = [0, 1]$

Remark: (1)  $\|\cdot\| : \mathcal{B}(S) \rightarrow \mathbb{R}_{\geq 0}$

$$(2) \|f\| = 0 \Leftrightarrow f \equiv 0$$

$$(3) \|f + g\| \leq \|f\| + \|g\|$$

$$(4) \|\alpha f\| = |\alpha| \|f\| \quad \forall \alpha \in \mathbb{R}$$

$$(5) \|fg\| \leq \|f\| \|g\|$$

Submultiplicative.

$$\Delta\text{-inequality.}$$

$$\sup_S \left( \frac{|f(x)|}{1} \left| \frac{g(x)}{1} \right| \right) = \sup_S |f| \times \sup_S |g|$$

Looks like  $|\cdot|$  on  $\mathbb{R}$ !!

Remark: Indeed,  $\|\cdot\|$  on  $\mathcal{B}(S)$  plays the role of  $|\cdot|$  on  $\mathbb{R}$ !!

Def:  $\forall f, g \in \mathcal{B}(S)$ , define the distance between  $f$  &  $g$

as:

$$d(f, g) = \|f - g\|$$

Also known as metric on  $\mathcal{B}(S)$ .

Remark:  $\therefore d: \mathcal{B}(S) \times \mathcal{B}(S) \rightarrow \mathbb{R}_{\geq 0}$ . And:

$$(1) d(f, g) = \sup_{x \in S} |f(x) - g(x)|$$

$$(2) d(f, g) \leq d(f, h) + d(h, g)$$

$$(3) d(f, g) = 0 \Leftrightarrow f = g$$

$\forall f, g \in \mathcal{B}(S)$

(6)

$= \{g: S \rightarrow \mathbb{R}\}$

# Let  $\{f_n\} \subseteq \mathcal{F}(S)$  &  $f \in \mathcal{F}(S)$ . Recall:  $f_n \xrightarrow{u} f$  <sup>on S</sup> if  $\forall \varepsilon > 0 \exists N \in \mathbb{N}$  s.t.

$$|f_n(x) - f(x)| < \varepsilon \quad \forall x \in S, n \geq N.$$

$$\Leftrightarrow (\because f_n - f \in \mathcal{B}(S) \quad \forall n \geq N)$$

$$\|f_n - f\| < \varepsilon \quad \forall n \geq N.$$

$N \geq n$   
 $|x_n - x| < \varepsilon$



Now this "looks" like modulus.

Ensures that  $f_n - f \in \mathcal{B}(S) \quad \forall n \geq N$ .

eg:

Let  $S = [-1, 1]$ . Define  $f_n \in \mathcal{F}(S)$  by

$$f_n(x) = \begin{cases} \frac{1}{n} & \text{if } |x| \leq \frac{1}{n} \\ |x| & \text{if } \frac{1}{n} < |x| \leq 1 \end{cases}$$



Then  $\lim_{n \rightarrow \infty} f_n(x) = |x| \quad \forall x \in S.$

i.e.  $f_n \xrightarrow{p} f$  on  $S$ , where

$$f(x) = |x| \quad \forall x \in S.$$

\* fixed  $x \in [-1, 1]$

$$|f_n(x) - f(x)| = \begin{cases} \left| \frac{1}{n} - |x| \right| & \text{if } |x| \leq \frac{1}{n} \\ 0 & \text{if } \frac{1}{n} < |x| \leq 1 \end{cases} \Rightarrow f_n(x) \rightarrow f(x) \quad \forall x.$$

In fact:  $\|f_n - f\| = \sup_{x \in [-1, 1]} |f_n(x) - f(x)|$

$$= \frac{1}{n} \quad \forall n. \quad \leftarrow \because \text{the max occurs at } x=0.$$

$\therefore$  For  $\varepsilon > 0 \exists N \in \mathbb{N}$  s.t.

$$\|f_n - f\| < \varepsilon \quad \forall n \geq N.$$

$$\Rightarrow f_n \xrightarrow{u} f \text{ on } [-1, 1].$$

□

Remark:  $f_n \xrightarrow{u} f \Rightarrow f_n \xrightarrow{p} f$ . (i.e. unif  $\Rightarrow$  point.)

Proof: Suppose that  $f_n \xrightarrow{u} f$  uniformly on  $S$ .

Let  $\varepsilon > 0$ . Then  $\exists N \in \mathbb{N}$  s.t.

$$\|f_n - f\| < \varepsilon \quad \forall n \geq N.$$

$$\text{i.e. } \sup_x |f_n(x) - f(x)| < \varepsilon \quad \forall n \geq N.$$

$$\text{i.e. } |f_n(x) - f(x)| < \varepsilon \quad \forall n \geq N, x \in S.$$

$$\Rightarrow f_n(x) \rightarrow f(x) \quad \forall x \in S.$$

$$\therefore f_n \rightarrow f \text{ pointwise.} \quad \square$$

#  $\therefore$  For u.c. it is perhaps a good idea to compute the pointwise limit (if exists) first !!

[If pointwise limit fails to exist, then u.c. also fails.]

Remark: " $\Leftarrow$ " of the above is NOT true! i.e. pointwise  $\not\Rightarrow$  unif.

eg:  $f_n(x) = x^n$  on  $[0, 1]$ .  $\leftarrow \{f_n\}$  is NOT u.c. (HW).

# Now, we develop some useful tools for convergence !!

The above examples suggest a pattern!

Although computation of pointwise limit may be diff. occasionally.

(8)

Thm: (Cauchy's criterion): Let  $\{f_n\} \subseteq \mathcal{F}(S)$ .

~~$$\lim_{n \rightarrow \infty} f_n(x) = f(x) \quad \forall x \in S.$$~~

~~$$[i.e. f_n \xrightarrow{p} f].$$~~

Then  $\{f_n\}$  is u.c.

(This is a must for u.c.)

Then  $f_n \xrightarrow{u} f \iff \forall \varepsilon > 0 \exists N \in \mathbb{N} \text{ s.t.}$

$$\|f_m - f_n\| < \varepsilon \quad \forall m, n \geq N.$$

No need to worry about limit  $f_n \rightarrow f$ .

(Also ensures that  $f_m - f_n \in \mathcal{B}(S) \quad \forall m, n$ ).

Proof: " $\Rightarrow$ " Let  $f_n \xrightarrow{u} f$ , and let  $\varepsilon > 0$ .

$$\therefore \exists N \in \mathbb{N} \text{ s.t.} \quad \|f_n - f\| < \varepsilon/2 \quad \forall n \geq N.$$

$\therefore \forall m, n \geq N$ , we have:

$$\|f_n - f_m\| = \|(f_n - f) + (f - f_m)\|$$

$$\begin{aligned} &\leq \|f_n - f\| + \|f - f_m\| \\ &< \varepsilon/2 + \varepsilon/2 = \varepsilon. \end{aligned}$$

$\Delta$ -ineq.

Standard proof like for  $\{x_n\} \subseteq \mathbb{R}$

" $\Leftarrow$ " Let  $\varepsilon > 0$ .

$$\therefore \exists N \in \mathbb{N} \text{ s.t.}$$

$$\|f_n - f_m\| < \varepsilon/2 \quad \forall m, n \geq N.$$

$$= \sup_{x \in S} |f_n(x) - f_m(x)|$$

i.e.  ~~$\{f_n(x)\}_{n \geq 1}$  is Cauchy  $\forall x \in S$ .~~

$$i.e. \quad |f_n(x) - f_m(x)| < \varepsilon/2 \quad \forall m, n \geq N \quad \forall x \in S. \quad (*)$$

$\Rightarrow \{f_n(x)\}$  is Cauchy  $\forall x \in S. \Rightarrow \lim_{n \rightarrow \infty} f_n(x) := f(x)$  exists  $\forall x \in S$ .

~~$$i.e. f_n \xrightarrow{p} f.$$~~

$$\text{Also } (*) \Rightarrow f_n(x) - \varepsilon/2 < f_m(x) < f_n(x) + \varepsilon/2 \quad \forall m, n \geq N, x \in S.$$



But  $\lim_{m \rightarrow \infty} f_m(x) = f(x) \quad \forall x \in S.$

⑨  
Jaydeb Sarkar  
 $f(x) := \lim_{n \rightarrow \infty} f_n(x).$

$\therefore$  Taking limit as  $m \rightarrow \infty$ , we have.

$$f_n(x) - \varepsilon/2 \leq f(x) \leq f_n(x) + \varepsilon/2 \quad \forall n \geq N \quad \forall x \in S.$$

$$\Rightarrow |f_n(x) - f(x)| \leq \varepsilon/2 < \varepsilon$$

$$\forall n \geq N \quad \forall x \in S.$$

$$\Rightarrow \|f_n - f\| < \varepsilon \quad \forall n \geq N.$$

i.e.  $f_n \xrightarrow{u} f$  on  $S.$

$= \{f_n : n \in \mathbb{N}\}.$

$f : \mathbb{N} \rightarrow \mathbb{Q} \cap [0, 1]$

# Cauchy Criterion is often useful.

Eg: Let  $\{r_n\}_{n=1}^{\infty}$  be an enumeration of rationals  $\mathbb{Q} \cap [0, 1].$

For each  $n \in \mathbb{N}$ , define  $f_n : [0, 1] \rightarrow \mathbb{R}$  by

$$f_n(x) = \begin{cases} 0 & \text{if } x = r_1, \dots, r_n \\ 1 & \text{if } x \neq r_1, \dots, r_n. \end{cases}$$

$y=1$   
 $r_1, \dots, r_n$

Then  $\{f_n\}$  is not u.c. on  $[0, 1].$

Indeed, choose  $\varepsilon = \frac{1}{2}.$

Observe, for each  $m \in \mathbb{N}$ ,  $f_m(r_m) = 0$  &  $f_m(r_{m+1}) = 1$

we have  $f_n(r_{n+1}) = 1$  &  $f_{n+1}(r_{n+1}) = 0.$

$$\Rightarrow |f_n(r_{n+1}) - f_{n+1}(r_{n+1})| = 1. \quad \forall n.$$

$$\therefore \nexists N \in \mathbb{N} \text{ s.t. } |f_m(x) - f_n(x)| < \varepsilon = \frac{1}{2}.$$

$$\forall m, n \geq N.$$

$\therefore$  By Cauchy Criterion,  $\{f_n\}$  is NOT u.c.

[Q:  $\{f_n\}$  Converges pointwise?]

Thm: (M-test): Let  $\{f_n\} \subseteq \mathcal{F}(S)$  & suppose  $f_n \xrightarrow{p} f$ .

Set  $M_n := \sup_{x \in S} |f_n(x) - f(x)|$ .

Then  $f_n \xrightarrow{u} f \text{ on } S \iff M_n \longrightarrow 0$ .

Proof: " $\Rightarrow$ " Let  $f_n \xrightarrow{u} f$ , & let  $\varepsilon > 0$ .

$\therefore \exists N \in \mathbb{N} \cdot \forall n \geq N, \|f_n - f\| < \varepsilon$

$\stackrel{\text{By def.}}{=} M_n$

i.e.  $M_n < \varepsilon \quad \forall n \geq N$ .

$\Rightarrow M_n \longrightarrow 0$ .

" $\Leftarrow$ " Let  $\varepsilon > 0$ . Then  $\exists N \in \mathbb{N} \cdot \forall n \geq N, M_n < \varepsilon$

$\Rightarrow \|f_n - f\| < \varepsilon \quad \forall n \geq N$ .

$\Rightarrow f_n \xrightarrow{u} f$ .

$\Rightarrow f_n \xrightarrow{u} f$ .

eg:

$\forall n \in \mathbb{N}, x \in [0, 1]$ , define

$$f_n(x) = \frac{nx}{1+n^2x^2}$$

Note that:  $\frac{nx}{1+n^2x^2} < 1 \quad \forall n$  & fixed  $x \in [0, 1]$ .

$\Rightarrow \lim_{n \rightarrow \infty} f_n(x) = f(x) \quad \text{i.e.} \quad f_n \xrightarrow{p} f$ ,

where  $f(x) \equiv 0$ .

$\therefore M_n = \|f_n - f\| = \sup_{x \in [0, 1]} \left[ \frac{nx}{1+n^2x^2} \right] \quad \forall n$ .

$\frac{nx}{1+n^2x^2} \rightarrow 0$  as  $n \rightarrow \infty$   
as  $x$  is fixed.

$f_n(x) < 1 \quad \forall n$   
 $\Rightarrow \lim_{n \rightarrow \infty} f_n(x) = 0$

fix  $x \in [0, 1]$ ,  $\lim_{n \rightarrow \infty} f(x) = 0$

But,  $x=0 \Rightarrow f_n(x) = 0 \quad \forall n$ .

$$\& x > 0 \Rightarrow \frac{\frac{1}{nx} + nx}{2} \geq \sqrt{\frac{1}{nx} \times nx} = 1.$$

$$\Rightarrow \frac{1}{2} \times \frac{1+n^2x^2}{nx} \geq 1.$$

$$\Rightarrow \frac{nx}{1+n^2x^2} \leq \frac{1}{2}.$$

And, "=" occurs at  $x = \frac{1}{n}$ .

$$\therefore M_n = \frac{1}{2} \quad \forall n \geq 1.$$

$$\Rightarrow M_n \not\rightarrow 0.$$

$\therefore$  By M-test,  $\{f_n\}$  is not unif. convergent on  $[0,1]$ .

□

eg:  $\forall n \in \mathbb{N}$ , define  $f_n(x) = x^n(1-x)$ .  $x \in [0,1]$ .

$$\left. \begin{aligned} f_n(x) &= \begin{cases} \therefore f_n(1) = 0 = f_n(0) \quad \forall n. \\ \& \text{for } 0 < x < 1, f_n(x) \rightarrow 0. \end{cases} \end{aligned} \right\}$$


$$\therefore f_n \xrightarrow{p} 0 \quad \text{the zero fn.}$$

$$\therefore M_n = \sup_{x \in [0,1]} [x^n(1-x)].$$

$$\text{But } f'_n(x) = x^{n-1} [n - (n+1)x].$$

$$\therefore \underline{f'_n(x) \geq 0} \quad \forall x \in [0, \frac{n}{n+1}]$$

$$\& \underline{f'_n(x) \leq 0} \quad \forall x \in [\frac{n}{n+1}, 1].$$



(12)

$\therefore f_n(0) = f_n(1) = 0$ , it follows that  $f_n$  has a max.

at  $x = \frac{n}{n+1}$ .

$$\text{Now } f_n\left(\frac{n}{n+1}\right) = \left(\frac{n}{n+1}\right)^n \times \frac{1}{n+1} < \frac{1}{n+1}.$$

$$\text{i.e. } M_n < \frac{1}{n+1}.$$

$$\Rightarrow M_n \rightarrow 0.$$

$\therefore$  By M-test,  $\{f_n\}$  is u.c. ( $\& f_n \xrightarrow{u} 0$ ).  $\square$

~~Eg~~ Consider  $S = \mathbb{R}$ .

Remark: Suppose  $|f_n(x) - f(x)| \leq M_n \quad \forall x \in S, \quad n \geq 1.$

If  $M_n \rightarrow 0$ , then  $f_n \xrightarrow{u} f$  on  $S$ .

all  $n$  but finitely many.

Proof: Follows from M-test.

Eg: Let  $r > 0$ .  $f_n(x) := e^{-nx} \quad \forall x \in [r, \infty).$

Now  $f_n \downarrow$  on  $\mathbb{R} \quad \forall n$ .

$$\Rightarrow 0 < f_n(x) = e^{-nx} \leq e^{-nr} \quad \forall n \geq 1, \& x \geq r.$$

But  $e^{-nr} \rightarrow 0$  as  $n \rightarrow \infty$ .

$\Rightarrow f_n \rightarrow 0$  uniformly on  $[r, \infty).$

$\square$