

Thm: (Limit Comparison test - I)



Let $f(x), g(x) \geq 0 \quad \forall x \in [a, b)$.

Suppose $\lim_{x \rightarrow b^-} \frac{f(x)}{g(x)} = l$.

If $l \neq 0, \infty$, then $\int_a^b f$ & $\int_a^b g$ Converge or diverge together at b.

Proof:

Suppose $l > 0$. Choose $\varepsilon > 0$ small s.t. $l - \varepsilon > 0$.

$\therefore \lim_{x \rightarrow b^-} \frac{f(x)}{g(x)} = l, \quad \exists c \in [a, b)$ s.t.

$$\left| \frac{f(x)}{g(x)} - l \right| < \varepsilon \quad \forall x \in (c, b).$$

$$\Rightarrow l - \varepsilon < \frac{f(x)}{g(x)} < l + \varepsilon \quad \forall x \in (c, b).$$

$$\Rightarrow (l - \varepsilon) g(x) < f(x) < (l + \varepsilon) g(x).$$

$\therefore l - \varepsilon > 0$ & $g(x) \geq 0 \quad \forall x$, it follows that

$$0 \leq (l - \varepsilon) g(x) \leq f(x) \quad \forall x \in (c, b).$$

\therefore By Comparison test, if $\int_a^b f$, or equivalently, if

$\int_c^b f$ Converges at b, then $(l - \varepsilon) \int_c^b g$, or equivalently,

$$(l - \varepsilon) \int_a^b g \text{ Converges at b.} \Rightarrow \int_a^b g \text{ Converges at b.}$$

Now suppose $\int_a^b g$, or equiv., $\int_c^b g$ converges at b .

Again by $f(x) < (1+\epsilon)g(x) \quad \forall x \in (c, b)$ &
by the comparison test, it follows that $\int_a^b f$ converges at b .

$\therefore \int_a^b f$ converges at $b \iff \int_a^b g$ converges at b .

Illy; $\int_a^b f$ diverges at $b \iff \int_a^b g$ diverges at b .



eg: (1) $\int_0^1 \frac{\sin x}{x^2} dx$. \leftarrow An I.I. at $x=0$.

Set $f(x) = \frac{\sin x}{x^2}$ & $g(x) = \frac{1}{x}$.

$$\therefore \frac{f(x)}{g(x)} = \frac{\sin x}{x}.$$

$$\text{Now } \lim_{x \rightarrow 0^+} \frac{f(x)}{g(x)} = \lim_{x \rightarrow 0^+} \frac{\sin x}{x} = 1 > 0.$$

But $\int_0^1 g = \int_0^1 \frac{1}{x} dx$ diverges \leftarrow why?

\therefore By Limit comparison test, $\int_0^1 \frac{\sin x}{x^2} dx$ diverges.

(2) $\int_0^1 \frac{e^{\sqrt{x}} - 1}{x} dx$. \leftarrow I.I. at $x=0$.

Set $f(x) = \frac{e^{\sqrt{x}} - 1}{x}$ & $g(x) = \sqrt{x}$.

$$\therefore \frac{f(x)}{g(x)} = \frac{\sqrt{x}(e^{\sqrt{x}} - 1)}{x} = \frac{\sqrt{x}(\sqrt{x} + h(x) \times x)}{x}$$

$\rightarrow 1$ as $x \rightarrow 0^+$.

\leftarrow why?

$\therefore \int_0^1 \sqrt{x} dx$ Converges, by Limit-Comparison test,
 $\int_0^1 \frac{e^{\sqrt{x}} - 1}{x} dx$ Converges.

(72)
 Jaydeb Sarkar.

14

(3) $\int_1^2 \frac{x^2 + x + 1}{(x^2 - 1)^{1/3}} dx.$

$f(x) := \frac{x^2 + x + 1}{(x^2 - 1)^{1/3}}$ ← Rational fn.

$g(x) := \frac{1}{(1-x)^{1/3}}$ ∴ We can deal with $\frac{1}{x+1}$ factor.

$\therefore \lim_{x \rightarrow 1^+} \frac{f(x)}{g(x)} = \lim_{x \rightarrow 1^+} \frac{(x^2 + x + 1)}{(x+1)^{1/3}} = \frac{3}{2^{1/3}} > 0.$

Now $\int_1^2 g = \int_1^2 \frac{1}{(1-x)^{1/3}} dx$

$= \int_0^1 \frac{1}{x^{1/3}} dx$ ← $\begin{cases} 1-x \mapsto x \\ \Rightarrow 1 \rightarrow 0 \\ \quad 2 \rightarrow 1 \end{cases}$

$\therefore p = \frac{1}{3} < 1, \int_1^2 g$ Converges.

$\Rightarrow \int_1^2 \frac{x^2 + x + 1}{(x^2 - 1)^{1/3}} \underline{\text{Converges.}}$ □

Def: An I.I. $\int_a^b f$ is said to be absolutely convergent if the I.I. $\int_a^b |f|$ is convergent.

Thm: Absolute Convergent \Rightarrow Convergent. ← Q: " \Leftarrow "?

Proof: Let $\int_a^b f$ be an I.I. at $x=a$.

$$\text{Now } -|f(x)| \leq f(x) \leq |f(x)| \quad \forall x \in (a, b].$$

$$\Rightarrow 0 \leq f(x) + |f(x)| \leq 2|f(x)| \quad \forall x \in (a, b].$$

\therefore the I.I. $\int_a^b |f|$ is absolutely conv. at a , by Comparison test, it follows that

$$\int_a^b (f(x) + |f(x)|) dx \text{ converges.}$$

Finally, $\forall a < c < b$, we have:

$$\begin{aligned} \int_c^b f &= \int_c^b ((f(x) + |f(x)|) - |f(x)|) dx \\ &= \int_c^b (f(x) + |f(x)|) dx - \int_c^b |f(x)| dx \end{aligned}$$

~~(both I.I. exist)~~

~~\therefore~~ $\int_a^b (f(x) + |f(x)|) dx$ & $\int_a^b |f(x)| dx$ both exist, it follows that

$\int_a^b f$ also exists.



eg: (Recall from page 72).

$$\int_0^1 \frac{\sin x}{x^2} \cdot dx.$$

$$\therefore \left| \frac{\sin x}{x^2} \right| \leq |\sin x| \quad \forall x \in (0, 1]$$

$\int_0^1 |\sin x| dx$ is convergent, it follows that

$$\int_0^1 \frac{\sin x}{x^2} \text{ is A.C. } \& \text{ hence } \underline{\text{convergent}}.$$



def: (I.I of type II):

Fix $a \in \mathbb{R}$ & suppose $f \in \mathcal{R}[a, b] \quad \forall b > a$.

if $\lim_{r \rightarrow +\infty} \int_a^r f$ exists, then we say that

$\int_a^{+\infty} f$ converges & we write:

$$\int_a^{+\infty} f = \lim_{r \rightarrow +\infty} \int_a^r f.$$

if $\lim_{r \rightarrow +\infty} \int_a^r f$ diverges, then we say $\int_a^{+\infty} f$ diverges.

114 $\int_{-\infty}^b f = \lim_{r \rightarrow -\infty} \int_r^b f$. \leftarrow if Converges.

Def: $\int_{-\infty}^{\infty} f := \int_{-\infty}^c f + \int_c^{\infty} f$

~~whenever the R.~~

Def: Let $f \in \mathcal{R}[a, b]$ $\forall a < b$ in \mathbb{R} . If $\exists c \in \mathbb{R}$ s.t.

both $\int_{-\infty}^c f$ & $\int_c^{\infty} f$ Converge, then we say that $\int_{-\infty}^{\infty} f$

Converges & write

$$\int_{-\infty}^{\infty} f = \int_{-\infty}^c f + \int_c^{\infty} f . \quad (*)$$

HW: If $\int_{-\infty}^{\infty} f$ Converges, then $(*)$ is independent of the
Choice of $c \in \mathbb{R}$.

eg: ① $\int_0^{\infty} \sin x \, dx$.

Let $f(x) = \sin x$. $x \in [0, \infty)$.

Note that $f \in \mathcal{R}[0, R]$ $\forall R \geq 0$.

$$\therefore \lim_{R \rightarrow \infty} \int_0^R f = \lim_{R \rightarrow \infty} \int_0^R \sin x \, dx$$

$$= \lim_{R \rightarrow \infty} [1 - \cos R]$$

But this limit DNE!

$\Rightarrow \int_0^{\infty} \sin x \, dx$ diverges.

OR Does not Converge.

$$\textcircled{2} \int_{-\infty}^0 e^{-x} dx.$$

Again, $e^{-x} \in \mathcal{R}[-R, 0] \quad \forall R > 0.$

Now, for $R > 0,$

$$\int_{-R}^0 e^{-x} dx = \int_0^R e^x dx \quad [x \rightarrow -x]$$

$$= e^R - 1.$$

$$\therefore \lim_{R \rightarrow \infty} \int_{-R}^0 e^{-x} dx = \lim_{R \rightarrow \infty} (e^R - 1) = \infty.$$

$$\therefore \int_{-\infty}^0 e^{-x} dx \text{ diverges .}$$

$$\textcircled{3} \int_{-\infty}^{\infty} \frac{dx}{1+x^2}.$$

We observe:

$$\int_0^{\infty} \frac{dx}{1+x^2} = \lim_{R \rightarrow \infty} \int_0^R \frac{dx}{1+x^2}$$

$$= \lim_{R \rightarrow \infty} [\tan^{-1}(R) - \tan^{-1}(0)].$$

why? $\Rightarrow \pi/2 - 0 = \pi/2.$

Also,

$$\int_{-\infty}^0 \frac{dx}{1+x^2} = \lim_{R \rightarrow \infty} \int_{-R}^0 \frac{dx}{1+x^2}$$

$$= \lim_{R \rightarrow \infty} \int_0^R \frac{dx}{1+x^2} = \pi/2.$$

$$\therefore \int_{-\infty}^{\infty} \frac{dx}{1+x^2} = \pi.$$

This is an even fcn.

[14]

Fact: (HW). Let $a > 0$. Then

$$\int_a^{\infty} \frac{1}{x^p} dx$$

Converges $\forall p > 1$ & diverges $\forall p \leq 1$.

Notation: For a fixed $a \in \mathbb{R}$, we say $f \in \mathcal{R}[a, \infty)$ if $f \in \mathcal{R}[a, R]$ $\forall R > a$.

Thm: (Comparison test - II).

Let $a \in \mathbb{R}$, $f, g \in \mathcal{R}[a, \infty)$ & let

$$0 \leq f(x) \leq g(x) \quad \forall x \in [a, \infty).$$

(i) If $\int_a^{\infty} g$ converges, then $\int_a^{\infty} f$ converges.

(ii) If $\int_a^{\infty} g$ diverges, then $\int_a^{\infty} f$ diverges.

CAUTION !!

We are not saying that f is Riemann integrable.

Proof: $\because 0 \leq f(x) \leq g(x) \quad \forall x \in [a, \infty)$ &

$f, g \in \mathcal{R}[a, \infty)$, it follows that

$$0 \leq \int_a^t f(x) dx \leq \int_a^t g(x) dx \quad \forall t > a.$$

$$\text{Set } F(t) := \int_a^t f(x) dx \quad \& \quad G(t) := \int_a^t g(x) dx. \quad \forall t > a.$$

$\therefore F, G \in \mathcal{C}(a, \infty)$ & monotonically increasing.

The result now follows immediately.

(Cont. of F & G is not needed).