

# **ECON-GA 2001**

## **First Half**

Quang Vuong

New York University

Revised October 2024

I. Probability Spaces and Random Variables

II. Measurable Functions and Lebesgue Integration

III. Expectations and Conditional Expectations

IV. Densities and Distributions

V. Modes of Convergence

VI. M-Estimation

VII. NLLS and ML Estimation

## I. Probability Spaces and Random Variables

$(\Omega, \mathcal{F}, P)$  Probability space

→  $\Omega$ : underlying space (nonempty)

→  $\mathcal{F}$ :  $\sigma$ -algebra or  $\sigma$ -field

→  $P$ : Probability measure

### I.1. $\sigma$ -algebra. (Bierens, Sections 1.1, 1.3–1.4)

DEFINITION: An *algebra/field* is a nonempty collection of subsets of  $\Omega$  satisfying

- (i) If  $A \in \mathcal{F}$  then  $A^c \in \mathcal{F}$  (closed under complementation)
- (ii) If  $A, B \in \mathcal{F}$  then  $A \cup B \in \mathcal{F}$  (closed under finite unions)

REMARKS:

- In the definition, (ii) can be replaced by

(ii') If  $A, B \in \mathcal{F}$  then  $A \cap B \in \mathcal{F}$  (closed under finite intersections)

- Examples:  $\mathcal{F}_1 = \{\emptyset, \Omega\}$  (trivial  $\sigma$ -algebra),  $\mathcal{F}_2 = \{\emptyset, A, A^c, \Omega\}$ ,  $\mathcal{F}_3 = 2^\Omega$  (power set)

DEFINITION: A  $\sigma$ -algebra/field is a nonempty collection of subsets of  $\Omega$  satisfying

- (i) If  $A \in \mathcal{F}$  then  $A^c \in \mathcal{F}$  (closed under complementation)
- (ii) If  $A_1, A_2, \dots \in \mathcal{F}$  then  $\cup_{i=1}^{\infty} A_i \in \mathcal{F}$  (closed under countable unions)

The pair  $(\Omega, \mathcal{F})$  is called a *measurable space*.

REMARKS:

- In the definition, (ii) can be replaced by

- (ii') If  $A_1, A_2, \dots \in \mathcal{F}$  then  $\cap_{i=1}^{\infty} A_i \in \mathcal{F}$  (closed under countable intersections)

- Examples:  $\mathcal{F}_1, \mathcal{F}_2, \mathcal{F}_3$  are  $\sigma$ -algebras

PROPERTIES OF  $\sigma$ -ALGEBRA:

- (a)  $\mathcal{F}$  is a  $\sigma$ -algebra  $\Rightarrow \mathcal{F}$  is an algebra. The reverse is not necessarily true.
- (b)  $\mathcal{F}$  is an algebra with a finite number of elements  $\Rightarrow \mathcal{F}$  is a  $\sigma$ -algebra.

- Let  $\mathcal{F}_\theta, \theta \in \Theta$  be a (possibly uncountable) collection of  $\sigma$ -algebras of  $\Omega$  indexed by  $\theta$ . Then
  - $\rightarrow \cap_{\theta \in \Theta} \mathcal{F}_\theta$  is a  $\sigma$ -algebra.
  - $\rightarrow \cup_{\theta \in \Theta} \mathcal{F}_\theta$  is not necessarily a  $\sigma$ -algebra.

**DEFINITION:** The smallest  $\sigma$ -algebra containing a collection  $\mathcal{C}$  of subsets of  $\Omega$  is called the  *$\sigma$ -algebra generated by  $\mathcal{C}$* . It is denoted  $\sigma(\mathcal{C})$  and satisfies  $\sigma(\mathcal{C}) = \cap_{\{\mathcal{F} : \mathcal{C} \subseteq \mathcal{F}\}} \mathcal{F}$ .

**EXAMPLES:**

- The  $\sigma$ -algebra generated by unions  $\vee_{\theta \in \Theta} \mathcal{F}_\theta \equiv \sigma(\cup_{\theta \in \Theta} \mathcal{F}_\theta)$
- Let  $\Omega = \mathbb{R}^k$  and  $\mathcal{C} = \{\times_{i=1}^k (a_i, b_i); a_i < b_i \in \mathbb{R}\}$ . Then
  - $\mathcal{B}^k \equiv \sigma(\mathcal{C})$  is the (Euclidean) *Borel  $\sigma$ -algebra* and any element of  $\mathcal{B}^k$  is called a *Borel set*.

**REMARKS:**

- $\rightarrow \mathcal{B}^k = \sigma(\{\times_{i=1}^k ([a_i, b_i]; a_i \leq b_i \in \mathbb{R})\}) = \sigma(\{\times_{i=1}^k (-\infty, a_i]; a_i \in \mathbb{R}\})$ , etc.
- $\rightarrow \mathcal{B}^k = \sigma(\{\text{Open sets of } \mathbb{R}^k\}) = \sigma(\{\text{Closed sets of } \mathbb{R}^k\})$
- $\rightarrow \mathcal{B}^k \neq 2^{\mathbb{R}^k}$  (Vitali example)

## I.2. Probability Measures. (Bierens, Sections 1.1, 1.5–1.7)

DEFINITION: A *measure*  $\mu$  on  $(\Omega, \mathcal{F})$  is a mapping  $\mathcal{F} \rightarrow \mathbb{R} \cup \{+\infty\}$  satisfying

- (i)  $\mu(A) \geq 0$  for all  $A \in \mathcal{F}$  (nonnegativity)
- (ii)  $\mu(\emptyset) = 0$
- (iii) If  $A_1, A_2, \dots \in \mathcal{F}$  are disjoint, then  $\mu(\cup_{i=1}^{\infty} A_i) = \sum_{i=1}^{\infty} \mu(A_i)$  (countable additivity).

REMARKS:  $(\Omega, \mathcal{F}, \mu)$  is called a *measured space*.

$\rightarrow \mu$  is *finite* if  $\mu(\Omega) < \infty$

$\rightarrow \mu$  is  $\sigma$ -*finite* if  $\Omega = \cup_{i=1}^{\infty} A_i$  with  $\mu(A_i) < \infty$

EXAMPLES:

- Let  $\mathcal{F} = 2^{\Omega}$  and  $\mu_c(A) \equiv \#A$ . Then  $\mu_c$  is called the *counting measure*. It is finite if  $|\Omega| < \infty$ .
- Let  $(\Omega, \mathcal{F}) = (\mathbb{R}^k, \mathcal{B}^k)$ . The *Lebesgue measure*  $\lambda$  on  $(\mathbb{R}^k, \mathcal{B}^k)$  is

$$\lambda(B) \equiv \inf_{B \subseteq \cup_{j=1}^{\infty} \{\times_{i=1}^k (a_{ij}, b_{ij})\}} \sum_{j=1}^{\infty} \prod_{i=1}^k (b_{ij} - a_{ij})$$

The Lebesgue measure  $\lambda$  is  $\sigma$ -finite.

DEFINITION: A *probability measure*  $P$  on a *measurable space*  $(\Omega, \mathcal{F})$  is a measure with (ii) replaced by (ii)  $P(\Omega) = 1$ . The triplet  $(\Omega, \mathcal{F}, P)$  is called a *probability space*.

REMARK: A probability measure  $P$  on an algebra  $\mathcal{A}$  is defined similarly by restricting  $\cup_{i=1}^{\infty} A_i \in \mathcal{A}$  in (iii). It can be uniquely extended to  $\sigma(\mathcal{A})$ . (Carathéodory Theorem)

EXAMPLE: Let  $(\Omega, \mathcal{F}) = (\{1, 2, 3, 4, 5, 6\}, 2^{\Omega})$ . The *uniform probability measure* is  $P_0(A) = \frac{\#A}{6}$ .

#### PROPERTIES OF MEASURES AND PROBABILITY MEASURES:

- (a)  $P(\emptyset) = 0$
- (b)  $P(A^c) = 1 - P(A)$  (only for probability measures)
- (c)  $A \subseteq B$  implies  $P(A) \leq P(B)$
- (d)  $P(A \cup B) = P(A) + P(B) - P(A \cap B)$
- (e) If  $A_1 \subseteq A_2 \subseteq \dots$ , then  $P(A_n) \uparrow P(\cup_{i=1}^{\infty} A_n)$
- (f) If  $A_1 \supseteq A_2 \supseteq \dots$ , then  $P(A_n) \downarrow P(\cap_{i=1}^{\infty} A_n)$  (only for finite measures)
- (g)  $P(\cup_{i=1}^{\infty} A_n) \leq \sum_{i=1}^{\infty} P(A_n)$

### I.3. Random Variables/Vectors. (Bierens, Sections 1.8, 1.10)

DEFINITION: Let  $(\Omega, \mathcal{F}, P)$  be a probability space and  $(\tilde{\Omega}, \tilde{\mathcal{F}})$  be a measurable space. A mapping  $X : \Omega \rightarrow \tilde{\Omega}$  is a random element iff  $X(\cdot)$  is  $\mathcal{F}/\tilde{\mathcal{F}}$ -measurable, i.e. iff

$$X^{-1}(\tilde{B}) \in \mathcal{F} \quad \text{for every} \quad \tilde{B} \in \tilde{\mathcal{F}}$$

where  $X^{-1}(\tilde{B}) \equiv \{\omega \in \Omega : X(\omega) \in \tilde{B}\}$ .

EXAMPLE: When  $\tilde{\Omega} = \{f(\cdot) : \mathcal{T} \rightarrow \mathbb{R}\}$  and  $\mathcal{F}$  is the  $\sigma$ -algebra generated by the open sets relative to a metric on  $\tilde{\Omega}$ , then  $X(\cdot)$  is a *random process* indexed by  $\mathcal{T}$  and denoted  $X(\omega; t)$ .

REMARKS:

- The collection  $\mathcal{F}_X \equiv \{X^{-1}(\tilde{B}) : \tilde{B} \in \tilde{\mathcal{F}}\}$  is a  $\sigma$ -algebra called the  *$\sigma$ -algebra generated by  $X$* .

$X(\cdot)$  is a random element iff  $\mathcal{F}_X \subseteq \mathcal{F}$ .

- Let  $P_X : \tilde{\mathcal{F}} \rightarrow \mathbb{R} \cup \{+\infty\}$  defined by  $P_X(\tilde{B}) \equiv P[X^{-1}(\tilde{B})]$  for every  $\tilde{B} \in \tilde{\mathcal{F}}$ . Then

$(\tilde{\Omega}, \tilde{\mathcal{F}}, P_X)$  is a probability space, and  $P_X$  is called the *probability measure induced by  $X$* .

DEFINITION: When  $(\tilde{\Omega}, \tilde{\mathcal{F}}) = (\mathbb{R}^k, \mathcal{B}^k)$ , the random element  $X : \Omega \rightarrow \mathbb{R}^k$  is called a *random variable/vector* (r.v.).

REMARKS:

- The  $\mathcal{F}/\mathcal{B}^k$ -measurability condition  $\{X^{-1}(\tilde{B}) : \tilde{B} \in \mathcal{B}^k\} \subseteq \mathcal{F}$  is equivalent to

$$\{X^{-1}((-\infty, x]) : x \in \mathbb{R}^k\} \subseteq \mathcal{F}$$

- Let  $F_X : \mathbb{R}^k \rightarrow [0, 1]$  defined by  $F(x) \equiv P_X[(-\infty, x]] = P\{X^{-1}[(-\infty, x]]\}$  for every  $x \in \mathbb{R}^k$ .

$F_X(\cdot)$  is called the *distribution function of X*.

PROPERTIES OF DISTRIBUTION FUNCTIONS:

(a) A distribution function  $F_X$  on  $\mathbb{R}^k$  corresponds to a unique probability measure  $P_X$  satisfying

$$P_X[(-\infty, x]] = F_X(x) \text{ for every } x \in \mathbb{R}^k.$$

(b) A distribution function  $F_X(\cdot)$  is monotone nondecreasing, right-continuous with left-limits

(*cadlag*), at most countable discontinuities,  $\lim_{x \downarrow -\infty} F_X(x) = 0$  and  $\lim_{x \uparrow +\infty} F_X(x) = 1$ .



DEFINITION: (i) The events  $A_1, A_2, \dots$  in  $\mathcal{F}$  of a probability space  $(\Omega, \mathcal{F}, P)$  are (mutually) *independent* iff for every finite collection  $A_{j_1}, \dots, A_{j_n}$  and every  $n$

$$P(\cap_{i=1}^n A_{j_i}) = \prod_{i=1}^n P(A_{j_i})$$

(ii) The  $\sigma$ -algebras  $\mathcal{F}_1, \mathcal{F}_2, \dots$  included in  $\mathcal{F}$  on a probability space  $(\Omega, \mathcal{F}, P)$  are (mutually) *independent* iff every sequence  $A_1, A_2, \dots$  with  $A_j \in \mathcal{F}_j$  is independent.

(iii) The r.v.  $X_1, X_2, \dots$  on a probability space  $(\Omega, \mathcal{F}, P)$  are (mutually) *independent* iff the sequence  $\mathcal{F}_{X_1}, \mathcal{F}_{X_2}, \dots$  is independent.

PROPERTY OF INDEPENDENCE:

- The r.v.  $X_1, X_2, \dots$  on a probability space  $(\Omega, \mathcal{F}, P)$  are (mutually) independent iff

$$F_X(x) = \prod_{i=1}^n F_{X_{j_i}}(x_{j_i})$$

for every  $x = (x_{j_1}, \dots, x_{j_n})$ ,  $(j_1, \dots, j_n)$  and  $n$ , where  $X = (X_{j_1}, \dots, X_{j_n})$ .

$F_X(\cdot)$  and  $F_{X_j}(\cdot)$  are called the *joint and marginal distribution functions*, respectively.

## II. Measurable Functions and Lebesgue Integration

### II.1. Measurable Functions. (Bierens, Sections 2.2, 2.4)

DEFINITION: A  $\mathcal{F}/\mathcal{B}^k$ -measurable function  $g(\cdot)$  is a mapping from a measurable space  $(\Omega, \mathcal{F})$  to  $(\mathbb{R}^k, \mathcal{B}^k)$  that satisfies  $g^{-1}(B) \in \mathcal{F}$  for every  $B \in \mathcal{B}^k$  where  $g^{-1}(B) \equiv \{\omega : g(\omega) \in B\}$ .

EXAMPLES:

- Same as a r.v. except that the latter is defined on  $(\Omega, \mathcal{F}, P)$ .
- When  $(\Omega, \mathcal{F}) = (\mathbb{R}^\ell, \mathcal{B}^\ell)$ , then  $g(\cdot)$  is simply called a *measurable function*.
  - Continuous functions from  $\mathbb{R}^\ell$  to  $\mathbb{R}^k$  are measurable functions. The reverse is not true.

PROPERTIES:

- (a)  $g(\cdot) = [g_1(\cdot), \dots, g_k(\cdot)]'$  is  $\mathcal{F}/\mathcal{B}^k$ -measurable iff  $g_i(\cdot)$  is  $\mathcal{F}/\mathcal{B}$ -measurable for every  $i = 1, \dots, k$ .
- (b) Let  $X(\cdot)$  and  $Y(\cdot)$  be two r.v. on  $(\Omega, \mathcal{F}, P)$ . Then  $\mathcal{F}_Y \subseteq \mathcal{F}_X$  iff there exists a measurable function  $g(\cdot) : \mathbb{R}^{\dim X} \rightarrow \mathbb{R}^{\dim Y}$  such that  $Y(\omega) = g[X(\omega)]$  for all  $\omega \in \Omega$ .

DEFINITION: A *simple function*  $g(\cdot)$  is a  $\mathcal{F}/\mathcal{B}$ -measurable function from  $(\Omega, \mathcal{F})$  to  $(\mathbb{R}, \mathcal{B})$  of the form

$$g(\omega) = \sum_{j=1}^m b_j \mathbb{I}(\omega \in A_j)$$

where  $m < \infty$ ,  $b_1, \dots, b_m \in \mathbb{R}$  and  $\{A_j\}_{j=1}^m$  a partition of  $\Omega$  with  $A_j \in \mathcal{F}$  for  $j = 1, \dots, m$ .

PROPERTIES:

- (a) A function  $g(\cdot) : (\Omega, \mathcal{F}) \rightarrow (\mathbb{R}, \mathcal{B})$  is  $\mathcal{F}/\mathcal{B}$ -measurable iff it is the (pointwise) limit of a sequence of simple functions  $\{g_n(\cdot)\}_{n=1}^\infty$ , i.e.  $\lim_{n \rightarrow \infty} g_n(\omega) = g(\omega)$  for every  $\omega \in \Omega$ .
- (b) If  $f(\cdot)$  and  $g(\cdot)$  are  $\mathcal{F}/\mathcal{B}$ -measurable, then  $f(\cdot) + g(\cdot)$ ,  $f(\cdot) - g(\cdot)$ ,  $f(\cdot) \times g(\cdot)$  and  $f(\cdot)/g(\cdot)$  (with  $g(\cdot) \neq 0$ ) are  $\mathcal{F}/\mathcal{B}$ -measurable,
- (c) Let  $\{g_n(\cdot)\}_{n=1}^\infty$  be a sequence of  $\mathcal{F}/\mathcal{B}$ -measurable functions. Provided the RHS are well-defined,
  - (c.1)  $\underline{f}(\cdot) \equiv \inf_{n \geq 1} g_n(\cdot)$  and  $\bar{f}(\cdot) \equiv \sup_{n \geq 1} g_n(\cdot)$  are  $\mathcal{F}/\mathcal{B}$ -measurable,
  - (c.2)  $\underline{h}(\cdot) \equiv \liminf_{n \rightarrow \infty} g_n(\cdot)$  and  $\bar{h}(\cdot) \equiv \limsup_{n \rightarrow \infty} g_n(\cdot)$  are  $\mathcal{F}/\mathcal{B}$ -measurable,
  - (c.3)  $g(\cdot) \equiv \lim_{n \rightarrow \infty} g_n(\cdot)$  is  $\mathcal{F}/\mathcal{B}$ -measurable.

## II.2. Lebesgue Integration. (Bierens, Sections 2.2, 2.4, 4.3–4.4)

DEFINITION: Let  $(\Omega, \mathcal{F}, \mu)$  be a measured space.

(i) The *integral* of a simple function  $g(\cdot)$  w.r.t.  $\mu$  is

$$\int_{\Omega} g(\omega) d\mu(\omega) \equiv \sum_{j=1}^m b_j \mu(A_j)$$

(ii) The *integral* of a  $(\mu$ -a.e.) nonnegative  $\mathcal{F}/\mathcal{B}$ -measurable function  $g(\cdot)$  w.r.t.  $\mu$  is

$$\int_{\Omega} g(\omega) d\mu(\omega) \equiv \sup_{0 \leq g_S(\cdot) \leq g(\cdot)} \int_{\Omega} g_S(\omega) d\mu(\omega)$$

where  $g_S(\cdot)$  is a simple function.

(iii) The *integral* of a  $\mathcal{F}/\mathcal{B}$ -measurable function  $g(\cdot)$  w.r.t.  $\mu$  is

$$\int_{\Omega} g(\omega) d\mu(\omega) \equiv \int_{\Omega} g_+(\omega) d\mu(\omega) - \int_{\Omega} g_-(\omega) d\mu(\omega)$$

where  $g_+(\cdot) \equiv \max\{g(\cdot), 0\}$  and  $g_-(\cdot) \equiv \max\{-g(\cdot), 0\}$  provided  $\int_{\Omega} g_+(\omega) d\mu(\omega) < \infty$  or  $\int_{\Omega} g_-(\omega) d\mu(\omega) < \infty$ . If both are finite, then  $g(\cdot)$  is  $\mu$ -integrable.

### REMARKS:

- $g(\cdot)$  is  $\mu$ -integrable iff  $|g(\cdot)|$  is  $\mu$ -integrable, in which case  $|\int_{\Omega} g(\omega)d\mu(\omega)| \leq \int_{\Omega} |g(\omega)|d\mu(\omega)$ .
- If  $g(\cdot)$  is  $\mu$ -integrable, then  $g(\cdot) < \infty$   $\mu$ -a.e. (allowing  $g(\cdot)$  to have  $\pm\infty$  values)
- Let  $\int_A g(\omega)d\mu(\omega) \equiv \int_{\Omega} \mathbb{I}(\omega \in A)g(\omega)d\mu(\omega)$  for  $A \in \mathcal{F}$ . If  $\mu(A) = 0$ , then  $\int_A g(\omega)d\mu(\omega) = 0$ .
- When  $(\Omega, \mathcal{F}) = (\mathbb{R}, \lambda)$ , then  $\int_A g(x)d\lambda(x)$  is called the *Lebesgue integral of  $g(\cdot)$  on  $A$* .
  - If a bounded function  $g(\cdot)$  is Riemann integrable, then  $g(\cdot)$  is Lebesgue integrable and  $\int_a^b g(x)dx = \int_{[a,b]} g(x)d\lambda(x) < \infty$ . Counterexample:  $g(\cdot)$  is the Dirichlet function.
  - A bounded function  $g(\cdot) : [a, b] \rightarrow \mathbb{R}$  is Riemann integrable iff its set of discontinuities is of  $\lambda$ -measure zero. (Lebesgue Theorem)

**PROPERTIES OF LEBESGUE INTEGRATION:** Let  $g(\cdot)$  and  $h(\cdot)$  be  $\mu$ -integrable. Then

- (a) If  $g(\cdot) = 0$   $\mu$ -a.e., then  $\int_{\Omega} g(\omega)d\mu(\omega) = 0$ .
- (b)  $\int_{\Omega} [\alpha g(\omega) + \beta h(\omega)]d\mu(\omega) = \alpha \int_{\Omega} g(\omega)d\mu(\omega) + \beta \int_{\Omega} h(\omega)d\mu(\omega)$  for  $\alpha, \beta \in \mathbb{R}$
- (c) If  $g(\cdot) \leq h(\cdot)$   $\mu$ -a.e., then  $\int_{\Omega} g(\omega)d\mu(\omega) \leq \int_{\Omega} h(\omega)d\mu(\omega)$

## LIMITS AND LEBESGUE INTEGRATION:

- Let  $g(\cdot)$  be  $\mu$ -integrable, and  $A_1, A_2, \dots \in \mathcal{F}$ .

If  $\lim_{n \rightarrow \infty} \mu(A_n) = 0$ , then  $\lim_{n \rightarrow \infty} \int_{A_n} g(\omega) d\mu(\omega) = 0$ .

- *Monotone Convergence Theorem*: Let  $\{g_n(\cdot)\}_{n=1}^{\infty}$  be a nondecreasing sequence of nonnegative  $\mathcal{F}/\mathcal{B}$ -measurable functions converging (pointwise) to a function  $g(\cdot)$   $\mu$ -a.e. Then

$$\lim_{n \rightarrow \infty} \int_{\Omega} g_n(\omega) d\mu(\omega) = \int_{\Omega} g(\omega) d\mu(\omega)$$

- *Lebesgue Dominated Convergence Theorem*: Let  $\{g_n(\cdot)\}_{n=1}^{\infty}$  be a sequence of  $\mathcal{F}/\mathcal{B}$ -measurable functions converging (pointwise) to a function  $g(\cdot)$  satisfying  $|g_n(\cdot)| \leq h(\cdot)$   $\mu$ -a.e. for every  $n$  where  $h(\cdot)$  is  $\mu$ -integrable. Then, the previous conclusion holds.

REMARK: Let  $Q(\theta) \equiv \int_{\Omega} g(\omega; \theta) d\mu(\omega)$  where  $g(\cdot; \theta)$  is  $\mu$ -integrable for every  $\theta \in \Theta \subset \mathbb{R}^p$ . Then

(i)  $Q(\cdot)$  is continuous and

(ii)  $Q'(\theta) = \int_{\Omega} g'(\omega; \theta) d\mu(\omega)$ , where  $g'(\omega; \cdot)$  is the derivative of  $g(\omega; \cdot)$ .

See e.g. Theorem 16.8 in Billingsley (1995) for conditions.

CHANGE-OF-VARIABLE: Let  $g(\cdot)$  and  $h(\cdot)$  be  $\mathcal{B}^k/\mathcal{B}$  and  $\mathcal{B}^k/\mathcal{B}^k$ -measurable. If  $h(\cdot)$  is one-to-one from  $A \in \mathcal{B}^k \rightarrow B \in \mathcal{B}^k$ , and continuously differentiable with Jacobian  $J(x) \equiv \partial h / \partial x'$ , then

$$\int_B g(y) d\lambda(y) = \int_A g[h(x)] \cdot |J(x)| d\lambda(x).$$

PRODUCT MEASURE AND FUBINI'S THEOREM. (Billingsley, Section 18)

Let  $(\Omega, \mathcal{F}, \mu)$  and  $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mu})$  be two measure spaces with  $\sigma$ -finite measures  $\mu$  and  $\tilde{\mu}$ .

DEFINITION: The *product  $\sigma$ -algebra* on  $\Omega \times \tilde{\Omega}$  is  $\mathcal{F} \times \tilde{\mathcal{F}} \equiv \sigma(\{A \times \tilde{A} : A \in \mathcal{F}, \tilde{A} \in \tilde{\mathcal{F}}\})$ .

The *product measure*  $\nu$  on  $\mathcal{F} \times \tilde{\mathcal{F}}$  is the unique  $\sigma$ -finite measure satisfying

$$\nu(A \times \tilde{A}) = \mu(A) \cdot \tilde{\mu}(\tilde{A}) \quad \text{for every } A \in \mathcal{F}, \tilde{A} \in \tilde{\mathcal{F}}.$$

The triplet  $(\Omega \times \tilde{\Omega}, \mathcal{F} \times \tilde{\mathcal{F}}, \nu)$  is called the *product measure space*.

PROPERTY: (Fubini's Theorem) Let  $g(\cdot) : \Omega \times \tilde{\Omega} \rightarrow \mathbb{R}$  be  $(\mathcal{F} \times \tilde{\mathcal{F}})/\mathcal{B}$ -measurable. Then

$$\int_{\Omega \times \tilde{\Omega}} g(\varpi) d\nu(\varpi) = \int_{\Omega} \left[ \int_{\tilde{\Omega}} g(\omega, \tilde{\omega}) d\tilde{\mu}(\tilde{\omega}) \right] d\mu(\omega) = \int_{\tilde{\Omega}} \left[ \int_{\Omega} g(\omega, \tilde{\omega}) d\mu(\omega) \right] d\tilde{\mu}(\tilde{\omega})$$

where  $\varpi = (\omega, \tilde{\omega})$ , provided one of the integrals is finite.

### III. Expectations and Conditional Expectations

#### III.1. Expectations. (Bierens, Sections 2.5–2.8, 5.1)

DEFINITION: Let  $(\Omega, \mathcal{F}, P)$  be a probability space,  $X(\cdot)$  a r.v. in  $\mathbb{R}^{\dim X}$ , and  $g(\cdot) : \mathbb{R}^{\dim X} \rightarrow \mathbb{R}^k$  measurable. Then  $E[g(X)] \equiv \int_{\Omega} g[X(\omega)]dP(\omega) \equiv \left[ \int_{\Omega} g_1[X(\omega)]dP(\omega), \dots, \int_{\Omega} g_k[X(\omega)]dP(\omega) \right]'$ .

EXAMPLES:

- When  $\dim X = 1$ , the *raw and central p-th moment* of  $X$  are  $E[X^p]$  and  $E[X - E(X)]^p$ .  
→  $E[X]$  and  $E[X - E(X)]^2$  are the *mean* and *variance* of  $X$ .
- The *variance-covariance matrix* of  $X$  is  $E[X - E(X)][X - E(X)]' = E[XX'] - E[X]E[X]'$ .

PROPERTIES:

- (a)  $E[g(X)] = \int_{\mathbb{R}^{\dim X}} g(x)dP_X(x) = \int_{\mathbb{R}^{\dim Y}} ydP_Y(y)$  where  $Y \equiv g(X)$ , provided one of the integrals is finite.
- (b) The r.v.  $X$  and  $Y$  are independent iff  $E[g(X)h(Y)] = E[g(X)] \cdot E[h(Y)]$  for all measurable functions  $g(\cdot)$  and  $h(\cdot)$  on  $\mathbb{R}^{\dim X}$  and  $\mathbb{R}^{\dim Y}$ .



### SOME INEQUALITIES:

- Markov inequality  $\Pr[|X| \geq \epsilon] \leq E[|X|^p]/\epsilon^p$  for every  $p > 0$ . ( $\Rightarrow$  Chebyshev inequality)
- Jensen inequality  $E[g(X)] \leq g(E[X])$  for  $g(\cdot)$  concave on  $\mathcal{S}_X$ . ( $\Rightarrow$  Liapounov inequality)
- Holder inequality  $E(|XY|) \leq (E[|X|^p])^{1/p}(E[|Y|^q])^{1/q}$  for  $1/p+1/q=1$ . ( $\Rightarrow$  Cauchy-Schwarz in.)
- Triangle inequality  $(E[|X + Y|^p])^{1/p} \leq (E[|X|^p])^{1/p} + (E[|Y|^p])^{1/p}$  for  $p \geq 1$ . ( $\Rightarrow$  Minkowski in.)

DEFINITION: The *moment generating function* of  $X$  is  $M_X(t) \equiv E[\exp(t'X)]$  for  $t \in \mathcal{D}_X \subseteq \mathbb{R}^{\dim X}$ . The *characteristic function* of  $X$  is  $\phi_X(t) \equiv E[\exp(it'X)]$  for  $t \in \mathbb{R}^{\dim X}$ .

### PROPERTIES:

(a) If  $(-t_o, t_o) \subseteq \mathcal{D}_X \subseteq \mathbb{R}$  with  $t_o > 0$ , then  $M^{(p)}(0) = E[X^p] < \infty$  for all  $p \in \mathbb{N}$ ,

$M_X(t) = \sum_{p=0}^{\infty} \frac{E[X^p]}{p!} t^p$  and  $P_X(\cdot)$  is uniquely determined by  $M_X(\cdot)$  (and by the moments of  $X$ ).

(b)  $P_X(\cdot)$  is uniquely determined by  $\phi_X(\cdot)$ .

When  $\dim X = 1$ ,  $\phi_X(\cdot)$  is uniformly continuous. Moreover,  $\phi^{(p)}(0) = i^p E[X^p]$  if  $E[X^p] < \infty$ .

(c)  $\phi_X(t) = M_X(it)$  and  $M_X(t) = \phi_X(-it)$  whenever  $M_X(\cdot)$  is well-defined.

### III.2. Conditional Expectations. (Bierens, Chapter 3)

→  $E[Y|X]$  where  $Y(\cdot)$  is a r.v. and  $X(\cdot)$  is a random element on  $(\Omega, \mathcal{F}, P)$

EXAMPLE:  $(\Omega, \mathcal{F}, P) = (\{1, 2, 3, 4, 5, 6\}, 2^\Omega, P_0)$  where  $P_0$  is the uniform probability measure.

Let  $Y(\omega) = \omega$  and  $X(\omega) = \mathbb{I}[\omega \in \{2, 4, 6\}]$ . Then  $E[Y|X] = 3 + X$ .

DEFINITION: Let  $Y$  be a random variable on  $(\Omega, \mathcal{F}, P)$  with  $E[Y] < \infty$ . Let  $\mathcal{F}_0 \subseteq \mathcal{F}$  be a  $\sigma$ -algebra. The *conditional expectation of  $Y$  given  $\mathcal{F}_0$*  is a mapping  $E[Y|\mathcal{F}_0](\cdot) : \Omega \rightarrow \mathbb{R}$  satisfying

- (i)  $E[Y|\mathcal{F}_0](\cdot)$  is  $\mathcal{F}_0/\mathcal{B}$ -measurable,
- (ii)  $\int_A E[Y|\mathcal{F}_0](\omega) dP(\omega) = \int_A Y(\omega) dP(\omega)$ , i.e.,  $E\{\mathbb{I}_A E[Y|\mathcal{F}_0]\} = E[\mathbb{I}_A Y]$  for every  $A \in \mathcal{F}_0$ .

REMARKS:

- $E[Y|\mathcal{F}_0](\cdot)$  is uniquely defined  $P$ -a.s.
- The *conditional expectation of  $Y$  given  $X$*  is  $E[Y|X](\cdot) \equiv E[Y|\mathcal{F}_X](\cdot)$ .
  - When  $X \in \mathbb{R}^k$ , then  $E[Y|X](\cdot) = g[X(\cdot)]$  for some  $\mathcal{B}^k/\mathcal{B}$ -measurable function  $g(\cdot)$ .
  - $E[Y|X=x] \equiv g(x) = E[Y|X](\omega)$  for all  $\omega \in X^{-1}(x)$ .

## PROPERTIES OF CONDITIONAL EXPECTATIONS:

- (a) If  $Y = c$ , then  $E[Y|\mathcal{F}_0] = c$ .
- (b)  $E[\alpha X + \beta Y|\mathcal{F}_0] = \alpha E[X|\mathcal{F}_0] + \beta E[Y|\mathcal{F}_0]$ .
- (c) If  $X(\cdot) \leq Y(\cdot)$ , then  $E[X|\mathcal{F}_0] \leq E[Y|\mathcal{F}_0]$ .
- (d) If  $\mathcal{F}_0 = \{\emptyset, \Omega\}$ , then  $E[Y|\mathcal{F}_0] = E[Y]$ .
- (e) If  $\mathcal{F}_Y \perp \mathcal{F}_0$ , then  $E[Y|\mathcal{F}_0] = E[Y]$ .
- (f) If  $\mathcal{F}_X \subseteq \mathcal{F}_0$ , then  $E[X|\mathcal{F}_0] = X$  and  $E[XY|\mathcal{F}_0] = XE[Y|\mathcal{F}_0]$ .
- (g) *Law of Iterated Expectations:* If  $\mathcal{F}_0 \subseteq \mathcal{F}_1 \subseteq \mathcal{F}$ , then

$$E\{E[Y|\mathcal{F}_0]|\mathcal{F}_1\} = E\{E[Y|\mathcal{F}_1]|\mathcal{F}_0\} = E[Y|\mathcal{F}_0].$$

- (h) The r.v.s  $X$  and  $Y$  are *conditionally independent* given  $\mathcal{F}_0$  iff  $E[g(X)h(Y)|\mathcal{F}_0] = E[g(X)|\mathcal{F}_0] \cdot E[h(Y)|\mathcal{F}_0]$  for all measurable functions  $g(\cdot)$  and  $h(\cdot)$  on  $\mathbb{R}^{\dim X}$  and  $\mathbb{R}^{\dim Y}$ .
- (i) *Best MSE:* If  $E[Y^2] < \infty$ , then  $E[Y|X] = \arg \min_{h \in \mathcal{H}} MSE(h) \equiv E[Y - h(X)]^2$  where  $\mathcal{H}$  is the set of squared-integrable (measurable) functions of  $X$ .

## LIMITS AND CONDITIONAL EXPECTATIONS:

- *Monotone Convergence Theorem:* Let  $\{Y_n(\cdot)\}_{n=1}^{\infty}$  be a nondecreasing sequence of nonnegative r.v.s with  $E[\sup_{n \geq 1} Y_n] < \infty$ . Then  $\lim_{n \rightarrow \infty} E[Y_n | \mathcal{F}_0] = E[\lim_{n \rightarrow \infty} Y_n | \mathcal{F}_0]$ .
- *Dominated Convergence Theorem:* Let  $\{Y_n(\cdot)\}_{n=1}^{\infty}$  be a sequence of r.v.s converging to  $Y(\cdot)$   $P$ -a.s. and satisfying  $|Y_n(\cdot)| \leq M(\cdot)$  with  $E[M | \mathcal{F}_0] < \infty$ . Then,  $\lim_{n \rightarrow \infty} E[Y_n | \mathcal{F}_0] = E[Y | \mathcal{F}_0]$ .
- Let  $\mathcal{F}_1 \subseteq \mathcal{F}_2 \subseteq \dots \subseteq \mathcal{F}$ . Then  $\lim_{n \rightarrow \infty} E[Y | \mathcal{F}_n] = E[Y | \mathcal{F}_{\infty}]$  where  $\mathcal{F}_{\infty} \equiv \bigvee_{n=1}^{\infty} \mathcal{F}_n = \sigma(\bigcup_{n=1}^{\infty} \mathcal{F}_n)$ .  
 $\rightarrow \lim_{n \rightarrow \infty} E[Y_t | Y_{t-1}, \dots, Y_{t-n}] = E[Y_t | Y_{t-1}, Y_{t-2}, \dots]$

DEFINITION: Let  $Y$  be a r.v. on  $(\Omega, \mathcal{F}, P)$ . The *conditional probability measure of  $Y \in \mathbb{R}^{\ell}$  given  $\mathcal{F}_0 \subseteq \mathcal{F}$*  is the mapping  $P_{Y|\mathcal{F}_0}(\cdot, \cdot) : (B, \omega) \in \mathcal{B}^{\ell} \times \Omega \rightarrow P_{Y|\mathcal{F}_0}(B, \omega) \equiv E[\mathbb{I}(Y \in B) | \mathcal{F}_0](\omega)$ .

## REMARKS:

- For each  $\omega \in \Omega$ ,  $(\mathbb{R}, \mathcal{B}^{\ell}, P_{Y|\mathcal{F}_0}(\cdot, \omega))$  is a probability space.
- The *cond. prob. measure of  $Y \in \mathbb{R}^{\ell}$  given  $X \in \mathbb{R}^k$*  is  $P_{Y|X}(\cdot | x) \equiv P_{Y|\mathcal{F}_X}(\cdot, \omega)$  for  $\omega \in X^{-1}(x)$ .  
 $\rightarrow \int_A P_{Y|X}(B | x) dP_X(x) = P[X \in A, Y \in B]$  for every  $(A, B) \in \mathcal{B}^k \times \mathcal{B}^{\ell}$ .

## IV. Densities and Distributions

### IV.1. Densities. (Bierens, Sections 1.9)

DEFINITION: Let  $X$  be r.v. from  $(\omega, \mathcal{F}, P)$  to  $(\mathbb{R}^k, \mathcal{B}^k, P_X)$ . Then  $P_X$  is *absolutely continuous w.r.t. a measure  $\mu$  on  $(\mathbb{R}^k, \mathcal{B}^k)$* , i.e.  $P_X \ll \mu$ , iff  $\mu(B) = 0 \Rightarrow P_X(B) = 0$  for every  $B \in \mathcal{B}^k$ .

EXAMPLES:  $k = 1$

- When  $\mu(B) = \mu_c(B) \equiv \#(B \cap \mathbb{N})$ , then  $P_X$  (or  $X$ ) is *discrete* and  $p_X(i) \equiv P[X = i]$ .
- When  $\mu(B) = \lambda(B)$ , then  $P_X$  (or  $X$ ) is *(absolutely) continuous*.

RADON-NIKODYM THEOREM: If  $P_X \ll \mu$  with  $\sigma$ -finite  $\mu$ , then there exists a nonnegative  $\mathcal{B}^k/\mathcal{B}$ -measurable function  $f_X(\cdot)$  satisfying  $P_X(B) = \int_B f_X(x) d\mu(x)$  for every  $B \in \mathcal{B}^k$ .

→ The *(probability) density*  $f_X(\cdot)$  is  $\mu$ -a.e. unique.

→  $\int_B g(x) dP_X(x) = \int_B g(x) f_X(x) d\mu(x)$  and  $F_X(\cdot) = \int_{(-\infty, \cdot]} f_X(x) d\mu(x)$ .

→ When  $\mu = \mu_c$ , then  $\int_B g(x) d\mu_c(x) = \sum_{i \in B} g(i) p_X(i)$  and  $F_X(x) = \sum_{i \leq x} p_X(i)$ .

→ When  $\mu = \lambda$  and  $F_X(\cdot)$  is continuously differentiable, then  $f_X(\cdot) = F'_X(\cdot)$ .

## IV.2. Some Discrete and Continuous R.V.s. (Bierens, Sections 4.1, 4.5–4.6)

### SOME UNIVARIATE DISCRETE R.V.s:

→ *Binomial Distribution*:  $X \sim \mathcal{B}(n, p)$  where  $n \in \mathbb{N}$  and  $0 < p < 1$ .

$$p_X(i) = \binom{n}{p} p^i (1-p)^{n-i} \text{ for } i = 0, 1, \dots, n.$$

$$E[X] = np, \text{ Var}[X] = np(1-p), \phi_X(t) = [1 - p + p \exp(it)]^n.$$

→ *Poisson Distribution*:  $X \sim \mathcal{P}(\lambda)$  where  $\lambda > 0$

$$p_X(i) = \frac{\lambda^i}{i!} \exp(-\lambda) \text{ for } i = 0, 1, 2, \dots$$

$$E[X] = \lambda, \text{ Var}[X] = \lambda, \phi_X(t) = \exp\{\lambda[\exp(it) - 1]\}.$$

### SOME UNIVARIATE CONTINUOUS R.V.s:

→ *Normal Distribution*:  $X \sim \mathcal{N}(\mu, \sigma^2)$  where  $\mu \in \mathbb{R}$  and  $\sigma^2 > 0$ .

$$f_X(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp\left[-\frac{1}{2} \frac{(x-\mu)^2}{\sigma^2}\right] \text{ for } x \in \mathbb{R}.$$

$$E[X] = \mu, \text{ Var}[X] = \sigma^2, \phi_X(t) = \exp(i\mu t - \sigma^2 t^2/2).$$

*Standard Normal Distribution Z*:  $\mu = 0$  and  $\sigma^2 = 1$ .

# SOME UNIVARIATE CONTINUOUS R.V.s: (continued)

→ *Chi-Square Distribution*:  $X \sim \chi_k^2 = \sum_{j=1}^k Z_j^2$  where  $k \in \mathbb{N}$  and  $Z_j$  are i.i.d.  $\mathcal{N}(0, 1)$ .

$$f_X(x) = \frac{1}{2^{k/2}\Gamma(k/2)} x^{k/2-1} \exp(-x/2) \text{ for } x \geq 0, \text{ where } \Gamma(\alpha) \equiv \int_0^\infty x^{\alpha-1} \exp(-x) dx.$$

$$E[X] = k, \text{ Var}[X] = 2k, \phi_X(t) = \frac{1}{(1-2it)^{k/2}}.$$

→ *Student-t Distribution*:  $X \sim t_k = \frac{Z}{\sqrt{\chi_k^2/k}}$  where  $k \in \mathbb{N}$  and  $Z \sim \mathcal{N}(0, 1) \perp \chi_k^2$ .

$$f_X(x) = \frac{1}{\sqrt{k}B(\frac{1}{2}, \frac{k}{2})} \left(1 + \frac{x^2}{k}\right)^{-\frac{k+1}{2}} \text{ for } x \in \mathbb{R}, \text{ where } B(\alpha, \beta) \equiv \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}.$$

$$E[X] = 0 \text{ for } k \geq 2, \text{ Var}[X] = \frac{k}{k-2} \text{ for } k \geq 3, \phi_X(t) = \frac{2 \cdot \Gamma[(k+1)/2]}{\sqrt{k\pi} \cdot \Gamma(k/2)} \int_0^\infty \frac{\cos(tx)}{(1+x^2/k)^{(k+1)/2}} dx.$$

*Cauchy Distribution*:  $k = 1$ .  $E[X^p]$  does not exist or is infinite for any  $p \in \mathbb{N}$ .

→ *F Distribution*:  $X \sim F(m, n) = \frac{\chi_m^2/m}{\chi_n^2/n}$  where  $m, n \in \mathbb{N}$  and  $\chi_m^2 \perp \chi_n^2$ .

$$f_X(x) = \frac{m^{m/2} n^{n/2}}{B(m/2, n/2)} \frac{x^{m/2-1}}{(n+mx)^{(m+n)/2}} \text{ for } x \in [0, +\infty].$$

$$E[X] = \frac{n}{n-2} \text{ for } n \geq 3, \text{ Var}[X] = \frac{2n^2(m+n-4)}{m(n-2)^2(n-4)} \text{ for } n \geq 5.$$

$E[X] = \infty$  for  $n = 1, 2$ ,  $\text{Var}[X]$  does not exist or is infinite for  $n = 1, 2, 3, 4$ .

### IV.3. The Multivariate Normal Distribution. (Bierens, Sections 5.2–5.3, 5.5)

DEFINITION: The r.v.  $X$  with values in  $\mathbb{R}^k$  has a *multivariate normal distribution*  $\mathcal{N}(\mu, \Sigma)$  where  $\mu \in \mathbb{R}^k$  and  $\Sigma$  is positive definite matrix iff its density w.r.t. Lebesgue measure is

$$f_X(x) = \frac{1}{(\sqrt{2\pi})^n \sqrt{\det(\Sigma)}} \exp \left[ -\frac{1}{2}(x - \mu)' \Sigma^{-1}(x - \mu) \right] \quad \text{for } x \in \mathbb{R}^k.$$

PROPERTIES:

- (a)  $E[X] = \mu$ ,  $\text{Var}[X] = \Sigma$ ,  $\phi_X(t) = \exp(it'\mu - t'\Sigma t/2)$ .
- (b) If  $Y = A + BX$  where  $X \sim \mathcal{N}(\mu, \Sigma)$ , then  $Y \sim \mathcal{N}(A + B\mu, B\Sigma B')$ .
- (c) If  $X \sim \mathcal{N}(\mu, \Sigma)$ , then its components are independent iff  $\Sigma$  is diagonal.
- (d) If  $Y \in \mathbb{R}$  and  $X \in \mathbb{R}^k$  are (jointly) normal distributed with  $E[Y] = \mu_Y$ ,  $E[X] = \mu_X$ ,  $\text{Var}[Y] = \sigma_Y^2$ ,  $\text{Var}[X] = \Sigma_{XX}$  nonsingular, and  $E[XY] = \Sigma_{XY}$ , then  $P_{Y|X}(\cdot|x)$  is normal with  $E[Y|X] = \mu_Y + (X - \mu_X)' \Sigma_{XX}^{-1} \Sigma_{XY}$  and  $\text{Var}[Y|X] = \sigma_Y^2 - \Sigma_{YX} \Sigma_{XX}^{-1} \Sigma_{XY}$  with  $\Sigma'_{YX} = \Sigma_{XY}$ .
- (e) If  $X \sim \mathcal{N}(0, \Sigma)$ , then  $X' \Sigma^{-1} X \sim \chi_k^2$ .



## V. Modes of Convergence (Bierens, Sections 6.2–6.3, 6.5–6.7)

DEFINITION: Let  $X$  and  $\{X_n\}_{n=1}^{\infty}$  be random variables on  $(\Omega, \mathcal{F}, P)$ .

- (i)  $X_n$  converges to  $X$  almost surely, i.e.  $X_n \xrightarrow{a.s.} X$ , iff  $P[\lim_{n \rightarrow \infty} X_n = X] = 1$ ,
- (ii)  $X_n$  converges to  $X$  in probability, i.e.  $X_n \xrightarrow{p} X$ , iff  $\forall \epsilon > 0 \lim_{n \rightarrow \infty} P[|X_n - X| \leq \epsilon] = 1$ ,
- (iii)  $X_n$  converges to  $X$  in distribution, i.e.  $X_n \xrightarrow{d} X$ , iff  $\lim_{n \rightarrow \infty} F_{X_n}(x) = F_X(x)$   
for all continuity points  $x$  of  $F_X(\cdot)$ .

REMARKS:

- Alternative definitions:

- (i)  $X_n \xrightarrow{a.s.} X$  iff  $\forall \epsilon > 0 \lim_{n \rightarrow \infty} P[\sup_{m \geq n} |X_m - X| \leq \epsilon] = 1$ ,
- (iii)  $X_n \xrightarrow{d} X$  iff  $\lim_{n \rightarrow \infty} E[g(X_n)] = E[g(X)]$  for all bounded continuous  $g(\cdot) : \mathbb{R} \rightarrow \mathbb{R}$ .

- The multivariate case:  $X \in \mathbb{R}^k$  and  $X_n \in \mathbb{R}^k$

- (i)-(ii)  $X_n \xrightarrow{a.s./p} X$  iff  $X_n \xrightarrow{a.s./p} X$  componentwise,
- (iii)  $X_n \xrightarrow{d} X$  iff  $\lambda' X_n \xrightarrow{d} \lambda' X$  for all  $\lambda \in \mathbb{R}^k$  iff  $\lim_{n \rightarrow \infty} \phi_{X_n}(t) = \phi_X(t)$  for all  $t \in \mathbb{R}^k$ .

PROPERTIES:

(a)  $X_n \xrightarrow{a.s.} X \Rightarrow X_n \xrightarrow{p} X \Rightarrow X_n \xrightarrow{d} X$ .

The reverse is not true except:  $X_n \xrightarrow{p} c \Leftrightarrow X_n \xrightarrow{d} c$  where  $c$  is a constant.

(b) *Continuous Mapping (Slutzky) Theorem*: Let  $g(\cdot) : \mathbb{R}^k \rightarrow \mathbb{R}^\ell$  be continuous ( $P_X$ -a.s.).

If  $X_n \xrightarrow{a.s./p/d} X$ , then  $g(X_n) \xrightarrow{a.s./p/d} g(X)$ .

Examples: Let  $X = (X'_1, X'_2)'$  and  $X_n = (X'_{1n}, X'_{2n})'$ . If  $X_n \xrightarrow{a.s./p/d} X$ , then

$X_{1n} + X_{2n} \xrightarrow{a.s./p/d} X_1 + X_2$ ,  $X_{1n} X_{2n} \xrightarrow{a.s./p/d} X_1 X_2$ ,  $X_{1n}/X_{2n} \xrightarrow{a.s./p/d} X_1/X_2$  when  $P[X_2=0]=0$ .

BASIC LLN AND CLT: Let  $X_i, i = 1, 2, \dots$  be i.i.d. and  $\bar{X}_n \equiv \frac{1}{n} \sum_{i=1}^n X_i$ .

- Strong LLN (Kolmogorov): If  $E[X_i] \equiv \mu < \infty$ , then  $\bar{X}_n \xrightarrow{a.s.} \mu$ .

- CLT: If  $E[X_i] \equiv \mu < \infty$  and  $\text{Var}[X_i] \equiv \Sigma < \infty$  nonsingular, then  $\sqrt{n} \Sigma^{-1/2}(\bar{X}_n - \mu) \xrightarrow{d} \mathcal{N}(0, I)$ .

REMARKS: Extensions to non i.i.d  $X_i$ . See Bierens, Sections 7.2-7.3, 7.5. Let  $\mu_i \equiv E[X_i]$ .

- SLLNs:  $\frac{1}{n} \sum_{i=1}^n (X_i - \mu_i) \xrightarrow{a.s.} 0 \Rightarrow$  WLLNs:  $\frac{1}{n} \sum_{i=1}^n (X_i - \mu_i) \xrightarrow{p} 0 \left( \Leftrightarrow \frac{1}{n} \sum_{i=1}^n (X_i - \mu_i) \xrightarrow{d} 0 \right)$ .

- CLTs: When  $\dim X_i = 1$ ,  $\frac{\sum_{i=1}^n (X_i - E[X_i])}{\sqrt{\text{Var}(\sum_{i=1}^n X_i)}} \xrightarrow{d} \mathcal{N}(0, 1)$ .

## THE LINEAR REGRESSION MODEL (Bierens, Section 5.7)

- Data:  $(Y_i, X_i) \in \mathbb{R} \times \mathbb{R}^k, i = 1, 2, \dots$  i.i.d. on  $(\Omega, \mathcal{F}, P_0)$ .
- Model:  $E[Y|X] = X'\beta_0$  or  $Y_i = X_i'\beta_0 + \epsilon_i$  where  $E[\epsilon_i|X_i] = 0$ .
- *OLS estimator*:  $\hat{\theta}_{OLS} = \arg \min_{\beta} (1/n) \sum_{i=1}^n (Y_i - X_i'\beta)^2$ . Provided  $\mathbf{X}$  is *full-column rank*

$$\hat{\beta}_{OLS} = \left( \frac{1}{n} \sum_{i=1}^n X_i X_i' \right)^{-1} \left( \frac{1}{n} \sum_{i=1}^n X_i Y_i \right) = (\mathbf{X}'\mathbf{X})^{-1} \mathbf{X}'\mathbf{Y}$$

- Properties of  $\hat{\beta}_{OLS}$ : Assume  $E[XX'] \equiv \Sigma_{XX}$  is p.d., and  $0 < \sigma_{\epsilon}^2(X) \equiv E[\epsilon^2|X] < \infty$
- (a) Finite sample property:  $E[\hat{\beta}_{OLS}] = \beta_0$ , i.e.  $\hat{\beta}_{OLS}$  *unbiased estimator* of  $\beta_0$ , etc.
- (b) Asymptotic properties:  $\hat{\beta}_{OLS} \xrightarrow{a.s.} \beta_0$  and  $\sqrt{n}(\hat{\beta}_{OLS} - \beta_0) \xrightarrow{d} \mathcal{N}(0, \Sigma_{XX}^{-1} E[\sigma_{\epsilon}^2(X) XX'] \Sigma_{XX}^{-1})$ .
- (c) Approximation:  $\hat{\beta}_{OLS} \approx \mathcal{N}(\beta_0, (\mathbf{X}'\mathbf{X})^{-1} [\sum_{i=1}^n e_i^2 X_i X_i'] (\mathbf{X}'\mathbf{X})^{-1})$  where  $e_i \equiv Y_i - X_i'\hat{\beta}_{OLS}$ .  
 $\rightarrow (\hat{\beta}_j - \beta_{j0})/\hat{\sigma}_j \approx \mathcal{N}(0, 1)$  where  $\hat{\sigma}_j^2 \equiv [(\mathbf{X}'\mathbf{X})^{-1} [\sum_{i=1}^n e_i^2 X_i X_i'] (\mathbf{X}'\mathbf{X})^{-1}]_{jj}$  for  $j = 1, \dots, k$ .  
 $\hat{\sigma}_j$  is called White (1980) *Heteroscedasticity Robust Standard Error*.
- (d) Homoscedastic case  $\sigma_{\epsilon}^2(\cdot) = \sigma_{\epsilon 0}^2$ :  $\hat{\beta}_{OLS} \approx \mathcal{N}(\beta_0, \tilde{\sigma}_{\epsilon}^2 (\mathbf{X}'\mathbf{X})^{-1})$  where  $\tilde{\sigma}_{\epsilon}^2 \equiv \frac{1}{n-k} \sum_{i=1}^n e_i^2$ .  
 $\rightarrow$  When  $\epsilon_i \sim \mathcal{N}(0, \sigma_{\epsilon 0}^2)$ , then  $(\hat{\beta}_j - \beta_{j0})/\tilde{\sigma}_j \sim t_{n-k}$  where  $\tilde{\sigma}_j^2 \equiv \tilde{\sigma}_{\epsilon}^2 [(\mathbf{X}'\mathbf{X})^{-1}]_{jj}$ .

## VI. M-Estimation (Bierens, Sections 6.4, 6.9)

**DEFINITION:** An *extremum* or *M-estimator*  $\hat{\theta}_n \in \Theta \subseteq \mathbb{R}^k$  is a solution of  $\max_{\theta \in \Theta} Q_n(\theta)$  or  $\min_{\theta \in \Theta} \tilde{Q}_n(\theta)$ , where  $Q_n(\theta) = -\tilde{Q}_n(\theta)$  is a (measurable) function of the data.

**EXAMPLES:**

- OLS estimator  $\hat{\beta}_{OLS} \equiv \arg \min_{\beta \in \mathbb{R}^k} \tilde{Q}_n(\beta) = \frac{1}{n} \sum_{i=1}^n (Y_i - X_i' \beta)^2$ .
- NLLS estimator  $\hat{\theta}_{NLLS} \equiv \arg \min_{\theta \in \mathbb{R}^k} \tilde{Q}_n(\beta) = \frac{1}{n} \sum_{i=1}^n [Y_i - m(X_i; \theta)]^2$ .
- IV estimator  $\hat{\beta}_{IV} \equiv \arg \min_{\beta \in \mathbb{R}^k} \tilde{Q}_n(\beta) = \left\| \frac{1}{n} \sum_{i=1}^n Z_i (Y_i - X_i' \beta) \right\|_{A_n}$ .
- GMM estimator  $\hat{\theta}_{GMM} \equiv \arg \min_{\theta \in \Theta} \tilde{Q}_n(\beta) = \left\| \frac{1}{n} \sum_{i=1}^n h(Y_i, X_i, Z_i; \theta) \right\|_{A_n}$ .
- ML estimator  $\hat{\theta}_{ML} \equiv \arg \max_{\theta \in \Theta} Q_n(\beta) = \frac{1}{n} \sum_{i=1}^n \log f(Y_i | X_i; \theta)$ .

**Master Consistency Theorem:** For a nonstochastic function  $Q_\infty(\cdot)$  on  $\Theta$  and  $\theta_* \in \Theta$ . If

- (i) *Separation:*  $\sup_{\theta \in \Theta \cap \mathcal{N}^c} Q_\infty(\theta) < Q_\infty(\theta_*)$ , for any neighborhood  $\mathcal{N}$  of  $\theta_*$ ,
- (ii) *Uniform convergence:*  $\sup_{\theta \in \Theta} |Q_n(\theta) - Q_\infty(\theta)| \xrightarrow{a.s./p} 0$ ,

then  $\hat{\theta}_n \xrightarrow{a.s./p} \theta_*$  and  $\theta_*$  is called the *pseudo-true parameter value*.

*Proof of Strong Consistency:* Consider only the  $\omega$  for which  $\lim_{n \rightarrow \infty} \sup_{\theta \in \Theta} |Q_n(\theta) - Q_\infty(\theta)| = 0$ .

Thus for  $\epsilon > 0$  and  $n$  sufficiently large,  $Q_\infty(\theta) - \epsilon/3 < Q_n(\theta) < Q_\infty(\theta) + \epsilon/3$  for all  $\theta \in \Theta$ .

Hence, (a)  $Q_\infty(\theta_*) - \epsilon/3 < Q_n(\theta_*)$  and (b)  $Q_n(\hat{\theta}_n) < Q_\infty(\hat{\theta}_n) + \epsilon/3$ .

Because  $Q_n(\hat{\theta}_n) \geq Q_n(\theta)$  for all  $\theta \in \Theta$ , then  $Q_n(\hat{\theta}_n) > Q_\infty(\theta_*) - \epsilon/3$  by (a). Hence, by (b)

$$Q_\infty(\hat{\theta}_n) > Q_n(\hat{\theta}_n) - \epsilon/3 > Q_\infty(\theta_*) - 2\epsilon/3 > Q_\infty(\theta_*) - \epsilon.$$

Let  $\mathcal{N}$  be an arbitrary neighborhood of  $\theta_*$  and set  $\epsilon = Q_\infty(\theta_*) - \sup_{\theta \in \Theta \cap \mathcal{N}^c} Q_\infty(\theta)$ . Thus,

$$Q_\infty(\hat{\theta}_n) > \sup_{\theta \in \Theta \cap \mathcal{N}^c} Q_\infty(\theta) \Rightarrow \hat{\theta}_n \in \Theta \cap \mathcal{N}.$$

REMARKS:

- Condition (i)  $\Rightarrow \theta_*$  is the unique maximizer of  $Q_\infty(\cdot)$  on  $\Theta$ . See Lemma below for the reverse.
- Condition (ii) is often proved using a uniform SLLN. See below.
- $\theta_* \neq \theta_0$ . If condition (i) is satisfied by  $\theta_0$ , then  $\theta_* = \theta_0$ .
- Neither  $Q_n(\cdot)$  nor  $Q_\infty(\cdot)$  need to be continuous on  $\Theta$ .
- Generalization to  $\Theta \subseteq \mathcal{G}$  with a metric  $d$  on  $\mathcal{G}$ .

LEMMA: Suppose (i)  $\Theta$  is compact, (ii)  $Q_\infty(\cdot)$  is continuous on  $\Theta$ , and (iii) *Identification*:  $\theta_*$  is the unique maximizer of  $Q_\infty(\cdot)$  on  $\Theta$ . Then condition (i) in the basic consistency theorem holds.

*Proof of Lemma:* By (i)  $\Theta \cap \mathcal{N}^c$  is compact for any neighborhood  $\mathcal{N}$  of  $\theta_*$ . Thus, by (ii)  $\sup_{\theta \in \Theta \cap \mathcal{N}^c} Q_\infty(\theta) = Q_\infty(\bar{\theta})$  for some  $\bar{\theta} \in \Theta$ . By (iii)  $Q_\infty(\bar{\theta}) < Q_\infty(\theta_*)$ .

EXAMPLE:  $\tilde{Q}_\infty(\beta) \equiv E[Y - X'\beta]^2$  and  $\beta_* \equiv \arg \min_{\beta \in B} \tilde{Q}_\infty(\beta)$ . If  $E[Y^2] < \infty$  and  $E[XX'] < \infty$  nonsingular, then  $\tilde{Q}_\infty(\cdot)$  is continuous on  $B$  and  $\beta_* = \arg \min_{\beta \in B} E[E(Y|X) - X'\beta]^2$ .

$\rightarrow \beta_* = (E[XX'])^{-1}E[XY]$  if the latter  $\in B \Rightarrow$  If  $E[Y|X] = X'\beta_0$  with  $\beta_0 \in B$ , then  $\beta_* = \beta_0$ .

BASIC UNIFORM SLLN: Let  $X_1, X_2, \dots$  be i.i.d. r.v.s in  $\mathbb{R}^p$  and  $g(\cdot; \cdot): \mathbb{R}^p \times \Theta \rightarrow \mathbb{R}$  be  $\mathbb{R}^p/\mathbb{R}$ -measurable and continuous on compact  $\Theta \subset \mathbb{R}^k$ . If  $\sup_{\theta \in \Theta} |g(\cdot; \theta)| \leq M(\cdot)$  with  $E[M(X_i)] < \infty$ , then  $\sup_{\theta \in \Theta} \left| \frac{1}{n} \sum_{i=1}^n \{g(X_i; \theta) - E[g(X_i; \theta)]\} \right| \xrightarrow{a.s.} 0$  and  $E[g(X_i; \cdot)]$  is continuous on  $\Theta$ .

REMARKS:

- Extensions to non i.i.d  $X_i$ . See Bierens, Section 7.4.
- Extensions to *empirical processes*:  $\sup_{g \in \mathcal{G}} \left| \frac{1}{n} \sum_{i=1}^n \{g(X_i) - E[g(X_i)]\} \right| \xrightarrow{a.s.} 0$ .

EXAMPLE (CONTINUED):  $\tilde{Q}_n(\beta) \equiv \frac{1}{n} \sum_{i=1}^n (Y_i - X_i' \beta)^2$  with  $B$  compact. We have  $\tilde{g}(y, x; \beta) \equiv (y - x' \beta)^2 \leq 2[y^2 + (x' \beta)^2] \leq 2(y^2 + \|x\|^2 \|\beta\|^2) \leq M(y, x) \equiv 2(y^2 + M_o^2 \|x\|^2)$ . If  $E[Y_i^2] < \infty$  and  $E[X_i X_i'] < \infty$ , then  $E[M(Y_i, X_i)] < \infty$ . Hence,  $\sup_{\beta \in B} |\tilde{Q}_n(\beta) - \tilde{Q}_\infty(\beta)| \xrightarrow{a.s./p} 0$ .

**Master Asymptotic Normality Theorem:** Let  $\hat{\theta}_n \xrightarrow{a.s./p} \theta_*$ . If

- (i)  $\theta_* \in \text{int}(\Theta)$ , and  $Q_n(\cdot)$  is twice continuously differentiable in a neighborhood  $\mathcal{N}$  of  $\theta_*$ ,
  - (ii)  $\sup_{\theta \in \mathcal{N}} \|\partial^2 Q_n(\theta) / \partial \theta \partial \theta' - H(\theta)\| \xrightarrow{a.s./p} 0$  where  $H(\cdot)$  is continuous and nonsingular at  $\theta_*$ ,
  - (iii)  $\sqrt{n} \partial Q_n(\theta_*) / \partial \theta \xrightarrow{d} \mathcal{N}(0, \Sigma)$  for  $\Sigma$  nonsingular,
- then  $\sqrt{n}(\hat{\theta}_n - \theta_*) \xrightarrow{d} \mathcal{N}(0, H^{-1} \Sigma H^{-1})$  where  $H \equiv H(\theta_*)$ .

*Proof of Asymptotic Normality:* By (i)  $\partial Q_n(\hat{\theta}_n) / \partial \theta = 0$  where  $\hat{\theta}_n \in \mathcal{N}$  with probability one or approaching one. Taking a second-order Taylor expansion at  $\theta_*$  on the neighborhood  $\mathcal{N}$  gives

$$0 = \sqrt{n} \frac{\partial Q_n(\theta_*)}{\partial \theta} + \frac{\partial^2 Q_n(\bar{\theta}_n)}{\partial \theta \partial \theta'} \sqrt{n}(\hat{\theta}_n - \theta_*) \quad \text{where } \bar{\theta}_n \in [\theta_*, \hat{\theta}_n] \text{ (componentwise)}$$

By (ii)  $\partial^2 Q_n(\bar{\theta}_n) / \partial \theta \partial \theta' \xrightarrow{a.s./p} H = H(\theta_*)$  since  $\bar{\theta}_n \xrightarrow{a.s./p} \theta_*$ . By (iii) and the continuous mapping theorem, the result follows since  $H$  is nonsingular.

REMARK: Condition (ii) by USLLN while condition (iii) from CLT at  $\theta_*$ . See NLLS and MLE.

EXAMPLE (CONTINUED):  $\frac{\partial^2 \tilde{Q}_n(\beta)}{\partial \beta \partial \beta'} = \frac{2}{n} \sum_{i=1}^n X_i X_i' \xrightarrow{a.s./p} 2\Sigma_{XX}$ ,  $\sqrt{n} \frac{\partial \tilde{Q}_n(\beta_*)}{\partial \beta} = \frac{-2}{\sqrt{n}} \sum_{i=1}^n X_i (Y_i - X_i' \beta_*) \xrightarrow{d}$

$\mathcal{N}(0, 4E[\sigma_{\epsilon_*}^2(X)XX'])$ , where  $\sigma_{\epsilon_*}(X) \equiv E[\epsilon_*^2|X]$  and  $\epsilon_* \equiv Y - X'\beta_*$ . Thus,

$\sqrt{n}(\hat{\beta}_{OLS} - \beta_*) \xrightarrow{d} \mathcal{N}(0, \Sigma_{XX}^{-1} E[\sigma_{\epsilon_*}^2(X)XX'] \Sigma_{XX}^{-1})$  same result as before with  $\beta_*$  instead of  $\beta_0$ .

→ Approximation:  $\hat{\beta}_{OLS} \approx \mathcal{N}(\beta_*, (\mathbf{X}'\mathbf{X})^{-1} [\sum_{i=1}^n e_i^2 X_i X_i'] (\mathbf{X}'\mathbf{X})^{-1})$  where  $e_i \equiv Y_i - X_i' \hat{\beta}_{OLS}$ .

THEOREM (DELTA METHOD): Let  $\sqrt{n}(X_n - c) \xrightarrow{d} \mathcal{N}_p(0, \Omega)$  and  $g(\cdot) : \mathbb{R}^p \rightarrow \mathbb{R}^m$  be continuously differentiable on a neighborhood of  $c \in \mathbb{R}^p$ . Then  $\sqrt{n}[g(X_n) - g(c)] \xrightarrow{d} \mathcal{N}_m\left(0, \frac{\partial g(c)}{\partial c'} \Omega \frac{\partial g(c)'}{\partial c}\right)$ .

REMARKS:

- Different from continuous mapping theorem.

-  $n[g(X_n) - g(c)]' \left[ \frac{\partial g(X_n)}{\partial c'} \hat{\Omega} \frac{\partial g(X_n)'}{\partial c} \right]^{-1} [g(X_n) - g(c)] \xrightarrow{d} \chi_p^2$  where  $\hat{\Omega} \xrightarrow{a.s./p} \Omega$  if  $\frac{\partial g(c)}{\partial c'} \Omega \frac{\partial g(c)'}{\partial c}$  nonsingular.

→ Application to testing  $H_0 : g(\theta_*) = 0$  vs.  $H_A : g(\theta_*) \neq 0$ . If  $\sqrt{n}(\hat{\theta}_n - \theta_*) \xrightarrow{d} \mathcal{N}_k(0, \Omega)$ ,

then  $n g(\hat{\theta}_n)' \left[ \frac{\partial g(\hat{\theta}_n)}{\partial \theta'} \hat{\Omega} \frac{\partial g(\hat{\theta}_n)'}{\partial \theta} \right]^{-1} g(\hat{\theta}_n) \xrightarrow{d} \chi_p^2$  under  $H_0$  and  $\xrightarrow{a.s./p} +\infty$  under  $H_1$

where  $\hat{\Omega} \equiv \hat{H}^{-1} \hat{\Sigma} \hat{H}^{-1} \xrightarrow{a.s./p} \Omega = H^{-1} \Sigma H^{-1}$  and  $\hat{H} = \frac{\partial^2 Q_n(\hat{\theta}_n)}{\partial \theta \partial \theta'}$ . See below for  $\hat{\Sigma}$ .



## VII. NLLS and ML Estimation (Bierens, Sections 6.4, 8.1-8.2, 8.4)

**VII.1. Non-Linear Least Squares:**  $\hat{\theta}_{NLLS} \equiv \arg \min_{\theta \in \Theta} \tilde{Q}_n(\theta) = \frac{1}{n} \sum_{i=1}^n [Y_i - m(X_i; \theta)]^2$

**THEOREM (NLLS):** Let  $(Y_1, X_1), (Y_2, X_2), \dots$  be i.i.d. r.v.s in  $\mathbb{R} \times \mathbb{R}^p$  and  $m(\cdot; \cdot) : \mathbb{R}^p \times \Theta \rightarrow \mathbb{R}$  be measurable on  $\mathbb{R}^p$  and continuous on  $\Theta \subset \mathbb{R}^k$ . Suppose that  $E[Y^2] < \infty$  and  $\theta_*$  uniquely solves  $\min_{\theta \in \Theta} Q_\infty = E[Y - m(X; \theta)]^2$ .

- (i) If  $\Theta$  is compact and  $\sup_{\theta \in \Theta} |m(\cdot; \theta)| \leq M(\cdot)$  with  $E[M^2(X)] < \infty$ , then  $\hat{\theta}_{NLLS} \xrightarrow{a.s./p} \theta_*$ .
- (ii) If in addition  $\theta_* \in \text{int}(\Theta)$  with  $m(\cdot, \cdot)$  twice cont. differentiable on a neighborhood  $\mathcal{N}$  of  $\theta_*$ ,  
 $\sup_{\theta \in \mathcal{N}} \left\| \frac{\partial m(\cdot; \theta)}{\partial \theta} \right\| \leq M_1(\cdot)$  with  $E[M_1^2(X)] < \infty$ ,  $\sup_{\theta \in \mathcal{N}} \left\| \frac{\partial^2 m(\cdot; \theta)}{\partial \theta \partial \theta'} \right\| \leq M_2(\cdot)$  with  $E[M_2^2(X)] < \infty$ ,  
then  $\sqrt{n}(\hat{\theta}_{NLLS} - \theta_*) \xrightarrow{d} \mathcal{N}(0, H^{-1} \Sigma H^{-1})$  provided  $H = 2E \left[ \epsilon_* \frac{\partial^2 m(X; \theta_*)}{\partial \theta \partial \theta'} - \frac{\partial m(X; \theta_*)}{\partial \theta} \frac{\partial m(X; \theta_*)}{\partial \theta'} \right]$   
is nonsingular and  $\Sigma = 4E \left[ \sigma_{\epsilon_*}^2(X) \frac{\partial m(X; \theta_*)}{\partial \theta} \frac{\partial m(X; \theta_*)}{\partial \theta'} \right]$  with  $\sigma_{\epsilon_*}^2(X) = E[\epsilon_*^2 | X]$  and  $\epsilon_* = Y - m(X; \theta_*)$ .

**REMARKS:** Let  $e_i \equiv Y_i - m(X_i; \hat{\theta}_{NLLS})$ . Then  $\hat{\Sigma} = \frac{4}{n} \sum_{i=1}^n e_i^2 \frac{\partial m(X_i; \hat{\theta}_{NLLS})}{\partial \theta} \frac{\partial m(X_i; \hat{\theta}_{NLLS})}{\partial \theta'} \xrightarrow{a.s./p} \Sigma$   
and  $\hat{H} = \frac{2}{n} \sum_{i=1}^n \left[ e_i \frac{\partial^2 m(X; \hat{\theta}_{NLLS})}{\partial \theta \partial \theta'} - \frac{\partial m(X; \hat{\theta}_{NLLS})}{\partial \theta} \frac{\partial m(X; \hat{\theta}_{NLLS})}{\partial \theta'} \right] \xrightarrow{a.s./p} H$ .

- If  $E[Y|X] = m(X; \theta_0)$  for  $\theta_0 \in \Theta$ , then  $\theta_* = \theta_0$ . If, in addition  $\sigma_{\epsilon_*}^2(X) = \sigma_0^2$ , then

$$H = -2E \left[ \frac{\partial m(X; \theta_0)}{\partial \theta} \frac{\partial m(X; \theta_0)}{\partial \theta'} \right], \quad \Sigma = 4\sigma_0^2 E \left[ \frac{\partial m(X; \theta_0)}{\partial \theta} \frac{\partial m(X; \theta_0)}{\partial \theta'} \right], \quad H^{-1} \Sigma H^{-1} = \sigma_0^2 \left( E \left[ \frac{\partial m(X; \theta_0)}{\partial \theta} \frac{\partial m(X; \theta_0)}{\partial \theta'} \right] \right)^{-1}$$

**VII.2. Maximum Likelihood:**  $\hat{\theta}_{ML} \equiv \arg \max_{\theta \in \Theta} Q_n(\beta) = \frac{1}{n} \sum_{i=1}^n \log f(Y_i|X_i; \theta)$

**THEOREM (ML):** Let  $(Y_1, X_1), (Y_2, X_2), \dots$  be i.i.d. r.v.s in  $\mathbb{R}^m \times \mathbb{R}^p$ . Let the *model* be a family  $\{f(\cdot|\cdot; \theta) : \theta \in \Theta\}$  of conditional densities w.r.t. a measure  $\mu$  on  $\mathbb{R}^m$  and continuous on  $\Theta \subset \mathbb{R}^k$ . Suppose that  $\theta_*$  uniquely solves  $\max_{\theta \in \Theta} Q_\infty = E[\log f(Y|X; \theta)]$ .

- (i) If  $\Theta$  is compact and  $\sup_{\theta \in \Theta} |\log f(\cdot|\cdot; \theta)| \leq M(\cdot, \cdot)$  with  $E[M(Y, X)] < \infty$ , then  $\hat{\theta}_{ML} \xrightarrow{a.s./p} \theta_*$ .
- (ii) If in addition  $\theta_* \in \text{int}(\Theta)$  with  $f(\cdot|\cdot; \cdot)$  twice continuously differentiable on a neighborhood  $\mathcal{N}$  of  $\theta_*$ ,  $\sup_{\theta \in \mathcal{N}} \left\| \frac{\partial^2 \log f(\cdot|\cdot; \theta)}{\partial \theta \partial \theta'} \right\| \leq M_2(\cdot, \cdot)$ ,  $E[M_2(Y, X)] < \infty$  and  $E \left[ \frac{\partial^2 \log f(Y|X; \theta_*)}{\partial \theta \partial \theta'} \right]$  nonsingular, then

$$\sqrt{n}(\hat{\theta}_{ML} - \theta_*) \xrightarrow{d} \mathcal{N}(0, A^{-1}BA^{-1})$$

$$\text{where } A = A(\theta_*) \equiv E \left[ \frac{\partial^2 \log f(Y|X; \theta_*)}{\partial \theta \partial \theta'} \right] \text{ and } B = B(\theta_*) \equiv E \left[ \frac{\partial \log f(Y|X; \theta_*)}{\partial \theta} \frac{\partial \log f(Y|X; \theta_*)}{\partial \theta'} \right].$$

**REMARKS:**

- If  $\sup_{\theta \in \mathcal{N}} \left\| \frac{\partial \log f(\cdot|\cdot; \theta)}{\partial \theta} \right\| \leq M_1(\cdot)$  with  $E[M_1(X)] < \infty$ , then

$$\hat{A} = \frac{1}{n} \sum_{i=1}^n \frac{\partial^2 \log f(Y_i|X_i; \hat{\theta}_{ML})}{\partial \theta \partial \theta'} \xrightarrow{a.s./p} A \text{ and } \hat{B} = \frac{1}{n} \sum_{i=1}^n \frac{\partial \log f(Y_i|X_i; \hat{\theta}_{ML})}{\partial \theta} \frac{\partial \log f(Y_i|X_i; \hat{\theta}_{ML})}{\partial \theta'} \xrightarrow{a.s./p} B.$$

- *Kullback-Leibler Divergence:*  $\theta_* = \arg \min_{\theta \in \Theta} KLIC[f_{Y|X}(\cdot|\cdot); f(\cdot|\cdot; \theta)] \equiv E \left[ \log \frac{f_{Y|X}(Y|X)}{f(Y|X; \theta)} \right] \geq 0.$

DEFINITION:  $\{f(\cdot|\cdot; \theta) : \theta \in \Theta\}$  is *correctly specified* iff  $f_{Y|X}(\cdot|\cdot) = f(\cdot|\cdot; \theta_0)$  a.s. for  $\theta_0 \in \Theta$ .

It is *one-to-one parameterized* iff  $f(\cdot|\cdot; \theta_1) = f(\cdot|\cdot; \theta_2)$  a.s.  $\Leftrightarrow \theta_1 = \theta_2$  for any  $\theta_1, \theta_2 \in \Theta$ .

LEMMA: If  $\{f(\cdot|\cdot; \theta) : \theta \in \Theta\}$  is correctly specified and one-to-one parameterized, then  $\theta_0$  uniquely solves  $\max_{\theta \in \Theta} E[\log f(Y|X; \theta)]$ .

*Proof of Lemma:* If  $f(y|x; \theta_0) > 0$ , then  $\log \frac{f(y|x; \theta)}{f(y|x; \theta_0)} \leq \frac{f(y|x; \theta)}{f(y|x; \theta_0)} - 1$  with equality iff  $\frac{f(y|x; \theta)}{f(y|x; \theta_0)} = 1$ . Thus  $E \left[ \log \frac{f(y|x; \theta)}{f(y|x; \theta_0)} \right] \leq E \left[ \frac{f(y|x; \theta)}{f(y|x; \theta_0)} \right] - 1 \leq \int_{\{f(y|x; \theta_0) > 0\}} f(y|x; \theta) d\mu(y) dF_X(x) - 1 \leq 0$ . Hence,  $\theta_0 = \arg \max_{\theta \in \Theta} E[\log f(Y|X; \theta)]$ . Moreover,  $E \left[ \log \frac{f(y|x; \theta)}{f(y|x; \theta_0)} \right] = 0 \Rightarrow E \left[ \log \frac{f(y|x; \theta)}{f(y|x; \theta_0)} - \frac{f(y|x; \theta)}{f(y|x; \theta_0)} + 1 \right] = 0 \Rightarrow \log \frac{f(y|x; \theta)}{f(y|x; \theta_0)} = \frac{f(y|x; \theta)}{f(y|x; \theta_0)} - 1$  a.s.  $\Rightarrow \frac{f(y|x; \theta)}{f(y|x; \theta_0)} = 1$  a.s.  $\Rightarrow \theta = \theta_0$ , i.e.  $\theta_0$  is the unique maximizer.

LEMMA (INFORMATION MATRIX EQUALITY): Assume that  $\{f(\cdot|\cdot; \theta) : \theta \in \Theta\}$  is correctly specified. If one can switch integration and differentiation, then  $A(\theta_0) + B(\theta_0) = 0$ .

*Proof of Lemma:* Differentiating twice  $\int f(y|x; \theta) d\mu(y) = 1$  w.r.t.  $\theta$  gives

$$\int \frac{\partial^2 \log f(y|x; \theta)}{\partial \theta \partial \theta'} f(y|x; \theta) d\mu(y) + \int \frac{\partial \log f(y|x; \theta)}{\partial \theta} \frac{\partial \log f(y|x; \theta)}{\partial \theta'} f(y|x; \theta) d\mu(y) = 0 \text{ for every } \theta \in \Theta.$$

Letting  $\theta = \theta_0$  in  $A(\theta)$  and  $B(\theta)$  gives the result because  $f_{Y|X}(\cdot|\cdot) = f(\cdot|\cdot; \theta_0)$ .

REMARKS:

- *Information Matrix Test*: White (1982)  $H_0 : A(\theta_*) + B(\theta_*) = 0$  vs.  $H_A : A(\theta_*) + B(\theta_*) \neq 0$ .
- If  $\{f(\cdot|\cdot; \theta) : \theta \in \Theta\}$  is correctly specified and one-to-one parameterized, then
  - $\rightarrow \sqrt{n}(\hat{\theta}_{ML} - \theta_0) \xrightarrow{d} \mathcal{N}(0, B^{-1}) = \mathcal{N}(0, -A^{-1}) \Rightarrow \hat{\theta}_{ML} \approx \mathcal{N}\left(\theta_0, -\left[\sum_{i=1}^n \frac{\partial^2 \log f(Y_i|X_i; \hat{\theta}_{ML})}{\partial \theta \partial \theta'}\right]^{-1}\right)$ .
  - $\rightarrow \hat{\theta}_{ML}$  is *asymptotically efficient* among consistent estimators  $\hat{\theta}_n$ , i.e.,  $\lim_{n \rightarrow \infty} \text{Var}[\sqrt{n}\hat{\theta}_n] \geq B^{-1}$ .
  - $B = E\left[\frac{\partial \log f(Y|X; \theta_0)}{\partial \theta} \frac{\partial \log f(Y|X; \theta_0)}{\partial \theta'}\right]$  is the *Fisher Information Matrix* for one observation.

THEOREM (CRAMER-RAO): If  $\hat{\theta}_n$  is an unbiased estimator of  $\theta_0$ , then  $\text{Var}[\sqrt{n}\hat{\theta}_n] \geq B^{-1}$ .

*Proof of Cramer-Rao Bound*: We have  $\int g_n(y_1, x_1, \dots, y_n, x_n) \Pi_{i=1}^n f(y_i|x_i; \theta_0) d\mu(y_i) dF_X(x_i) = \theta_0$  for any  $\theta_0 \in \Theta$ , where  $\hat{\theta}_n = g_n(Y_1, X_1, \dots, Y_n, X_n)$ . Differentiating w.r.t.  $\theta_0$  gives  $\int g_n(y_1, x_1, \dots, y_n, x_n) \left[\sum_{i=1}^n \frac{\partial \log f(y_i|x_i; \theta_0)}{\partial \theta'}\right] \Pi_{i=1}^n f(y_i|x_i; \theta_0) d\mu(y_i) dF_X(x_i) = I$ , i.e.,  $E[\hat{\theta}_n S'_n] = I$  where  $S_n \equiv \sum_{i=1}^n \frac{\partial \log f(Y_i|X_i; \theta_0)}{\partial \theta}$ . But  $E[S_n] = 0$  by differentiating  $\int f(y_i|x_i; \theta_0) d\mu(y_i) dF_X(x_i) = 1$  w.r.t.  $\theta_0$ . Hence  $\text{Cov}[\hat{\theta}_n, S_n] = I$  and  $\text{Var}[S_n] = nB$ . Now,  $\text{Var}[\hat{\theta}_n] = \text{Var}[(nB)^{-1}S_n + \hat{\theta}_n - (nB)^{-1}S_n] = (nB)^{-1} + \text{Var}[\hat{\theta}_n - (nB)^{-1}S_n] \geq (nB)^{-1}$  since  $\text{Cov}[(nB)^{-1}S_n, \hat{\theta}_n - (nB)^{-1}S_n] = 0$ .

**That's All Folks !**