

Topological Spaces

Subspace Topology

Product Topology

Closed Sets and Limit Points

Hausdorff Spaces

Continuous Functions

Finite Product

Infinite Product Topology

Quotient Topology

Connectedness

Compactness

Fundamental Groups.

Topological Spaces.

Def. (Topology). Let X be a space. A topology on X is a set of subsets of X , \mathcal{T} , following the properties:

- (1). $\emptyset \in \mathcal{T}$.
- (2). $X \in \mathcal{T}$.
- (3). For any collection $\{T_\alpha\}_{\alpha \in I} \subset \mathcal{T}$, $\bigcup_{\alpha \in I} T_\alpha \in \mathcal{T}$.
- (4). For any $T_1, \dots, T_m \in \mathcal{T}$, $\bigcap_{i=1}^m T_i \in \mathcal{T}$.

A topology is a legitimate definition of which sets in X is open. \emptyset and X should be considered open; intersection of finitely many open sets is open; union of arbitrarily open sets is open.

Now we look at it first in metric space with focus on \mathbb{R}^n .

Def. (Metric Space). A metric space is a set X with a function $d: X \times X \rightarrow \mathbb{R}$ such that

- (1). $d(x, y) \geq 0 \quad \forall x, y \in X$; $d(x, y) = 0$ iff $x = y$.
- (2). $d(x, y) = d(y, x)$.
- (3). $d(x, z) \leq d(x, y) + d(y, z)$.

Def. (Open ball in metric space) $B_d(x, \varepsilon) = \{y \in X \mid d(x, y) < \varepsilon\}$.

Def. (d -Open). $U \subset X$ is d -open if for every $x \in U$, $\exists \varepsilon > 0$ such that $B_d(x, \varepsilon) \subset U$.

Proposition. (a). Open balls are d -open.

- (b). A set is d -open iff it is equal to a union of d -open balls.
- (c). Any union of d -open sets is d -open.
- (d). Any intersection of finitely many d -open sets is d -open.
- (e). Given (X, d_X) , (Y, d_Y) , $f: X \rightarrow Y$ is continuous in the ε - δ sense iff for every d_Y -open $U \subset Y$, $f^{-1}(U)$ is d_X -open in X .

Proof. (a). Let $B_d(x, \varepsilon)$ be an open ball. Given $x' \in B_d(x, \varepsilon)$, $x' \in B_d(x, \varepsilon - d(x, x')) \subset B_d(x, \varepsilon)$.

(b). Let A be d -open. Then, for every point $x \in A$, $\exists \varepsilon_x > 0$ s.t. $x \in B_d(x, \varepsilon_x) \subset A$.
Then $A = \bigcup_{x \in A} B_d(x, \varepsilon_x)$.

Let A be a union of d -open balls. Then, for each point, there is a d -open ball around it.

(c). Let A be a union of d -open sets. Then A is a union of d -open balls by (b).

Then, for every point $a \in A$, a is in some open balls in A , thus A is d -open.

(d). Let $A = \bigcap_{i=1}^n A_i$, then, for each $a \in A$, $a \in A_i \quad \forall i \{1, \dots, n\}$. A_i is d -open.

Then, for each A_i , we can find d -open ball $B_d(a, \varepsilon_i)$ s.t. $a \in B_d(a, \varepsilon_i)$.

Then $a \in \bigcap_{i=1}^n B_d(a, \varepsilon_i)$. $a \in B_d(a, \min\{\varepsilon_i\}) \subset \bigcap_{i=1}^n B_d(a, \varepsilon_i) \subset A$.

Thus A is d -open.

(e) (\Rightarrow): We now have $\forall \varepsilon > 0, \exists \delta > 0$, s.t. $d_X(x, y) < \delta \Rightarrow d_Y(f(x), f(y)) < \varepsilon$.

Let U be d_Y -open in Y . WTS $f^{-1}(U)$ is d_X -open in X .

Let $x \in f^{-1}(U)$. Then $f(x) \in U$. Since U is d_Y -open, $\exists \varepsilon > 0$, $B_{d_Y}(f(x), \varepsilon) \subset U$.

Then, $\exists \delta > 0$, so that for all $x'' \in X$, $d_X(x, x'') < \delta \Rightarrow d_Y(f(x), f(x'')) < \varepsilon$.

Thus, $x \in B_{d_X}(x, \delta)$ and $f(x) \in B_{d_Y}(f(x), \varepsilon) \subset U$. Then $B_{d_X}(x, \delta) \subset f^{-1}(U)$. $f^{-1}(U)$ is d_X -open.

(\Leftarrow) . Let $x \in X$ be such that $f(x) \in U$.

Then, $\exists \varepsilon > 0$ s.t. $f(x) \in B_{df}(f(x), \varepsilon) \subset U$.

Since $f^{-1}(B_{df}(f(x), \varepsilon))$ is open in X , $\exists \delta > 0$ and $B_{dx}(x, \delta)$ such that $f(B_{dx}(x, \delta)) \subset B_{df}(f(x), \varepsilon) \subset U$.

Therefore $d_X(x, x') < \delta \Rightarrow d_f(f(x), f(x')) < \varepsilon$. Continuous in ε - δ sense.

□

Define Euclidean metric in \mathbb{R}^n as

$$d(\vec{x}, \vec{y}) = \left(\sum_{i=1}^n (x_i - y_i)^2 \right)^{\frac{1}{2}}$$

Define p metric as

$$p(\vec{x}, \vec{y}) = \max_i |x_i - y_i|$$

We first prove that p is a proper metric.

① $p(\vec{x}, \vec{y}) \geq 0$ is trivial

② (\Rightarrow) : suppose that for some $x_i \neq y_i$, then $p > 0$. (\Leftarrow) is again trivial.

$$\text{③ } p(\vec{x}, \vec{y}) + p(\vec{y}, \vec{z}) = \max_i |x_i - y_i| + \max_i |y_i - z_i|$$

$$\geq \max_i |x_i - z_i|$$

$$\geq \max_i |x_i - z_i| \\ = p(\vec{x}, \vec{z}).$$

Define $B_p(x, r) = \{y \in \mathbb{R}^n \mid p(x, y) < r\}$, then

$$\begin{aligned} B_p(x, r) &= \{(y_1, \dots, y_n) \mid |y_i - x_i| < r \ \forall i\} \\ &= \prod_{i=1}^n \{y_i \in \mathbb{R} \mid |x_i - y_i| < r\}. \end{aligned}$$

Then we prove $p(\vec{x}, \vec{y}) \leq d(\vec{x}, \vec{y}) \leq \sqrt{n} p(\vec{x}, \vec{y})$.

$$\begin{aligned} d(\vec{x}, \vec{y}) &= \left(\sum_{i=1}^n (x_i - y_i)^2 \right)^{\frac{1}{2}} \\ &\geq (\max_i |x_i - y_i|^2)^{\frac{1}{2}} \\ &\geq \max_i |x_i - y_i| \\ &= p(\vec{x}, \vec{y}). \end{aligned}$$

$$\begin{aligned} d(\vec{x}, \vec{y}) &= \left(\sum_{i=1}^n (x_i - y_i)^2 \right)^{\frac{1}{2}} \\ &\leq \left(\sum_{i=1}^n (\max_i |x_i - y_i|)^2 \right)^{\frac{1}{2}} \\ &= \sqrt{n} p(\vec{x}, \vec{y}). \end{aligned}$$

Next we prove that $B(\vec{x}, r) \subset B_p(\vec{x}, r)$, $B_p(\vec{x}, \frac{r}{\sqrt{n}}) \subset B(\vec{x}, r)$.

$\forall \vec{y} \in B(\vec{x}, r)$, $p(\vec{x}, \vec{y}) \leq d(\vec{x}, \vec{y}) < r$, so $\vec{y} \in B_p(\vec{x}, r)$. $\forall y \in B_p(\vec{x}, \frac{r}{\sqrt{n}})$, $d(\vec{x}, \vec{y}) \leq \sqrt{n} p(\vec{x}, \vec{y}) < r$, so $B_p(\vec{x}, \frac{r}{\sqrt{n}}) \subset B(\vec{x}, r)$.

Next we prove that U is p -open iff it is open.

(\Leftarrow) : Since U is open, $\forall \vec{x} \in U$, $\exists r > 0$ s.t. $B(\vec{x}, r) \subset U$. Then $B_p(\vec{x}, \frac{r}{\sqrt{n}}) \subset B(\vec{x}, r) \subset U$.

(\Rightarrow) : Since U is p -open, $\forall \vec{x} \in U$, $\exists r > 0$ s.t. $B_p(\vec{x}, r) \subset U$. Then $B(\vec{x}, r) \subset B_p(\vec{x}, r) \subset U$.

Define another matrix for $X \subset \mathbb{R}^n$: $q(\vec{x}, \vec{y}) = \begin{cases} 0 & \text{if } \vec{x} = \vec{y} \\ 1 & \text{if } \vec{x} \neq \vec{y} \end{cases}$

Then,

$$B_q(\vec{x}, r) = \begin{cases} \{\vec{x}\} & \text{if } r=1 \\ X & \text{if } r>1 \end{cases}$$

Thus, every set is q -open.

Back to the definition of topology.

For examples:

- (1). If (X, d) is a metric space, then the set of all d -open sets in X is a topology.
- (2). Discrete topology: $\mathcal{T}_d = \{\text{all subsets of } X\}$. In (X, q) , this topology is induced from $q(\vec{x}, \vec{y}) = 1 \iff \vec{x} = \vec{y}$.
- (3). Indiscrete topology: $\mathcal{T}_i = \{\emptyset, X\}$.
- (4). Topologies on $X = \{a, b\}$: $\mathcal{T}_1 = \{\emptyset, \{a\}, \{b\}, X\}$

$$\mathcal{T}_2 = \{\emptyset, X\}$$

$$\mathcal{T}_3 = \{\emptyset, \{ab\}, X\}$$

$$\mathcal{T}_4 = \{\emptyset, \{b\}, X\}.$$

- (5). Let (X, γ) be a metric space. " U is open iff $\forall x \in U, \exists \varepsilon > 0$ s.t. $B_d(x, \varepsilon) \subset U$ ". This is the metric topology induced by metric d . A different metric may induce a different \mathcal{T} .

(d and ρ induce the same topology).

Not every topology comes from a metric. For example, if X is a finite set, then for many metrics on X , the induced topology is the discrete topology. This is because if all one-point sets are open, then all subsets are open. To verify, $\forall \vec{x} \in X$, take $r = \min\{d(x, y) | y \neq x\}$. Then $B_d(x, r) = \{x\} \subset \{x\}$.

Def. (Finite Complement Topology / Cofinite Topology). Let X be a set.

$$\mathcal{T}_f = \{U \subset X \mid X - U \text{ is finite or } U = \emptyset\}.$$

Proposition. \mathcal{T}_f is a topology.

Proof: (1) $\emptyset \in \mathcal{T}_f$ (2) $X = X - \emptyset \in \mathcal{T}_f$

(3). Let $\{U_\alpha\} \subset \mathcal{T}_f$ such that $X - U_\alpha$ is finite.

$$X - \left(\bigcup_{\alpha} U_\alpha \right) = \bigcap_{\alpha} (X - U_\alpha)$$

is also finite.

(4). Let $\{U_i\}_{i=1}^n \subset \mathcal{T}_f$ such that $X - U_i$ is finite.

$$X - \bigcap_{i=1}^n U_i = \bigcup_{i=1}^n (X - U_i)$$

is also finite. □

Def (Finer). Let X be a set. Suppose \mathcal{T} and \mathcal{T}' are topologies on X . \mathcal{T}' is finer than \mathcal{T} if $\mathcal{T}' \subset \mathcal{T}$. Strictly finer if $\mathcal{T}' \neq \mathcal{T}$.

Proposition. Let γ be the standard topology on \mathbb{R} . Let \mathcal{T}_f be the finite complement topology. Then $\mathcal{T}_f \not\subset \mathcal{T}$.

Proof: First we prove $\mathcal{T} \neq \mathcal{T}_f$. $(-1, 1) \in \mathcal{T}$, but $(-1, 1) \notin \mathcal{T}_f$ since $\mathbb{R} - (-1, 1)$ is infinite.

Let U be $\mathbb{R} - U = \{a_1, \dots, a_k\}$. Then, $U = \mathbb{R} - \{a_1, \dots, a_k\}$. For any $y \in U$, let

$$r := \{d(y, a_i) \mid i=1, \dots, k\}, \text{ then } B_\gamma(y, r) \subset U. \text{ Thus } U \in \mathcal{T}$$

Therefore $\mathcal{T}_f \not\subset \mathcal{T}$. □

Then, we revisit the def of a continuous function.

Def. (Continuous function). Let (X, τ) and (Y, ν) be topological spaces. A function $f: X \rightarrow Y$ is continuous if for every open set $U \in \nu$, $f^{-1}(U)$ is open in X .
(i.e. $\forall U \in \nu$, $f^{-1}(U) \in \tau$). preimage of open set is open.

Def. (Homeomorphism). f is called a homeomorphism if it is bijective and both f and f^{-1} are continuous.
 (X, τ) and (Y, ν) are homeomorphic if \exists a homeomorphism between them.

For example, $X = \{a, b\}$

$$\begin{aligned} \tau_3 &= \{\emptyset, \{a, b\}, \{a\}\} \\ \tau_4 &= \{\emptyset, \{a, b\}, \{b\}\} \end{aligned} \quad \text{incomparable}$$

define $f: X \rightarrow X$ by $f(a) = b$, $f(b) = a$, then $f^{-1}(\{b\}) = \{a\}$, so f is continuous.
 $g := f^{-1}$ then $g^{-1}(\{a\}) = \{b\}$, so g is continuous.

Basis for a topology.

Def. (Basis). Let X be a set, \mathcal{B} a collection of subsets of X . \mathcal{B} is a topological basis if
1. For every $x \in X$, $\exists B \in \mathcal{B}$ such that $x \in B$.
2. For any $B_1, B_2 \in \mathcal{B}$ and any $x \in B_1 \cap B_2$, $\exists B_3 \in \mathcal{B}$ such that $x \in B_3$, $B_3 \subset B_1 \cap B_2$.


Given a topological basis \mathcal{B} , we obtain a topology τ by declaring $T \in \tau$ iff T is a union of sets in \mathcal{B} (i.e. $\forall x \in T$, $\exists B \in \mathcal{B}$ s.t. $x \in B \subset T$).

Def. (Topology generated by \mathcal{B}). A subset T of X is said to be open in X ($T \in \tau$) if for every $x \in T$, $\exists B \in \mathcal{B}$ such that $x \in B \subset T$. (τ is generated by \mathcal{B}).

e.g. (X, d) be a metric space. $B(x, r) \cap B(y, s)$. Let $z \in B(x, r) \cap B(y, s)$.

Take $t = \min(r - d(x, z), s - d(z, y))$. If $w \in B_d(z, t)$, then

$$\begin{aligned} d(w, x) &\leq d(w, z) + d(z, x) \\ &< t + d(z, x) \\ &\leq r - d(x, z) + d(x, z) \\ &= r. \end{aligned}$$



$$\begin{aligned} d(w, y) &\leq d(w, z) + d(z, y) \\ &< t + d(z, y) \\ &\leq s - d(z, y) + d(z, y) \\ &= s. \end{aligned}$$

Therefore $B_d(z, t) \subset B(x, r) \cap B(y, s)$.

Theorem: τ is a topology.

pf. First, if U is empty set, then it satisfies the defining condition of openness.

Likewise, $X \in \tau$ since for all $x \in X$, $\exists B_x \in \mathcal{B}$ such that $x \in B_x \subset X$.

Then, let $\{T_\alpha\}_{\alpha \in J} \subset \gamma$. WTS: $T = \bigcup_{\alpha \in J} T_\alpha \in \gamma$.

Given $x \in T$, $\exists \alpha \in J$ such that $x \in T_\alpha$. Since $T_\alpha \in \gamma$, there is a basis element B such that $x \in B \subset T_\alpha$. Then $x \in B \subset T$, so that $T \in \gamma$ by definition.

Then, given $T_1 \dots T_k \in \gamma$, $x \in T_1 \cap \dots \cap T_k$. There exists $B_1 \subset T_1, \dots, B_k \subset T_k$ such that $x \in \bigcap_{i=1}^k B_i$. Since by definition $\forall x \in \bigcap_{i=1}^k B_i, \exists B' \in \mathcal{B}$ such that $x \in B'$, therefore $\exists B_{k+1} \subset \bigcap_{i=1}^k B_i$ such that $x \in B_{k+1} \subset B' \subset T_1 \cap \dots \cap T_k$. Therefore $T_1 \cap \dots \cap T_k \in \gamma$.

□

e.g. $X = \mathbb{R}$, $\mathcal{B} = \{\text{all open intervals } (a, b) \text{ where } a < b\}$.

γ is the standard topology.

$$(a, b) = B\left(\frac{a+b}{2}, \frac{b-a}{2}\right)$$

e.g. $X = \mathbb{R}$, $\mathcal{B} = \{(a, b) \mid a < b, a, b \in \mathbb{Q}\}$ gives the same topology as last one. Use the fact that \mathbb{Q} is dense in \mathbb{R} , i.e. $\forall x, y \in \mathbb{R}, x < y, \exists z \in \mathbb{Q}$ s.t. $x < z < y$.

On \mathbb{R}^n : $d(x, y) = \sqrt{\sum (x_i - y_i)^2}$; $\rho(x, y) = \max_i |x_i - y_i|$

Open sets generated by these two are the same notion.

Let γ be the topology from d , γ' from ρ .

Lemma: Suppose d, d' are both metrics on X . Let γ, γ' be the induced topologies.

Suppose $\forall x \in X, \forall \varepsilon > 0, \exists \delta > 0$ such that $B_{d'}(x, \delta) \subset B_d(x, \varepsilon)$, then $\gamma \subset \gamma'$.

Proof:

If U is open in γ , then for each $x \in U, \exists \varepsilon > 0$ such that $B_d(x, \varepsilon) \subset U$.

$\exists \delta > 0$ such that $B_{d'}(x, \delta) \subset B_d(x, \varepsilon) \subset U$, then U is also open in γ' .

q.e.d.

Theorem: Let X be a set. Let γ and γ' be topologies generated by bases \mathcal{B} and \mathcal{B}' .

TFAE.

1. $\gamma \subset \gamma'$

2. $\forall B \in \mathcal{B}$ is open in γ' .

3. $\forall B \in \mathcal{B}, \exists B' \in \mathcal{B}'$ s.t. $x \in B' \subset B$.

Proof: 3 \Rightarrow 1. Given $T \in \gamma$. WTS $T \in \gamma'$.

Let $x \in U$. Since \mathcal{B} generates γ , $\exists B \in \mathcal{B}$ such that $x \in B \subset U$.

by (3): $\exists B' \in \mathcal{B}', x \in B' \subset B$. Then $x \in B' \subset U$. Therefore $U \in \gamma'$ by definition.

1 \Rightarrow 3: We are given $x \in X$ and $B \in \mathcal{B}$ with $x \in B$. Since $B \in \gamma$ by definition and $\gamma \subset \gamma'$ by (1), we have $B \in \gamma'$. Since γ' is generated by \mathcal{B}' , $\exists B' \in \mathcal{B}'$ such that $x \in B' \subset B$.

□

Def. (lower limit topology). Let $X = \mathbb{R}$. The lower limit topology on \mathbb{R} is the topology generated by the basis $\{[a, b) \mid a < b\}$.

Def. (ℓ -open). A set $U \subset \mathbb{R}$ is lower limit open (ℓ -open) if $\forall x \in U, \exists a < b$ such that $a \leq x < b, [a, b) \subset U$. (i.e. $\exists \varepsilon > 0$ s.t. $(x, x+\varepsilon) \subset U$).

Subspace topology.

Def. Let (X, τ) be a topological space. $Y \subset X$ be a subset. Define a topology on Y : τ_Y the subspace topology by $\tau_Y = \{U \cap Y \mid U \in \tau\}$. That is, $V \subset Y$ is open in Y iff $V = Y \cap U$ for some open $U \subset X$.

Theorem. The subspace topology is a topology.

Proof: ① $\emptyset = \emptyset \cap Y$, $\Rightarrow \emptyset$ is open in Y .

② $Y = X \cap Y$, $\Rightarrow Y$ is open in Y .

③ Suppose $\{V_\alpha\}_{\alpha \in I}$ are open in Y . For each α : $V_\alpha = U_\alpha \cap Y$ for some U_α open in X .

$$\bigcup_{\alpha \in I} V_\alpha = \bigcup_{\alpha \in I} (U_\alpha \cap Y)$$

$$= \underbrace{\left(\bigcup_{\alpha \in I} U_\alpha \right)}_{\text{open in } X} \cap Y$$

Thus $\bigcup_{\alpha \in I} V_\alpha$ open in Y .

④ Suppose $\{V_i\}_{i=1}^n$ open in Y . For each i : $V_i = U_i \cap Y$ for some U_i open in X .

$$\bigcap_{i=1}^n V_i = \bigcap_{i=1}^n (U_i \cap Y)$$

$$= \left(\bigcap_{i=1}^n U_i \right) \cap Y$$

Thus, $\bigcap_{i=1}^n V_i$ open in Y .

In topological sense, we define subspace as any subset equipped with the subspace topology.

e.g. $X = \mathbb{R}$, τ standard topology. $Y = [0, 1]$

$[0, 1]$ is open in Y . $A = [0, \frac{1}{2}] = Y \cap (-\frac{1}{2}, \frac{1}{2})$, open in Y .

e.g. $X = \mathbb{R}$, with lower limit topology. τ_L . $Y = [0, 1]$

$[\frac{1}{2}, 1] = Y \cap [\frac{1}{2}, 2]$ is open in subspace topology.

$\{1\} = Y \cap [1, 2]$ is open in subspace topology.

$\{\frac{1}{2}\}$ is not open since $\nexists U$ open in τ_L such that $Y \cap U = \{\frac{1}{2}\}$.

Def. (Basis for the subspace topology of Y). If \mathcal{B} is a basis of X , then $\mathcal{B}_Y = \{B \cap Y \mid B \in \mathcal{B}\}$ is a basis for the subspace topology of Y .

e.g. \mathcal{B} for the subspace topology of $[0, 1] \subset \mathbb{R}$ standard consists of 3 kinds of sets.

(a, b) where $0 < a < b < 1$, $[0, b)$ $0 < b \leq 1$, $(a, 1]$ $0 \leq a < 1$.

e.g. $S^1 = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 = 1\} = \{(\cos t, \sin t) \mid t \in \mathbb{R}\}$

$$S^n = \{\vec{x} \in \mathbb{R}^{n+1} \mid \|\vec{x}\| = 1\}$$

Use the standard topology coming from the standard topology of \mathbb{R} .

For $a < b$, define $A_{a,b} := \{(\cos t, \sin t) \mid a \leq t \leq b\}$

• If $b - a > 2\pi$, then $A_{a,b} = S^1$.

• If $b - a < 2\pi$, then $A_{a,b}$ = open arc from $(\cos a, \sin a)$ to $(\cos b, \sin b)$.

Claim: $\{A_{a,b}\}$ are a basis for the topology S' .

proof: $\forall \vec{x} \in S'$, any $r > 0$. $B(\vec{x}, r) \cap S' = A_{a,b}$ for some a, b .

The $\{A_{a,b}\}$'s are intersections of S' with a basis for the topology of \mathbb{R}^2 .

Proposition Let X be a metric space. $Y \subset X$ be a subspace of X .

1. Take the metric topology on X and then the subspace topology on Y , γ_1 .

2. Consider $d_Y = \text{restriction of } d \text{ on } Y$, then take the metric topology, γ_2 .

Then $\gamma_1 = \gamma_2$

proof: The basis for the subspace topology γ_1 is $\{B_d(x, r) \cap Y \mid x \in X, r > 0\}$.

define $B_d^Y(y, r) = \{z \in Y \mid d(y, z) < r, y \in Y, r > 0\}$

then $B_d^Y(y, r) = B_d(y, r) \cap Y$

the basis for the metric topology on Y , γ_2 is $\{B_d^Y(y, r) \mid y \in Y, r > 0\} = \{B_d(y, r) \cap Y \mid y \in Y, r > 0\}$.

Given $B_d^Y(y, r)$ and $x \in B_d^Y(y, r)$, $x \in B_d(y, r) \cap Y \subset B_d^Y(y, r)$, therefore $\gamma_2 \subset \gamma_1$.

Conversely, WTS $\gamma_1 \subset \gamma_2$, that is, WTS $B_d(x, r) \cap Y$ is open in the metric topology γ_2 .

Given $y \in B_d(x, r) \cap Y$, $\exists s > 0$ such that $B_d(y, s) \subset B_d(x, r)$. Then $B_d(y, s) \cap Y \subset B_d(x, r) \cap Y$

Thus $\gamma_1 \subset \gamma_2$.

$\Rightarrow \gamma_1 = \gamma_2$.

□

Product Topology

Def. (Product Topology). Let X, Y be topological spaces. The product topology on $X \times Y$ is the topology with basis $\{U \times V \mid U \subset X \text{ open}, V \subset Y \text{ open}\}$.

Given a basis \mathcal{B}_X for the topology of X , and a basis \mathcal{B}_Y for the topology of Y , the sets of the form $\{U \times V \mid U \in \mathcal{B}_X, V \in \mathcal{B}_Y\}$ is a (smaller) basis for the same topology.

e.g. $X = Y = \mathbb{R}$ with standard topology.

The basis for the product topology on $\mathbb{R} \times \mathbb{R}$ is $\{(a, b) \times (c, d)\}$.

Claim: This is just another way of describing the usual topology on \mathbb{R}^2 .

$$\mathcal{Y}_{\text{prod}} = \mathcal{Y}_{\text{standard}}$$

Proof: We use the fact that $\mathcal{T}_{\text{standard}} = \mathcal{T}_p$.

$$\forall \vec{x} \in \mathbb{R}^2, \vec{x} \in B_p(\vec{x}, r) = (x_1 - r, x_1 + r) \times (x_2 - r, x_2 + r) \subset B(\vec{x}, \sqrt{2}r)$$

$$\text{So } \mathcal{Y}_{\text{standard}} \subset \mathcal{T}_p.$$

But open rectangles are also open in standard topology. Given any rectangle, given any point in it, there is a square around it

$$\Rightarrow \mathcal{Y}_{\text{product}} \subset \mathcal{Y}_{\text{standard}}$$

□

Def (Projection Map) Given $X \times Y$. Let $\pi_1: X \times Y \rightarrow X$, $\pi_2: X \times Y \rightarrow Y$ be $\pi_1(x \times y) = x$, $\pi_2(x \times y) = y$.

Claim: π_1 and π_2 are continuous. Indeed. For $U \subset X$ open, $\pi_1^{-1}(U) = U \times Y$ open in product topology.

For $V \subset Y$ open, $\pi_2^{-1}(V) = X \times V$ open in product topology.

We want $U \times Y$ open $\wedge U \subset X$ open; $X \times V$ open $\wedge V \subset Y$ open, in order that π_1, π_2 is open. and then $U \times Y \cap X \times V = U \times V$ is open.

The product topology on $X \times Y$ is the coarsest topology for which π_1 and π_2 are continuous.

Lemma. If $A \subset X$ and $B \subset Y$, then the subspace topology on $A \times B$ as the subspace of $X \times Y$ agrees with the product topology coming from the subspace topologies of A and B .

Proof: $U \times V$ where $U \subset X$ open and $V \subset Y$ open is the general basis element for $X \times Y$.

Therefore $(U \times V) \cap (A \times B)$ is the general basis element for the subspace topology on $A \times B$.

$$(U \times V) \cap (A \times B) = (U \cap A) \times (V \cap B)$$

Since $U \cap A$ and $V \cap B$ are the general open sets for the subspace topologies on A and B , the set $(U \times V) \cap (A \times B)$ is the general basis element for the product topology coming from the subspace topology of A and B .

The bases agree \Rightarrow topologies are the same.

Lemma If (X, d_X) , (Y, d_Y) are metric spaces, then define a metric ρ on $X \times Y$ as

$$\rho((x, y), (x', y')) = \max\{d_X(x, x'), d_Y(y, y')\}. \text{ This is a metric. Product topology} = \text{Metric topology}.$$

Closed Sets and Limit Points

Def. (Closed). Let X be a topological space. A set $A \subset X$ is closed iff $X - A$ is open.

e.g. in \mathbb{R} : $[a, b]$ closed since $\mathbb{R} - [a, b] = (-\infty, a) \cup (b, +\infty)$ is open.
 $(-\infty, a]$ closed since $\mathbb{R} - (-\infty, a] = (a, +\infty)$ is open.

Notes. a). Not every set is open or closed.

b). \emptyset, X are both open and closed.

c). In \mathbb{R} : $[0, +\infty)$ is open. It is also closed since $\mathbb{R} - [0, +\infty) = (-\infty, 0) = \bigcup_{n \in \mathbb{N}} (-n, 0)$ open

d). Arbitrary intersection of closed sets are closed

$$X - \bigcap_{a \in A} A_a = \bigcup_{a \in A} (X - A_a) \text{ open} \Rightarrow \bigcap_{a \in A} A_a \text{ closed.}$$

e). Finite union of closed sets are closed.

$$X - \bigcup_{i=1}^n A_i = \bigcap_{i=1}^n (X - A_i) \text{ open} \Rightarrow \bigcup_{i=1}^n A_i \text{ closed.}$$

Def (Limit Point). Let $A \subset X$. A point $x \in X$ is called a limit point of A if for every open set $U \subset X$, $x \in U$, $\exists a \in A - \{x\}$ such that $a \in U$. ($a \in U \cap (A - \{x\}) \neq \emptyset$)

e.g. $A = (0, 1) \subset \mathbb{R}$. $\frac{1}{2}, 0$ are limit points, -4 is not.

$X = \mathbb{R}$. $A = \mathbb{Z}$. First it is easy to see that non-integers are not limit points.

Second, for $n \in \mathbb{Z}$, it is not a limit point. Given $U = (n - \frac{1}{2}, n + \frac{1}{2})$, there is no $a \in \mathbb{Z} - \{n\}$ such that $a \in U = (n - \frac{1}{2}, n + \frac{1}{2})$

Theorem. A is closed iff it contains all of its limit points.

Proof: (\Rightarrow). If A is closed, then $X - A$ is open. Given $x \in X - A$, there exists open set U_x s.t. $x \in U_x$ such that $U_x \cap A = \emptyset \Rightarrow U_x \cap (A - \{x\}) = \emptyset$. Thus no points in $X - A$ can be a limit point of A . Therefore A contains all its limit point.

(\Leftarrow). Suppose A contains all of its limit points. WTS $X - A$ is open.

Let $x \in X - A$. Since x is not a limit point, $\exists U_x$ open, $x \in U_x$,

$$U_x \cap (A - \{x\}) = \emptyset \Rightarrow U_x \cap A = \emptyset \Rightarrow U_x \subset X - A.$$

Therefore $X - A$ is open. A is closed. □.

Def. (Closure). For $A \subset X$, the closure of A (\bar{A}) is the smallest closed set that contains A .

It is the intersection of all closed sets that contains A .

- \bar{A} is closed.
- If C is any closed set s.t. $A \subset C$, then $\bar{A} \subset C$.

Def. (Interior). The interior of A ($\text{int}(A)$ or A°) is the union of all open sets that are contained in A .

$$\cdot \text{int}(A) \subset A \subset \bar{A}.$$

Lemma $x \in \bar{A} \Leftrightarrow \forall$ open set U s.t. $x \in U$, $U \cap A \neq \emptyset$.

Proof: (a). if $x \in A$ then obvious.

(b). if $x \notin A$, then, suppose \exists open set U , $x \in U$, $U \cap A = \emptyset$. Then, $A \subset X - U$.

Since $X - U$ is closed, $\bar{A} \subset X - U$. Then $x \notin \bar{A}$. Contradiction.

□.

Proposition: $\bar{A} = A \cup \{\text{limit points of } A\}$.

Proof: If $x \in \{\text{limit points of } A\}$, then $\forall U \ni x$, U open, $U \cap (A - \{x\}) \neq \emptyset \Rightarrow U \cap A \neq \emptyset$. Thus $x \in \bar{A}$.
Therefore $A \cup \{\text{limit points of } A\} \subset \bar{A}$.

If $x \in \bar{A}$, then, (a). $x \in A$ is trivial

(b). $x \notin A$. Since $x \in \bar{A}$, then $\forall U \ni x$ open, $U \cap A \neq \emptyset$. Thus, $\exists a \in A$, $a \in U \cap \{x\}$.
Therefore, x is a limit point of A .

Therefore $\bar{A} \subset A \cup \{\text{limit points of } A\}$

□

In general: A is closed $\Leftrightarrow A = \bar{A}$

$x \in \bar{A} \Leftrightarrow \forall$ open set $U \ni x$, $U \cap A \neq \emptyset$

$\bar{A} = A \cup \{\text{limit points of } A\}$

$\text{int}(A) = X - \overline{(X - A)}$

e.g. in \mathbb{R} . $\overline{[a, b]} = [a, b]$.

Def (Boundary). Boundary of A $\text{bd}(A) := \bar{A} - \text{int}(A)$

Lemma. $x \in \text{bd}(A)$ iff every open set $U \ni x$ intersects both A and $X - \bar{A}$.

Proof: Let $x \in \bar{A} - \text{int}(A)$. Then $x \in \bar{A}$, thus $\forall U \ni x$ open, $U \cap A \neq \emptyset$.

Suppose that $\exists U_x \ni x$, $U_x \cap (X - \bar{A}) = \emptyset$, then $U_x \subset \bar{A}$, and \bar{A} is open. Contradiction.

□

e.g. $A = \mathbb{Z} \subset \mathbb{R}$. $\bar{A} = A$. $\text{int}(A) = \emptyset \Rightarrow \bar{A} = A = \text{bd}(A)$.

$[0, 1] \times \{0\} \subset \mathbb{R}^2$ is closed, $\text{int} = \emptyset$.

$X = \mathbb{R}$. $A = \mathbb{Q}$. Claim: $\bar{A} = \mathbb{R}$.

Proof: WTS $\bar{A} \subset \mathbb{R}$. Let $x \in \bar{A}$. then since \bar{A} is the intersection of all the closed sets in \mathbb{R} containing A . $x \in \mathbb{R}$.

WTS $\mathbb{R} \subset \bar{A}$. Let $x \in \mathbb{R}$. Then, for any $a \in \mathbb{R}$, $b \in \mathbb{R}$, $a < x < b$, $(a, b) \cap A \neq \emptyset$ by the density of rational numbers. $x \in \bar{A}$.

□

Def (Dense) A set A is dense in X if $\bar{A} = X$.

$\Rightarrow \mathbb{Q}$ is dense in \mathbb{R} .

601 4

601 4

$$15 \times 2652 = 39780$$

6012 4

$$12 \times 2652 = 31824$$

6011 3

Hausdorff Spaces

Motivation:

Def (Convergence of sequence in analysis). Given x_1, x_2, x_3, \dots , $\lim_{n \rightarrow \infty} x_n = L$ if $\forall \epsilon > 0, \exists N \text{ s.t. } |x_n - L| \leq \epsilon \forall n \geq N$.

Def (Convergence of sequence in topology). Given x_1, x_2, x_3, \dots , we say the sequence converges to x if for every open set $U \ni x$, $\exists N > 0$ such that $x_n \in U$ for all $n > N$.

In analysis if $x_n \rightarrow x$, such x is unique. However, consider $X = \{a, b, c\}$, $\mathcal{T} = \{\{a, b, c\}, \{a, b\}, \{b, c\}, \{b\}, \emptyset\}$. Given a sequence $x_n = b \forall n$, by definition it converges to a, b , and c .

To rule out these strange cases, we introduce Hausdorff space.

Def. A topological space is called Hausdorff if for every $x, y \in X$ with $x \neq y$, there exists open sets U, V such that $x \in U, y \in V, U \cap V = \emptyset$.

Proposition. Any metric space is Hausdorff.

Proof: Let $x, y \in X$ with $x \neq y$. Let $\epsilon := \frac{d(x, y)}{2}$.

WTS $B_d(x, \epsilon) \cap B_d(y, \epsilon) = \emptyset$.

If $\exists z \in B_d(x, \epsilon)$ and $z \in B_d(y, \epsilon)$, then

$$d(x, y) \leq d(x, z) + d(z, y) < 2\epsilon = d(x, y)$$

Contradiction!

□.

Proposition. Let X be Hausdorff. If a sequence $\{x_n\}$ converges to x also to y , then $x = y$.

Proof: if $x \neq y$, then we can choose $U \ni x, V \ni y$ open such that $U \neq \emptyset, V \neq \emptyset, U \cap V = \emptyset$.

Since convergence, $\exists N$ such that $\forall n > N, x_n \in U, x_n \in V$. Contradiction!

□.

Lemma. If X is Hausdorff, then all 1-point sets are closed.

Proof: Fix $x_0 \in X$. For all y such that $y \neq x_0$, $\exists U_y, V_y$ open such that $x_0 \in U_y, y \in V_y, U_y \cap V_y = \emptyset$.

Thus $V_y \subset X - \{x_0\}$. Thus $X - \{x_0\}$ is open, $\{x_0\}$ is closed.

□.

Def. (T_1 and T_2). A space is called T_1 if 1-point sets are closed, called T_2 if it is Hausdorff.
 $T_2 \Rightarrow T_1$. $T_1 \not\Rightarrow T_2$. Since:

Proposition. If X is infinite with finite complement topology, then X is T_1 but not T_2 .

Proof. $\mathcal{Y}_{\text{finite}} = \{U \subset X \mid X - U \text{ is finite, or } U = \emptyset\}$.

Let $U = X - \{x_0\}$ for some $x_0 \in X$. Then $X - U = \{x_0\}$ is finite. Thus U is open, $\{x_0\}$ is closed.

Suppose that X is Hausdorff. Then given arbitrary $x, y \in X$ with $x \neq y$, $\exists U \ni x, V \ni y$ open such that $U \cap V = \emptyset$. Since $X - U$ and $X - V$ is finite, let $A := X - U, B := X - V$, then $U = X - A, V = X - B$. Then, since $U \cap V = \emptyset$,

$$(X - A) \cap (X - B) = X - (A \cup B) = \emptyset$$

$$\Rightarrow X = A \cup B$$

However, $A \cup B$ is finite. Contradiction!

□.

Def ($T_{1.5}$) A space is $T_{1.5}$ if for every $x \neq y$, \exists open sets U, V s.t. $x \in U$ and $x \notin V, y \in V$ and $y \notin U$.
 $T_2 \Rightarrow T_{1.5} \Rightarrow T_1$.

Continuous Functions

Def. (Continuous functions). Given topological spaces X and Y . A function $f: X \rightarrow Y$ is continuous if for every $U \subset Y$ open, $f^{-1}(U)$ is open in X .

Facts about continuous functions.

1. Any constant function is continuous.

$$f: X \rightarrow Y, f(x) = y_0 \forall x. \text{ Then, } f^{-1}(U) = \begin{cases} X & \text{if } y_0 \in U \\ \emptyset & \text{otherwise.} \end{cases}$$

2. Identity Function $\text{id}_X: X \rightarrow X$ $\text{id}_X(x) = x$ is continuous.

3. If $f: X \rightarrow Y$, $g: Y \rightarrow Z$ both continuous, then $g \circ f: X \rightarrow Z$ is continuous.

$$\text{for } U \subset Z \text{ open, } (g \circ f)^{-1}(U) = f^{-1}(g^{-1}(U))$$

4. If $A \subset X$ and $f: X \rightarrow Y$ is continuous, then $f|_A: A \rightarrow Y$ is continuous. For $x \in A$: $f|_A(x) = f(x)$.
For $U \subset Y$ open, $f|_A^{-1}(U) = A \cap f^{-1}(U)$ open in subspace topology.

inclusion function: $i_A: A \rightarrow X$: $i_A(x) = x$ if $x \in A$.

5. If $f: X \rightarrow Y$ continuous, $B \subset Y$ is a subspace, $\text{Im}(f) \subset B$, then f is continuous as a function $X \rightarrow B$.

6. Let $f: X \rightarrow Y$. If X has the discrete topology, then any function $f: X \rightarrow Y$ is continuous.
If X has the indiscrete topology, then very few are (constant function is).

If $f: X \rightarrow Y$ is continuous, and we choose a different topology on X that is finer,
 $f: X' \rightarrow Y$ is still continuous. If we choose a coarser topology on Y , f is still continuous.

Theorem. Let X, Y be topological spaces. TFAE.

(1). f is continuous

(2). $\forall A \subset X, f(\bar{A}) \subset \overline{f(A)}$

(3). \forall closed set $B \subset Y, f^{-1}(B)$ is closed in X

(4). For each $x \in X$ and each open set $V \subset Y$, there is an open set $U \subset X$ such that $f(U) \subset V$.

Proof: (1) \Rightarrow (2) WTS if $x \in \bar{A}$, then $f(x) \in \overline{f(A)}$.

Let V be open in Y such that $f(x) \in V$. Then $f^{-1}(V)$ is open in X and $x \in f^{-1}(V)$. Since $x \in \bar{A}$, $f^{-1}(V)$ intersects A in some point y . Thus, V intersects $f(A)$ in some point $f(y)$. Thus $f(x) \in \overline{f(A)}$.

(2) \Rightarrow (3) Let B be closed in Y . Let $A := f^{-1}(B) = \{x | f(x) \in B\}$. WTS A is closed, that is, $\bar{A} = A$.

WTS $\bar{A} \subset A$.

Since $f(A) = f(f^{-1}(B)) \subset B$, if $x \in \bar{A}$, then

$$f(x) \in f(\bar{A}) \subset \overline{f(A)} \subset \overline{B} = B$$

then $x \in f^{-1}(B) = A$. Thus $\bar{A} \subset A$. A is closed.

(3) \Rightarrow (1). Let $A \subset Y$ be open. Then $Y - A$ is closed. Then $f^{-1}(Y - A) = X - f^{-1}(A)$ is closed, $f^{-1}(A)$ is open. Thus f is continuous.

(1) \Rightarrow (4). Let $x \in X$, V be open in Y such that $f(x) \in V$. Since f is continuous, $f^{-1}(V)$ is open, and $x \in f^{-1}(V)$. Then, \exists open set $U \subset X$ s.t. $x \in U \subset f^{-1}(V)$. Then $f(U) \subset V$.

(4) \Rightarrow (1). Let $V \subset Y$ be open. Let $x \in X$ be $x \in f^{-1}(V)$. Then $f(x) \in V$. By hypothesis, \exists open set $U_x \subset X$, $f(U_x) \subset V$. Thus, $U_x \subset f^{-1}(V)$. Thus $f^{-1}(V) = \bigcup_{x \in f^{-1}(V)} U_x$, it is open. \square .

Finite Products

Def (Product topology): Let X_1, \dots, X_n be topological spaces. The product topology on $X_1 \times \dots \times X_n$ has the basis $\{U_1 \times \dots \times U_n \mid U_i \subset X_i \text{ is open}\}$.

Let $\pi_i : X_1 \times \dots \times X_n \rightarrow X_i$ be the projection map. For open $U \subset X_i$, $\pi_i^{-1}(U) = X_1 \times \dots \times X_{i-1} \times U \times X_{i+1} \times \dots \times X_n$. π_i is continuous.

Given $f_1 : X \rightarrow Y_1, \dots, f_n : X \rightarrow Y_n$, define $f_1 \times \dots \times f_n : X \rightarrow Y_1 \times \dots \times Y_n$ by $(f_1 \times \dots \times f_n)(x) = (f_1(x), \dots, f_n(x))$

Theorem: $f_1 \times \dots \times f_n$ is continuous iff each f_i is continuous.

proof: (\Rightarrow). $f_i = \pi_i \circ f = \pi_i(f_1 \times \dots \times f_n)$

f continuous $\Rightarrow f_i$ continuous.

(\Leftarrow). Assume that all f_i is open. WTS for any basic open set $U \subset Y_1 \times \dots \times Y_n$, $f^{-1}(U)$ is open.
 $U = U_1 \times \dots \times U_n$ where $U_i \subset Y_i$ is open.

$$\begin{aligned} f^{-1}(U) &= f^{-1}(U_1 \times \dots \times U_n) \\ &= \{x \mid f_i(x) \in U_i \forall i\} \\ &= \bigcap_{i=1}^n \{x \mid f_i(x) \in U_i\} \\ &= \bigcap_{i=1}^n f_i^{-1}(U_i) \text{ open.} \end{aligned}$$

□

(claim). If X is a topological space, given $f, g : X \rightarrow \mathbb{R}$, we can define $f \pm g : X \rightarrow \mathbb{R}$ by
 $(f \pm g)(x) = f(x) \pm g(x)$, $f \cdot g$. If $g(x) \neq 0$, we can also define f/g .

If f and g are continuous, then so are $f \pm g$, $f \cdot g$, f/g .

proof. for $f+g$: define $+ : \mathbb{R}^2 \rightarrow \mathbb{R}$ by $+(x, y) = x+y$, $+$ is continuous.

Given f, g :

$$\begin{array}{ccccc} X & \xrightarrow{\substack{f \times g \\ \text{ctn}}} & \mathbb{R} \times \mathbb{R} & \xrightarrow{\substack{+ \\ \text{ctn}}} & \mathbb{R} \\ & \underbrace{\hspace{10em}}_{\text{ctn.}} & & & \end{array}$$

By this we can also define $-$, \cdot .

For division $/ : \mathbb{R} \times (\mathbb{R} - \{0\}) \rightarrow \mathbb{R}$: given continuous f, g , $f : X \rightarrow \mathbb{R}$, $g : X \rightarrow \mathbb{R} - \{0\}$,

$$\begin{array}{ccccc} X & \xrightarrow{\substack{f \times g \\ \text{ctn}}} & \mathbb{R} \times (\mathbb{R} - \{0\}) & \xrightarrow{\substack{/ \\ \text{ctn}}} & \mathbb{R} \\ & \underbrace{\hspace{10em}}_{\text{ctn.}} & & & \end{array}$$

□

The set of continuous functions $X \rightarrow \mathbb{R}$ is a commutative ring.

Lemma (The pasting lemma): Let $X = A \cup B$, where A and B are closed in X .

$f : A \rightarrow Y$, $g : B \rightarrow Y$ be continuous. If $f(x) = g(x) \forall x \in A \cap B$, then

f and g combine to give a new continuous function $h : X \rightarrow Y$

$$h(x) = \begin{cases} f(x) & \text{if } x \in A \\ g(x) & \text{if } x \in B \end{cases}$$

Proof: Let $C \subset Y$ be closed.

$$\begin{aligned} h^{-1}(C) &= (A \cup B) \cap h^{-1}(C) \\ &= (A \cap h^{-1}(C)) \cup (B \cap h^{-1}(C)) \\ &= h^{-1}(f(A) \cap C) \cup h^{-1}(g(B) \cap C) \\ &= f^{-1}(C) \cup g^{-1}(C) \quad \text{is closed} \end{aligned}$$

Thus h is continuous. \square

Q: If $X = A \cup B$ in general, $f|_A$ and $f|_B$ are continuous, is f necessarily continuous?

A: No. Consider on \mathbb{R} : $f(x) = \begin{cases} 1, & x > 0 \\ 0, & x \leq 0 \end{cases}$ $A = (0, +\infty)$ $B = (-\infty, 0]$.

Theorem. (1) If $X = \bigcup_{i=1}^n A_i$ where A_i is closed, $f: X \rightarrow Y$ is a function such that $f|_{A_i}$ is continuous for each i , then f is continuous.

(2) If $X = \bigcup_{\alpha} A_{\alpha}$ - each A_{α} is open, $f|_{A_{\alpha}}$ is continuous for each α , then f is continuous.

Proof: (1). To prove that f is continuous, WTS $f^{-1}(\text{closed}) = \text{closed}$.

$$\begin{aligned} \text{Given } C \subset Y \text{ closed}, \quad f^{-1}(C) &= (f^{-1}(C) \cap A_1) \cup \dots \cup (f^{-1}(C) \cap A_n) \\ &= f|_{A_1}^{-1}(C) \cup \dots \cup f|_{A_n}^{-1}(C). \end{aligned}$$

Each $f|_{A_i}^{-1}(C) = A_i \cap B_i$ for some $B_i \subset X$ closed, thus $f^{-1}(C)$ is closed.

(2). For U open in Y ,

$$\begin{aligned} f^{-1}(U) &= \bigcup_{\alpha} (f^{-1}(U) \cap A_{\alpha}) \\ &= \bigcup_{\alpha} f|_{A_{\alpha}}^{-1}(U) \end{aligned}$$

Each $f|_{A_{\alpha}}^{-1}(U) = A_{\alpha} \cap B_{\alpha}$ for some B_{α} open, thus $f^{-1}(U)$ is open. \square .

Infinite Product Topology

Def (Infinite Products): Given $\{X_\alpha\}_{\alpha \in J}$, J is some index set. Define $\prod_{\alpha \in J} X_\alpha = \{(x_\alpha) \mid x_\alpha \in X_\alpha\}$.

e.g. If $J = \mathbb{Z}_+$, $X_i = \mathbb{R}$, then $\mathbb{R}^\omega = \mathbb{R} \times \mathbb{R} \times \dots$
 $X^\omega = \{\text{all sequences of elements of } X\}$.

Note: $\mathbb{R}^\infty = \{\vec{x} \mid x_i = 0 \text{ for } i > 0\}$

$$\mathbb{R}^\omega = \prod_{i=1}^{\infty} \mathbb{R}$$

Def: (Box and product topology). On $\prod_{\alpha \in J} X_\alpha$:

Option 1: Take as our basis $\{\prod_{\alpha \in J} U_\alpha \mid U_\alpha \subset X_\alpha \text{ open}\}$. In this topology, for $\mathbb{R}^\omega, (-1, 1) \times (1, 3) \times (0, \pi) \times \mathbb{R} \times (0, +\infty) \times \dots$ would be a basis. This is called box topology.

Option 2: Take as our basis $\{\prod_{\alpha \in J} U_\alpha \mid U_\alpha \subset X_\alpha \text{ open, } U_\alpha = X_\alpha \text{ for all but finitely many } \alpha\}$.
e.g. in $\mathbb{R}^\omega, (-1, 1) \times (1, 3) \times (2, +\infty) \times \mathbb{R} \times \mathbb{R} \times \dots$ is one basis.

The generated topology is called the product topology.

Proposition: Product Topology \neq Box topology.

prf: C is obvious.

†: Claim: in \mathbb{R}^ω , $U = (-1, 1) \times (-1, 1) \times (-1, 1) \times \dots$ is open in box topology, not in product topology.

Indeed, if it were, then for $\vec{U} \in U$, $\exists B = V_1 \times \dots \times V_k \times \mathbb{R} \times \mathbb{R} \times \dots$ such that $\vec{U} \in B \subset U$.

However, $(0, 0, \dots, \underset{k}{\cancel{0}}, 3, 3, \dots) \in B$, not in U . $B \notin U$. Contradiction. \square .

Def (Projection map). Given $\prod_{\alpha \in J} X_\alpha$, for each index β , we have a projection map $\pi_\beta: \prod_{\alpha \in J} X_\alpha \rightarrow X_\beta$:
 $\pi_\beta(\vec{x}) = x_\beta$.

e.g. in \mathbb{R}^ω , $\pi_i^{-1}(U) = \mathbb{R} \times \dots \times U \times \mathbb{R} \times \dots$ If U is open in \mathbb{R} , then $\pi_i^{-1}(U)$ is open in both product and box topology.

\Rightarrow Theorem π_β is continuous using either topology.

Theorem: Given functions $f_\alpha: Z \rightarrow X_\alpha$, the product function $f := \prod_{\alpha \in J} f_\alpha: Z \rightarrow \prod_{\alpha \in J} X_\alpha$ is continuous iff f_α is continuous, using product topology but not the box topology.

prf: (\Rightarrow). $f_\alpha = \pi_\alpha \circ f$.

(\Leftarrow). Suppose f_α is continuous for each α . Let U be open in $\prod_{\alpha \in J} X_\alpha$ using box topology.

WTS $f^{-1}(U)$ is open in Z .

$U = U_1 \times \dots \times U_k \times X_{k+1} \times X_{k+2} \times \dots$, U_i open in X_i

$$f^{-1}(U) = \{x \in Z \mid f_1(x) \in U_1, \dots, f_k(x) \in U_k, f_{k+1}(x) \in X_{k+1}, \dots\}$$

$$= \bigcap_{i=1}^k \{x \in Z \mid f_i(x) \in U_i\}$$

$$= \bigcap_{i=1}^k f_i^{-1}(U_i) \text{ open.}$$

Therefore f is continuous. \square

Homeomorphism & Embedding

Def (Homeomorphism). A map $f: X \rightarrow Y$ is called a homeomorphism if (1) f is bijective
(2). f is continuous. (3) f^{-1} is continuous.

Def (Homeomorphic) $X \cong Y$ if \exists homeomorphism between them.

Given that f is a bijection, f is continuous means $\forall U \subset Y$ open, $f^{-1}(U)$ is open in X .

f^{-1} is continuous means that given V open in X , $(f^{-1})^{-1}(V) = \{y \in Y \mid f^{-1}(y) \in V\} = \{y \in Y \mid y \in f(V)\} = f(V)$ is open in Y . Doing relabel between X and Y is one-to-one.

e.g. ① $(a, b) \cong (c, d)$ by $f(x) = \frac{d-c}{b-a}x + \frac{bc-ad}{b-a}$

② $[0, 1] \cong [0, 1]$ by $f(x) = 1-x$

③ $(-1, 1) \cong \mathbb{R}$ by $f(x) = \frac{x}{1-x^2}$ or $f(x) = \tan(\frac{\pi}{2}x)$

④ $\mathbb{R} \cong (0, +\infty)$ by $f(x) = e^x$. $f^{-1}(x) = \ln(x)$.

⑤ open ball in $\mathbb{R}^n \cong \mathbb{R}^n$

⑥ $[0, 1] \cong [0, +\infty)$ by $f(x) = \frac{x}{1-x}$.

Therefore, homeomorphism can lose information on boundedness.

Note that not every continuous (in one direction) bijection is a homeomorphism.

e.g. ① $\text{id}: X \rightarrow X'$ where X, X' have different topology on X may not be a homeomorphism.

② let $f: \mathbb{R} \rightarrow S^1$ be $f(t) = (\cos(2\pi t), \sin(2\pi t))$.

let $g = f|_{[0, 1]}$. Then $g: [0, 1] \rightarrow S^1$ is a bijection. g is continuous, but g^{-1} is not.

Take $[0, \frac{1}{4}]$ open in the subspace topology of $[0, 1]$. Then $(g^{-1})^{-1}(U) = g(U) = \underbrace{\text{arc}}_{\text{not open in } S^1}$, not open in subspace topology of S^1 , for $\#$ open arc containing $(1, 0)$ that is contained in $(g^{-1})^{-1}(U)$.

$\#$ homeomorphism between $[0, 1]$ and S^1 . $[0, 1] \not\cong S^1$. $[0, 1]$ not compact, S^1 compact

Def (Embedding). A function $f: X \rightarrow Y$ is called an embedding if (1) it is injective
(2) it restricts to a homeomorphism $X \rightarrow \text{Im}(f)$, where $\text{Im}(f)$ is given the subspace topology.

$$\begin{array}{ccc} f: X & \rightarrow & Y \\ \text{bijection} & \searrow & \uparrow \\ & & \text{Im}(f) \end{array}$$

e.g. View g from the last example as $g: [0, 1] \rightarrow \text{Im}(g) = S^1 \subset \mathbb{R}^2$. It is injective continuous map, but not an embedding. (not a homeomorphism $[0, 1] \rightarrow S^1$).

Proposition: Given X, Y , $g: X \rightarrow Y$ continuous. Define $f: X \rightarrow X \times Y$ by $f(x) = (x, g(x))$. Then this is an embedding of f to $X \times Y$.

Proof: First, it is trivial to see that f is injective.

$$\begin{aligned} \text{Then, } \text{Im}(f) &= \{(x, g(x)) \mid x \in X\} \\ &= \{(x, y) \mid y = g(x)\} \\ &= \text{graph}(g) \end{aligned}$$

Indeed, the inverse function $\text{graph}(g) \rightarrow f$ is $(x, y) \mapsto x = \pi_1|_{\text{graph}(g)}$ is continuous. thus f restricts to a homeomorphism $X \rightarrow \text{Im}(f)$.

e.g. In $X \times X = \Delta = \{(x, x) \mid x \in X\} = \text{graph}(\text{id})$.

Often, we want to know whether a space can be embedded into a familiar space, for example, \mathbb{R}^n .

Proposition: (a). If $X \cong Y$, X is Hausdorff, then Y is Hausdorff.

(b). If X is Hausdorff, then any subspace of X is Hausdorff.

(c). If X is not Hausdorff and Y is, then X does not embed Y .

Proof: (a). \exists homeomorphism $f: X \rightarrow Y$.

Let $y_1, y_2 \in Y$. Then, since f is bijective, $f^{-1}(y_1), f^{-1}(y_2) \in X$.

Since X is Hausdorff, $\exists U, V$ open, $f^{-1}(y_1) \in U, f^{-1}(y_2) \in V, U \cap V = \emptyset$.

$y_1 \in f(U), y_2 \in f(V), f(U) \cap f(V) = \emptyset$, both are open.

Therefore Y is Hausdorff.

(b). Trivial. Just take the subspace topology.

(c). Suppose it does. that is, $\exists f: X \rightarrow \text{Im}(f) \subset Y$ a homeomorphism.

Since Y is Hausdorff, $\text{Im}(f)$ also is. Then, according to (a): $\text{Im}(f) \cong X$, $\text{Im}(f)$ is Hausdorff, it should be that X is also Hausdorff. Contradiction!

□

e.g. Let $X = \mathbb{R}$, $Y = \text{line with two origins}$



with basis $\mathcal{D}(a, b)$ where $ab > 0$

② $(a, 0) \cup \{\text{Taylor}\} \cup (0, b)$

③ $(a, 0) \cup (\text{Travis}) \cup (0, b)$

define $f: \mathbb{R} \rightarrow Y$ $f(x) = \begin{cases} x, & x \neq 0 \\ \text{Taylor}, & x = 0 \end{cases}$ then $\text{Im}(f)$ is Hausdorff.

An application of embedding is knot theory.

Def. (knot) a knot is an embedding of S^1 into \mathbb{R}^3

e.g.



Quotient Topology

Def. (Binary Relation) Let X be a set. A binary relation is a function $X \times X \rightarrow \{T, F\}$ such that if $(x, y) \mapsto T$, we say $x \sim y$; if $(x, y) \mapsto F$, we say $x \not\sim y$.

Def (Equivalence Relation). An equivalence relation is a relation \sim satisfying

- (1). $\forall x \in X, x \sim x$ (Reflexive)
- (2). $\forall x, y \in X, x \sim y \Rightarrow y \sim x$ (Symmetric)
- (3). $\forall x, y, z \in X, \text{ if } x \sim y, y \sim z, \text{ then } x \sim z$ (transitive).

Def. (Equivalence Classes) An equivalence relation determines a partition of X into equivalence classes. For $x \in X$, $[x] = \{y \mid y \sim x\}$.

If $x \sim y$, then $[x] = [y]$. If $x \not\sim y$, then $[x] \cap [y] = \emptyset$.

e.g. $X = \mathbb{Z}$. $x \sim y$ iff $3 \mid (x-y)$ ($x-y$ is divisible by 3).

\sim is an equivalence relation. since: $x \sim x : x-x=0, 3 \mid 0 = 0$

$$x \sim y \Rightarrow 3 \mid (x-y) \Rightarrow 3 \mid (y-x) \Rightarrow y \sim x.$$

$$x \sim y \Rightarrow 3 \mid (x-y), y \sim z \Rightarrow 3 \mid (y-z).$$

$$x \sim z = x-y+y-z \cdot 3 \mid x-z \Rightarrow x \sim z.$$

$$\text{E. Classes: } [0] = \{\dots -6, -3, 0, 3, 6, \dots\}$$

$$[1] = \{\dots -5, -2, 1, 4, 7, \dots\}$$

$$[2] = \{\dots -4, -1, 2, 5, 8, \dots\}.$$

Def (Quotient Space) X/\sim (or X^*) is the set of equivalent classes of X .

Thus, there is a natural surjective map $p: X \rightarrow X/\sim : p(x) = [x]$.

Here we first define some general notions.

Def. (Quotient map). Let X, Y be topological spaces. $p: X \rightarrow Y$ be a surjective map.

It is a quotient map provided that $U \subset Y$ is open iff $p^{-1}(U)$ is open in X .

Equivalently, it can be defined as $A \subset Y$ is closed iff $p^{-1}(A)$ is closed in X .

Def (Saturated). $C \subset X$ is saturated w.r.t. the surjective map $p: X \rightarrow Y$ if $\forall p^{-1}(y)$ such that $p^{-1}(y) \cap C \neq \emptyset, p^{-1}(y) \subset C$.

In other words, $C \subset X$ is saturated w.r.t. p if it is the complete preimage of some set in Y . i.e. $\exists U \subset Y, C = p^{-1}(U)$.

Proposition: p is a quotient map $\Leftrightarrow p$ is continuous & p maps saturated open sets in X to open sets in Y .

Proof: (\Rightarrow). Continuity is obvious. U open $\Rightarrow p^{-1}(U)$ open.

$A \subset X$ saturated open. Then $A = p^{-1}(B)$ for some $B \subset Y$. Also $p(A)$ is open in Y

$$p(A) = p(p^{-1}(B)) = p(\{x \mid p(x) \in B\}) \subset B$$

Also, since p is surjective, $\forall x \in B, \exists y \in A$ s.t. $p(y) = x \in p(A)$, $\Rightarrow B \subset p(A)$

Then $B = p(A)$ is open.

(\Leftarrow). Suppose $V \subset Y$. $p^{-1}(V)$ is open in X . WTS V is open in Y .

First, notice that $p^{-1}(V)$ is automatically saturated wrt p .

Then, $p(p^{-1}(V))$ is open. However, $p(p^{-1}(V)) = V$ since p is surjective onto its image.
continuous \Rightarrow entire Y is p 's image.

Thus V is open in Y . □

Def. $f: X \rightarrow Y$ is an open (closed) map if $\forall U \subset X$ open (closed), $f(U)$ is open (closed) in Y .

Proposition. Surjective continuous open (closed) maps are quotient maps.

Proof. obvious, just use the definition.

Def (Quotient Topology) Let X be a space. A be a set. If $p: X \rightarrow A$ is a surjective map. then \exists one topology γ on A relative to which p is a quotient map. γ is the quotient topology of p .

$$\gamma = \{U \subset A \mid p^{-1}(U) \text{ is open in } X\}.$$

Proposition. γ is a topology.

Proof: \emptyset and A are open, $p^{-1}(\emptyset)$, $p^{-1}(A)$ is open.

Let $\{U_\alpha\}_{\alpha \in J}$ be open sets. Then

$$p^{-1}\left(\bigcup_{\alpha \in J} U_\alpha\right) = \bigcup_{\alpha \in J} p^{-1}(U_\alpha) \text{ is open.}$$

Let $\{U_i\}_{i=1}^n$ be open sets. Then

$$p^{-1}\left(\bigcap_{i=1}^n U_i\right) = \bigcap_{i=1}^n p^{-1}(U_i) \text{ is open.} \quad \square$$

Back to our main purpose:

Let X be a topological space. \sim be an equivalence relation. $p: X \rightarrow X/\sim : p(x) = [x]$. Then,

Def. The quotient topology on X/\sim is: $U \subset X/\sim$ is defined to be open if $p^{-1}(U)$ is open.

Therefore through p : open set on $X/\sim \leftrightarrow$ saturated open set in X .

Note: A subset $U \subset X/\sim$ is a collection of equivalent classes, and the set $p^{-1}(U)$ is just the union of equivalent classes belonging to U .

Thus, a typical open set of X/\sim is a collection of equivalent classes whose union is an open set of X , according to the definition $\gamma = \{U \mid p^{-1}(U) \text{ open in } X\}$.

Theorem. Let $p: X \rightarrow Y$ be a quotient map. Let $A \subset X$ be a subspace that is saturated wrt p .

Let $q: A \rightarrow p(A)$ be the map obtained by restricting p to A .

(a). If A is either open or closed in X , then q is a quotient map.

(b). If p is either an open or closed map, then q is a quotient map.

Proof: First we check the following equations.

$$\begin{aligned} \textcircled{1} \quad q^*(V) &= p^*(V) \quad \text{if } V \subset p(A). \\ \textcircled{2} \quad p(U \cap A) &= p(U) \cap p(A) \quad \text{if } U \subset X. \end{aligned}$$

\textcircled{1}: Since $V \subset p(A)$, A is saturated, by def. we have $p^*(V) \subset A$.

Also since $q = p|_A$, we have $p^*(V) = q^*(V) = \{x \in A \mid p(x) \in V\}$.

\textcircled{2}: Since $\forall U, A \subset X$, $p(U \cap A) \subset p(U) \cap p(A)$, it suffices to prove that $p(U) \cap p(A) \subset p(U \cap A)$.

Suppose $y = p(u) = p(a)$ for some $u \in U$, $a \in A$. Since A is saturated, $a \in p^*(p(a))$,

$\Rightarrow p^*(p(a)) \subset A$. Then $A \supset p^*(p(a)) = \{x \mid p(x) = p(a)\}$, so that $u \in A$. Then, $y = p(u) \in p(U \cap A)$.

Now suppose A is open or p is open. Given $V \subset p(A)$. Since p is a quotient map,

we already have $U \subset A$ open $\Rightarrow q(U) = p|_A(U)$ is open in $p(A)$. Then it suffices to show that given $q^*(V)$ open in A , V is open in X .

Suppose A is open. Since $q^*(V)$ is open in A , A is open in X , we have $q^*(V)$ open in X . Since $q^*(V) = p^*(V)$, $p^*(V)$ is also open in X , so that V is open in X by the fact that p is a quotient map. Then, V is open in $p(A)$ since $p(A) \cap V = V$.

Now suppose p is open. Since $q^*(V) = p^*(V)$ and $q^*(V)$ is open in A , we have $p^*(V) = U \cap A$ for some U open in X . Now, $p(p^*(V)) = V$ since p is surjective. Then

$$V = p(p^*(V)) = p(U \cap A) = \underbrace{p(U)}_{\text{open in } Y} \cap p(A)$$

Thus V is open in $p(A)$.

Similar for the case of closedness. \square .

Proposition. Composition of two quotient maps is a quotient map.

Proof: Let p, q be two quotient maps, U be open.

$$\begin{aligned} (q \circ p)^*(U) &= \{x \mid q(p(x)) \in U\} \\ &= \{x \mid p(x) \in \{x \mid q(x) \in U\}\} \\ &= p^*(\{x \mid q(x) \in U\}) \\ &= p^*(q^*(U)), \text{ it is open.} \end{aligned}$$

Let $(q \circ p)^*(U)$ be open, then

$$U = q(p(p^*(q^*(U)))) = q(p((q \circ p)^*(U))) \text{ is also open.} \quad \square$$

However, the cartesian product of two quotient maps may not be a quotient map.

One need local compactness or both maps being open map.

Theorem Let $p: X \rightarrow Y$ be a quotient map. Let Z be a space and let $f: X \rightarrow Z$ be a map that is constant on each set $p^{-1}(y) \quad \forall y \in Y$. Then, f induces a map: $\bar{f}: Y \rightarrow Z$ such that $\bar{f} \circ p = f$. \bar{f} is continuous iff f is continuous. \bar{f} is a quotient map iff f is a quotient map.

$$\begin{array}{ccc} X & \xrightarrow{f} & Z \\ p \downarrow & \nearrow \bar{f} & \\ Y & & \end{array}$$

Proof: $\forall y \in Y$, $f(p^{-1}(y))$ is a one-point set in Z . Denote this point as $\bar{f}(y)$, then we have defined a map $\bar{f}: Y \rightarrow Z$ by $\bar{f}(p(x)) = f(x)$, since $\bar{f}(y) = f(p^{-1}(y))$, let $x \in p^{-1}(y)$, then $y = p(x)$, $\bar{f}(y) = f(x) \Rightarrow \bar{f}(p(x)) = f(x)$.

If \bar{f} is continuous, then $f = \bar{f} \circ p$ is continuous.

Conversely, suppose f is continuous. Given $V \subset Z$ open, $f^{-1}(V)$ is open in X . However $f^{-1}(V) = p^{-1}(\bar{f}^{-1}(V))$. Since p is a quotient map, $\bar{f}^{-1}(V)$ is open in Y . Thus \bar{f} is continuous.

If \bar{f} is a quotient map, then $f = \bar{f} \circ p$ is a quotient map.

Conversely, suppose \bar{f} is a quotient map. Since f is surjective, so is \bar{f} .

Let $V \subset Z$. Since $\bar{f}^{-1}(V) = f(p^{-1}(V))$, f continuous, p^{-1} continuous $\Rightarrow \bar{f}$ continuous.

WTS V is open in Z if $\bar{f}^{-1}(V)$ is open in Y .

Since $\bar{f}^{-1}(V)$ is open in Y , we have $p^{-1}(\bar{f}^{-1}(V))$ is open in X since p is continuous.

Since $f^{-1}(V) = p^{-1}(\bar{f}^{-1}(V))$, we have $f^{-1}(V)$ is open in X . Since f is a quotient map, V is open in Z . □

Corollary: Let $f: X \rightarrow Z$ be a surjective continuous map. Let $Y = X/\sim$ be the following collection of equivalent classes of X : $X/\sim = \{f^{-1}(z) | z \in Z\}$ (in other word, $x \sim y$ in X iff $f(x) = f(y) = z$ for some $z \in Z$). Given X/\sim the quotient topology.

$$\begin{array}{ccc} X & \xrightarrow{f} & Z \\ p \downarrow & \nearrow \bar{f} & \\ X/\sim & & \end{array} \quad f = \bar{f} \circ p$$

(a). The map f induces a bijective continuous map $\bar{f}: X/\sim \rightarrow Z$, which is a homeomorphism iff f is a quotient map.

(b). If Z is Hausdorff, then so is X/\sim .

Proof: first, check that f is constant on $p^{-1}(y) \quad \forall y \in X/\sim$. For $[x] \in X/\sim$, $x \sim y$ if $p(x) = p(y) = [x]$ $p^{-1}([x]) = [x], [y] \Rightarrow f([x]) = f([y])$ by definition of the equivalence class here.

Then, by the previous theorem, f induces a continuous map $\bar{f}: X/\sim \rightarrow Z$. \bar{f} is bijective since if $\bar{f}([x]) = \bar{f}([y])$, then $f(p^{-1}([x])) = f(p^{-1}([y])) \Rightarrow p^{-1}([x])$ and $p^{-1}([y])$ are in the same equivalence class, therefore $[x] = [y]$.

Suppose \bar{f} is a homeomorphism, then both \bar{f} and p are quotient, which makes $f = \bar{f} \circ p$ quotient.
 Conversely, suppose f is a quotient map. By the previous theorem, \bar{f} is a quotient map.
 Also, since \bar{f} is bijective, \bar{f} is a homeomorphism.

Suppose Z is Hausdorff. Given $[x], [y] \in X/\sim$, $\bar{f}([x]) \neq \bar{f}([y])$. \exists open $U \ni \bar{f}([x]), V \ni \bar{f}([y])$, $U \cap V = \emptyset$. Since \bar{f} is bijectively continuous, $[x] \in \bar{f}^{-1}(U), [y] \in \bar{f}^{-1}(V)$, $\bar{f}^{-1}(U) \cap \bar{f}^{-1}(V) = \emptyset$.
 $\Rightarrow X/\sim$ Hausdorff. \square

e.g. $X = [0, 1]$. \sim : $x \sim y \iff x - y \in \mathbb{Q}$. $f: [0, 1] \rightarrow \mathbb{R}^2$: $f(x) = (\cos(2\pi x), \sin(2\pi x))$.

f is a continuous surjective quotient map onto its image S^1 . Conclusion: $X/\sim \cong S^1$.

$$\begin{array}{ccc} [0, 1] & \xrightarrow{f} & S^1 \\ p \downarrow & \nearrow \bar{f} \cong & \\ [0, 1]/\sim & & \end{array}$$

Generally, suppose given an \sim on X . Figure out something about X/\sim . If we can write a function $f: X \rightarrow Z$ such that $x \sim x'$ iff $f(x) = f(x')$, then $\bar{f}: X/\sim \rightarrow Z$ is injective since $\bar{f}([x]) = f(x)$. Then X/\sim is in bijection with $\text{Im}(f)$. By the previous corollary, if f is continuous, \bar{f} is. If f is a quotient map, then \bar{f} is a homeomorphism to $\text{Im}(f)$ and an embedding into Z .

More examples for open & closed maps:

e.g. $\pi_1: X \times Y \rightarrow X$ is open, but $\pi_1: \mathbb{R}^2 \rightarrow \mathbb{R}$ is not closed. Consider $C = \{(x, y) | xy = 1\}$, $\pi_1(C) = \mathbb{R} - \{0\}$.
 $f: \mathbb{R} \rightarrow \mathbb{R}$: $f(x) = x^2$ is closed but not open: $f(\mathbb{R}) = [0, +\infty)$.

e.g. $X = \mathbb{R}$, $\sim: x \sim y$ iff $x - y \in \mathbb{Q}$. Then, X/\sim is uncountable.

Suppose countable. $X/\sim = \{[x_i], [x_j], \dots\}$. $x_i - x_j \notin \mathbb{Q} \quad \forall i, j$. Then, since this is a partition of \mathbb{R} , $\mathbb{R} = \bigcup_{i=1}^{\infty} \{x | x \in [x_i]\}$, each $[x_i] = x + \mathbb{Q}$ countable $\Rightarrow \mathbb{R}$ countable. Contradiction!


The topology is the indiscrete topology. The only saturated open sets are \emptyset and \mathbb{R} .

$\phi = P^{-1}(\emptyset)$ is trivial.

To see why \mathbb{R} is, if $U \subset \mathbb{R}$ is non-empty saturated and open, then U contains some rational number $r \in \mathbb{Q}$. $r \in U = P(V)$ for some $V \subset X/\sim$. $P(r) = [r] = \mathbb{Q} = P(U) \subset V$. However, $\underbrace{\mathbb{Q} = P^{-1}(\mathbb{Q})}_{\text{saturated}} = P^{-1}(P(U)) \subset U = P(V)$. $\mathbb{Q} \subset U$. Also U open. $\Rightarrow U = \mathbb{R}$.

\Rightarrow Quotient = $\{U | P^{-1}(U) \text{ open}\} = \{U | P^{-1}(U) = \emptyset \text{ or } P^{-1}(U) = \mathbb{R}\} = \{\emptyset, X/\sim\}$
 $\Rightarrow X/\sim$ is not Hausdorff even if X is Hausdorff. \star .



e.g. $X = \mathbb{R}^2 - \{0\}$. \sim : $(x, y) \sim (x', y')$ if ① $x = x'$ or ② if $x = x' = 0$, then $\text{sign}(y) = \text{sign}(y')$




$X/\sim \cong$ line with two origins. $f(x, y) = \begin{cases} x & \text{if } x \neq 0 \\ \text{Top} & \text{if } x=0, y>0 \\ \text{Bottom} & \text{if } x=0, y<0 \end{cases}$

e.g. $X = [0, 1]^2$. $\sim : (0, y) \sim (1, y) \quad \forall y \in [0, 1]$.

$X/\sim = I \times I/\sim$, An annulus in \mathbb{R}^3 . $X/\sim \cong S^1 \times [0, 1]$ by $(x, y) \mapsto (\cos(2\pi x), \sin(2\pi x), y)$




Or: it embeds into \mathbb{R}^2 by $(x, y) \mapsto ((1+y)\cos(2\pi x), (1+y)\sin(2\pi x))$.



e.g. $X = [0, 1]^2$, $\sim : (0, y) \sim (1, 1-y) \quad \forall y \in [0, 1]$

$X/\sim = I \times I/\sim$ is the Möbius strip.

Using cyclical coordinates, this can be embedded into \mathbb{R}^3 .



e.g. $X = [0, 1]^2$. $\sim : (0, y) \sim (1, y) \quad (x, 0) \sim (x, 1)$

$X/\sim = I \times I/\sim \cong ([0, 1]/\sim) \times ([0, 1]/\sim)$



$\cong S^1 \times S^1 = \{(x, y, z, w) \mid x^2 + y^2 = 1 \text{ and } z^2 + w^2 = 1\}$.

$\subset \mathbb{R}^2 \times \mathbb{R}^2 = \mathbb{R}^4$

Or we can write an embedding $I \times I/\sim \rightarrow \mathbb{R}^3$ by



$$(\theta, r, z) = (2\pi y, 2 + \cos(2\pi x), \sin(2\pi x))$$

$$\Rightarrow (x, y) \mapsto ((2 + \cos(2\pi x)) \cos(2\pi y), (2 + \cos(2\pi x)) \sin(2\pi y), \sin(2\pi x))$$



e.g. $X = [0, 1]^2$. $\sim : (0, y) \sim (1, y) \quad (x, 0) \sim (-x, 1)$

$X/\sim = I \times I/\sim$, a Klein bottle, K^2 .



$$\subset \mathbb{R}^3 \times \{0\} \subset \mathbb{R}^4$$

We can write down a continuous function $K^2 \rightarrow \mathbb{R}^3$ but not surjective. However, we can write down an embedding $K^2 \rightarrow \mathbb{R}^4$ using trig functions.

Theorem. \nexists embedding $K^2 \rightarrow \mathbb{R}^3$.

prf: using tools from algebraic topology.