

VE320 HW1 Solution

1.

$$E = h\nu = \frac{hc}{\lambda} \Rightarrow \lambda = \frac{hc}{E}$$

Gold: $E = 4.90 \text{ eV} = (4.90)(1.6 \times 10^{-19}) \text{ J}$

So,

$$\lambda = \frac{(6.625 \times 10^{-34})(3 \times 10^{10})}{(4.90)(1.6 \times 10^{-19})} = 2.54 \times 10^{-5} \text{ cm}$$

or

$$\lambda = 0.254 \mu\text{m}$$

Cesium: $E = 1.90 \text{ eV} = (1.90)(1.6 \times 10^{-19}) \text{ J}$

So,

$$\lambda = \frac{(6.625 \times 10^{-34})(3 \times 10^{10})}{(1.90)(1.6 \times 10^{-19})} = 6.54 \times 10^{-5} \text{ cm}$$

or

$$\lambda = 0.654 \mu\text{m}$$

2.

$$(a) \quad p = \frac{h}{\lambda} = \frac{6.625 \times 10^{-34}}{550 \times 10^{-9}} = 1.205 \times 10^{-27} \text{ kg-m/s}$$

$$v = \frac{p}{m} = \frac{1.2045 \times 10^{-27}}{9.11 \times 10^{-31}} = 1.32 \times 10^3 \text{ m/s}$$

or $v = 1.32 \times 10^5 \text{ cm/s}$

$$(b) \quad p = \frac{h}{\lambda} = \frac{6.625 \times 10^{-34}}{440 \times 10^{-9}} = 1.506 \times 10^{-27} \text{ kg-m/s}$$

$$v = \frac{p}{m} = \frac{1.5057 \times 10^{-27}}{9.11 \times 10^{-31}} = 1.65 \times 10^3 \text{ m/s}$$

or $v = 1.65 \times 10^5 \text{ cm/s}$

(c) Yes

3.

(a)

Simple cubic: $a = 4.50 \text{ \AA}$

$$(i) \quad (100) \text{ plane, surface density,}$$

$$= \frac{1 \text{ atom}}{(4.50 \times 10^{-8})^2} \Rightarrow \underline{4.94 \times 10^{14} \text{ cm}^{-2}}$$

$$(ii) \quad (110) \text{ plane, surface density,}$$

$$= \frac{1 \text{ atom}}{\sqrt{2}(4.50 \times 10^{-8})^2} \Rightarrow \underline{3.49 \times 10^{14} \text{ cm}^{-2}}$$

$$(iii) \quad (111) \text{ plane, surface density,}$$

$$= \frac{3\left(\frac{1}{6}\right) \text{ atoms}}{\frac{1}{2}(a\sqrt{2})(x)} = \frac{\frac{1}{2}}{\frac{1}{2} \cdot a\sqrt{2} \cdot \frac{a\sqrt{3}}{\sqrt{2}}} = \frac{1}{\sqrt{3}a^2}$$

$$= \frac{1}{\sqrt{3}(4.50 \times 10^{-8})^2} \Rightarrow \underline{2.85 \times 10^{14} \text{ cm}^{-2}}$$

(b)

Body-centered cubic

$$(i) \quad (100) \text{ plane, surface density,}$$

Same as (a),(i); surface density $\underline{4.94 \times 10^{14} \text{ cm}^{-2}}$

$$(ii) \quad (110) \text{ plane, surface density,}$$

$$= \frac{2 \text{ atoms}}{\sqrt{2}(4.50 \times 10^{-8})^2} \Rightarrow \underline{6.99 \times 10^{14} \text{ cm}^{-2}}$$

$$(iii) \quad (111) \text{ plane, surface density,}$$

Same as (a),(iii), surface density $\underline{2.85 \times 10^{14} \text{ cm}^{-2}}$

(c)

Face centered cubic

$$(i) \quad (100) \text{ plane, surface density}$$

$$= \frac{2 \text{ atoms}}{(4.50 \times 10^{-8})^2} \Rightarrow \underline{9.88 \times 10^{14} \text{ cm}^{-2}}$$

$$(ii) \quad (110) \text{ plane, surface density,}$$

$$= \frac{2 \text{ atoms}}{\sqrt{2}(4.50 \times 10^{-8})^2} \Rightarrow \underline{6.99 \times 10^{14} \text{ cm}^{-2}}$$

$$(iii) \quad (111) \text{ plane, surface density,}$$

$$= \frac{\left(3 \cdot \frac{1}{6} + 3 \cdot \frac{1}{2}\right)}{\frac{\sqrt{3}}{2}a^2} = \frac{4}{\sqrt{3}(4.50 \times 10^{-8})^2}$$

or $\underline{1.14 \times 10^{15} \text{ cm}^{-2}}$

4.

$$E_{avg} = \frac{3}{2} kT = \left(\frac{3}{2}\right)(0.0259) = 0.03885 \text{ eV}$$

Now

$$p_{avg} = \sqrt{2mE_{avg}} \\ = \sqrt{2(9.11 \times 10^{-31})(0.03885)(1.6 \times 10^{-19})}$$

or

$$p_{avg} = 1.064 \times 10^{-25} \text{ kg-m/s}$$

Now

$$\lambda = \frac{h}{p} = \frac{6.625 \times 10^{-34}}{1.064 \times 10^{-25}} = 6.225 \times 10^{-9} \text{ m}$$

or

$$\lambda = 62.25 \text{ \AA}$$

5.

$$(a) \quad E_n = \frac{\hbar^2 n^2 \pi^2}{2ma^2} = \frac{n^2 (1.054 \times 10^{-34})^2 \pi^2}{2(9.11 \times 10^{-31})(10 \times 10^{-10})^2} \\ = n^2 (6.018 \times 10^{-20}) \text{ J}$$

$$\text{or} \quad E_n = \frac{n^2 (6.018 \times 10^{-20})}{1.6 \times 10^{-19}} = n^2 (0.3761) \text{ eV}$$

Then

$$E_1 = 0.376 \text{ eV}$$

$$E_2 = 1.504 \text{ eV}$$

$$E_3 = 3.385 \text{ eV}$$

$$(b) \quad \lambda = \frac{hc}{\Delta E}$$

$$\Delta E = (3.385 - 1.504)(1.6 \times 10^{-19}) \\ = 3.01 \times 10^{-19} \text{ J}$$

$$\lambda = \frac{(6.625 \times 10^{-34})(3 \times 10^8)}{3.01 \times 10^{-19}}$$

$$= 6.604 \times 10^{-7} \text{ m}$$

$$\text{or} \quad \lambda = 660.4 \text{ nm}$$

6.

$$(a) \quad p = \frac{h}{\lambda} = \frac{6.625 \times 10^{-34}}{85 \times 10^{-10}} \\ = 7.794 \times 10^{-26} \text{ kg-m/s}$$

$$v = \frac{p}{m} = \frac{7.794 \times 10^{-26}}{9.11 \times 10^{-31}} = 8.56 \times 10^4 \text{ m/s}$$

$$\text{or} \quad v = 8.56 \times 10^6 \text{ cm/s}$$

$$E = \frac{1}{2} mv^2 = \frac{1}{2} (9.11 \times 10^{-31})(8.56 \times 10^4)^2 \\ = 3.33 \times 10^{-21} \text{ J}$$

$$\text{or} \quad E = \frac{3.33 \times 10^{-21}}{1.6 \times 10^{-19}} = 2.08 \times 10^{-2} \text{ eV}$$

$$(b) \quad E = \frac{1}{2} (9.11 \times 10^{-31})(8 \times 10^3)^2 \\ = 2.915 \times 10^{-23} \text{ J}$$

$$\text{or} \quad E = \frac{2.915 \times 10^{-23}}{1.6 \times 10^{-19}} = 1.82 \times 10^{-4} \text{ eV}$$

$$p = mv = (9.11 \times 10^{-31})(8 \times 10^3) \\ = 7.288 \times 10^{-27} \text{ kg-m/s}$$

$$\lambda = \frac{h}{p} = \frac{6.625 \times 10^{-35}}{7.288 \times 10^{-27}} = 9.09 \times 10^{-8} \text{ m}$$

$$\text{or} \quad \lambda = 909 \text{ \AA}$$

7.

$$\int_{-1}^{+3} A^2 \cos^2\left(\frac{\pi x}{2}\right) dx = 1$$

$$A^2 \left[\frac{x}{2} + \frac{\sin(\pi x)}{2\pi} \right]_{-1}^{+3} = 1$$

$$A^2 \left[\frac{3}{2} - \left(\frac{-1}{2} \right) \right] = 1$$

$$\text{so} \quad A^2 = \frac{1}{2}$$

$$\text{or} \quad |A| = \frac{1}{\sqrt{2}}$$

8.

(a) Region I: Since $V_o > E$, we can write

$$\frac{\partial^2 \psi_1(x)}{\partial x^2} - \frac{2m(V_o - E)}{\hbar^2} \psi_1(x) = 0$$

Region II: $V = 0$, so

$$\frac{\partial^2 \psi_2(x)}{\partial x^2} + \frac{2mE}{\hbar^2} \psi_2(x) = 0$$

Region III: $V \rightarrow \infty \Rightarrow \psi_3 = 0$

The general solutions can be written, keeping in mind that ψ_1 must remain finite for $x < 0$, as

$$\psi_1(x) = B_1 \exp(k_1 x)$$

$$\psi_2(x) = A_2 \sin(k_2 x) + B_2 \cos(k_2 x)$$

$$\psi_3(x) = 0$$

where

$$k_1 = \sqrt{\frac{2m(V_o - E)}{\hbar^2}} \quad \text{and} \quad k_2 = \sqrt{\frac{2mE}{\hbar^2}}$$

(b) Boundary conditions

At $x = 0$: $\psi_1 = \psi_2 \Rightarrow B_1 = B_2$

$$\frac{\partial \psi_1}{\partial x} = \frac{\partial \psi_2}{\partial x} \Rightarrow k_1 B_1 = k_2 A_2$$

At $x = a$: $\psi_2 = \psi_3 \Rightarrow$

At $x = a$: $\psi_2 = \psi_3 \Rightarrow$

$$A_2 \sin(k_2 a) + B_2 \cos(k_2 a) = 0$$

or

$$B_2 = -A_2 \tan(k_2 a)$$

(c)

$$k_1 B_1 = k_2 A_2 \Rightarrow A_2 = \left(\frac{k_1}{k_2} \right) B_1$$

and since $B_1 = B_2$, then

$$A_2 = \left(\frac{k_1}{k_2} \right) B_2$$

From $B_2 = -A_2 \tan(k_2 a)$, we can write

$$B_2 = -\left(\frac{k_1}{k_2} \right) B_2 \tan(k_2 a)$$

or

$$1 = -\left(\frac{k_1}{k_2} \right) \tan(k_2 a)$$

This equation can be written as

$$1 = -\sqrt{\frac{V_o - E}{E}} \cdot \tan\left[\sqrt{\frac{2mE}{\hbar^2}} \cdot a\right]$$

or

$$\sqrt{\frac{E}{V_o - E}} = -\tan\left[\sqrt{\frac{2mE}{\hbar^2}} \cdot a\right]$$

This last equation is valid only for specific values of the total energy E . The energy levels are quantized.