
Studies of $e^+e^- \to b\bar{b}$ channel at International Linear Collider

Using the b-quark charge measurement

BILOKIN S., PÖSCHL R., RICHARD F.

Laboratoire de'Acceleráteur Linéare

bilokin@lal.in2p3.fr





Abstract

This poster presents studies for the International Linear Collider (ILC), a linear electron-positron collider with a nominal center-of-mass energy from 250 GeV to 500 GeV.

The results of the detector study allow for an estimation of the ILC precision on the b-quark electroweak couplings and form factors. The ILC will be able to resolve the LEP anomaly in the $b\bar{b}$ production process. The ILC precision on the right-handed $Z^0b\bar{b}$ coupling, a prime candidate for effects of new physics, is calculated to be at least 5 times better than the LEP experiments.

Introduction

Aliquam non lacus dolor, *a aliquam quam* [2]. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Nulla in nibh mauris. Donec vel ligula nisi, a lacinia arcu. Sed mi dui, malesuada vel consectetur et, egestas porta nisi. Sed eleifend pharetra dolor, et dapibus est vulputate eu. **Integer faucibus elementum felis vitae fringilla.** In hac habitasse platea dictumst. Duis tristique rutrum nisl, nec vulputate elit porta ut. Donec sodales sollicitudin turpis sed convallis. Etiam mauris ligula, blandit adipiscing condimentum eu, dapibus pellentesque risus.

Aliquam auctor, metus id ultrices porta, risus enim cursus sapien, quis iaculis sapien tortor sed odio. Mauris ante orci, euismod vitae tincidunt eu, porta ut neque. Aenean sapien est, viverra vel lacinia nec, venenatis eu nulla. Maecenas ut nunc nibh, et tempus libero. Aenean vitae risus ante. Pellentesque condimentum dui. Etiam sagittis purus non tellus tempor volutpat. Donec et dui non massa tristique adipiscing.

Main Objectives

- 1. Lorem ipsum dolor sit amet, consectetur.
- 2. Nullam at mi nisl. Vestibulum est purus, ultricies cursus volutpat sit amet, vestibulum eu.
- 3. Praesent tortor libero, vulputate quis elementum a, iaculis.
- 4. Phasellus a quam mauris, non varius mauris. Fusce tristique, enim tempor varius porta, elit purus commodo velit, pretium mattis ligula nisl nec ante.
- 5. Ut adipiscing accumsan sapien, sit amet pretium.
- 6. Estibulum est purus, ultricies cursus volutpat
- 7. Nullam at mi nisl. Vestibulum est purus, ultricies cursus volutpat sit amet, vestibulum eu.
- 8. Praesent tortor libero, vulputate quis elementum a, iaculis.

Description of the heavy quark production

Electroweak production of the fermion pairs proceeds through the $f\bar{f}X$ vertex, where X represents neutral vector bosons, photon or Z^0 boson. The current at the $f\bar{f}X$ vertex can be expressed via form factors F as

$$\Gamma_{\mu}^{f\bar{f}X}(k^2,q,\bar{q}) = ie\{\gamma_{\mu}(F_{1V}^X(k^2) + \gamma^5 F_{1A}^X(k^2)) - \frac{\sigma_{\mu\nu}(q-\bar{q})^{\nu}}{2m_f}(iF_{2V}^X(k^2) + \gamma^5 F_{2A}^X(k^2))\}, \tag{1}$$

where $k^2=(q+\bar{q})^2$ is the four momentum squared of the exchanged vector boson, q and \bar{q} are the four vectors of the fermion f and antifermion \bar{f} and m_f is the fermion mass. Further, γ_μ and γ_5 are the Dirac matrices, and $\sigma_{\mu\nu}=i/2(\gamma_\mu\gamma_\nu-\gamma_\nu\gamma_\mu)$.

The Standard Model values of the form factors are the following:

$$F_{1V}^{f\gamma} = Q^f, \ F_{1A}^{f\gamma} = 0, \ F_{1V}^{fZ} = \frac{I^f - 2Q^f \sin^2 \theta_W}{2\cos \theta_W \sin \theta_W}, \ F_{1A}^{fZ} = -\frac{I^f}{2\cos \theta_W \sin \theta_W},$$
 (2)

and all F_2 factor are zero. In the Eq. 2 I^f is the weak isospin number, $I^t = 1/2$ for top and $I^b = -1/2$ for bottom quark and Q^f is the electric charge, $Q^t = 2/3$ and $Q^b = -1/3$.

The following definition of the left-handed and right handed $Z^0b\bar{b}$ couplings is used throughout the thesis:

$$g_L^Z = I^f - Q^f \sin^2 \theta_W, \ g_R^Z = -Q^f \sin^2 \theta_W,$$
 (3)

In case of the polarized beams, the fermion form factors can be expressed in terms of the helicity of the initial electrons [?]:

$$\mathcal{F}_{ij}^{L} = -F_{ij}^{\gamma} + \frac{-1/2 + \sin^2 \theta_W}{\cos \theta_W \sin \theta_W} \frac{s}{s - M_Z^2 + i\Gamma_Z M_Z} F_{ij}^Z, \tag{4}$$

$$\mathcal{F}_{ij}^{R} = -F_{ij}^{\gamma} + \frac{\sin^2 \theta_W}{\cos \theta_W \sin \theta_W} \frac{s}{s - M_Z^2 + i\Gamma_Z M_Z} F_{ij}^Z \tag{5}$$

where i=1,2 and $j=V,A,M_Z$ and Γ_Z are the Z^0 boson mass and width, respectively. The key expression for the studies is the differential cross section of $f\bar{f}$ production for elect

The key expression for the studies is the differential cross section of $f\bar{f}$ production for electron beam polarization I=L,R, expressed via the defined form factors:

$$\frac{d\sigma^{I}}{d\cos\theta} = \frac{3}{4}\mathcal{A}N_{c}\beta[(1+\cos^{2}\theta)[(\mathcal{F}_{1V}^{I} + \mathcal{F}_{2V}^{I})^{2} + (\beta\mathcal{F}_{1A}^{I})^{2}] - 4\cos\theta(\mathcal{F}_{1V}^{I} + \mathcal{F}_{2V}^{I})\beta\mathcal{F}_{1A}^{I} + \sin^{2}\theta[\gamma^{-2}(\mathcal{F}_{1V}^{I} + \gamma^{2}\mathcal{F}_{2V}^{I})^{2}]]$$
(6)

where $A = 4\pi\alpha^2/3s$ with α as the electromagnetic running coupling, N_c is the number of quark colors, β and γ are the velocity and the Lorentz factor of the produced fermion, respectively.


Results

Donec faucibus purus at tortor egestas eu fermentum dolor facilisis. Maecenas tempor dui eu neque fringilla rutrum. Mauris *lobortis* nisl accumsan. Aenean vitae risus ante.

Phasellus imperdiet, tortor vitae congue bibendum, felis enim sagittis lorem, et volutpat ante orci sagittis mi. Morbi rutrum laoreet semper. Morbi accumsan enim nec tortor consectetur non commodo nisi sollicitudin. Proin sollicitudin. Pellentesque eget orci eros. Fusce ultricies, tellus et pellentesque fringilla, ante massa luctus libero, quis tristique purus urna nec nibh.

Nulla ut porttitor enim. Suspendisse venenatis dui eget eros gravida tempor. Mauris feugiat elit et augue placerat ultrices. Morbi accumsan enim nec tortor consectetur non commodo. Pellentesque

condimentum dui. Etiam sagittis purus non tellus tempor volutpat. Donec et dui non massa tristique adipiscing. Quisque vestibulum eros eu. Phasellus imperdiet, tortor vitae congue bibendum, felis enim sagittis lorem, et volutpat ante orci sagittis mi. Morbi rutrum laoreet semper. Morbi accumsan enim nec tortor consectetur non commodo nisi sollicitudin.






Figure 2: Generated b-quark polar angle distribution compared to the final reconstructed b-quarks polar angle in left-handed case (a) and right-handed case (b) with overlaid background processes.

In hac habitasse platea dictumst. Etiam placerat, risus ac. Adipiscing lectus in magna blandit:

Treatments	Response 1	Response 2
Treatment 1	0.0003262	0.562
Treatment 2	0.0015681	0.910
Treatment 3	0.0009271	0.296

 Table 1: Table caption

Vivamus sed nibh ac metus tristique tristique a vitae ante. Sed lobortis mi ut arcu fringilla et adipiscing ligula rutrum. Aenean turpis velit, placerat eget tincidunt nec, ornare in nisl. In placerat.



Figure 3: Figure caption

Conclusions

- Pellentesque eget orci eros. Fusce ultricies, tellus et pellentesque fringilla, ante massa luctus libero, quis tristique purus urna nec nibh. Phasellus fermentum rutrum elementum. Nam quis justo lectus.
- Vestibulum sem ante, hendrerit a gravida ac, blandit quis magna.
- Donec sem metus, facilisis at condimentum eget, vehicula ut massa. Morbi consequat, diam sed convallis tincidunt, arcu nunc.
- Nunc at convallis urna. isus ante. Pellentesque condimentum dui. Etiam sagittis purus non tellus tempor volutpat. Donec et dui non massa tristique adipiscing.

Forthcoming Research

Vivamus molestie, risus tempor vehicula mattis, libero arcu volutpat purus, sed blandit sem nibh eget turpis. Maecenas rutrum dui blandit lorem vulputate gravida. Praesent venenatis mi vel lorem tempor at varius diam sagittis. Nam eu leo id turpis interdum luctus a sed augue. Nam tellus.

References

[1] A. B. Jones and J. M. Smith. Article Title. *Journal title*, 13(52):123–456, March 2013. [2] J. M. Smith and A. B. Jones. *Book Title*. Publisher, 7th edition, 2012.

Acknowledgements

Etiam fermentum, arcu ut gravida fringilla, dolor arcu laoreet justo, ut imperdiet urna arcu a arcu. Donec nec ante a dui tempus consectetur. Cras nisi turpis, dapibus sit amet mattis sed, laoreet.