

TEAM economic evaluation: clinical data variable list

Patient data

study_number	Patient ID	
wardtype	0=MMHU,1=geriatric ward, 2=general ward	
age	Age at admission	
gender	Gender	M: Male F: Female

Baseline

pi_residencecode	Permanent care home resident at admission	0: Community 1: Care home
------------------	--	------------------------------

Cognition

pi_mmse1 – pi_mmse11		MMSE items at admission
----------------------	--	----------------------------

Delirium

pi_drs1	DRS sleep disturbance	Questions 1 to 14, range 0 to 3
pi_drs2	DRS hallucinations	
pi_drs3	DRS delusions	
pi_drs4	DRS lability of affect	
pi_drs5	DRS language	
pi_drs6	DRS thought process	
pi_drs7	DRS motor agitation	
pi_drs8	DRS motor retardation	
pi_drs9	DRS orientation	
pi_drs10	DRS attention	
pi_drs11	DRS st memory	
pi_drs12	DRS lt memory	
pi_drs13	DRS visuospatial ability	
pi_drs14	DRS temporal onset	
pi_drs15	DRS symptom fluctuation	Range 0 – 2
pi_drs16	DRS physical disorder	Range 0 – 2

EQ-5D

pi_eq5d_mob	Patient initial EQ-5D mobility	1: No problems 2: Some problems 3: Confined to bed
pi_eq5d_sc	Patient initial EQ-5D self care	1: No problems 2: Some problems 3: Unable
pi_eq5d_act	Patient initial EQ-5D usual	1: No problems

	activities	2: Some problems 3: Unable
pi_eq5d_pain	Patient initial EQ-5D pain	1: None 2: Moderate 3: Extreme
pi_eq5d_anx	Patient initial EQ-5D anxiety	1: None 2: Moderate 3: Extreme
pi_eq5d_mob_p	Proxy initial EQ-5D mobility	<i>See above for value coding</i>
pi_eq5d_sc_p	Proxy initial EQ-5D self care	
pi_eq5d_act_p	Proxy initial EQ-5D usual activities	
pi_eq5d_pain_p	Proxy initial EQ-5D pain	
pi_eq5d_anx_p	Proxy initial EQ-5D anxiety	
<i>Comorbidity</i>		
pi_med_con_tot	Number of medical conditions	
<i>Activities of daily living at admission</i>		
pi_feed_b	BADL feeding admission	Response set 0 1 2, see piq for coding
pi_groom_b	BADL grooming admission	Response set 0 1
pi_dress_b	BADL dressing admission	Response set 0 1 2
pi_bath_b	BADL bathing admission	Response set 0 1
pi_toilet_b	BADL toilet use admission	Response set 0 1 2
pi_bladd_b	BADL bladder admission	Response set 0 1 2
pi_bowel_b	BADL bowels admission	Response set 0 1 2
pi_trans_b	BADL transfer admission	Response set 0 1 2 3
pi_mob_b	BADL mobility admission	Response set 0 1 2 3
pi_stair_b	BADL stairs admission	Response set 0 1 2
<i>Neuropsychiatric inventory</i>		
pi_npi_del_f	NPI delusions frequency admission	0: None 1: Occasionally 2: Often 3: Frequent 4: V frequent
pi_npi_del_s	NPI delusions severity admission	0: None 1: Mild 2: Moderate 3: Marked
pi_npi_hal_f	NPI hallucinations frequency admission	
pi_npi_hal_s	NPI hallucinations severity admission	
pi_npi_agit_f	NPI agitation frequency admission	
pi_npi_agit_s	NPI agitation severity admission	
pi_npi_depress_f	NPI depression frequency admission	
pi_npi_depress_s	NPI depression severity admission	

pi_npi_nerv_f	NPI anxiety frequency admission
pi_npi_nerv_s	NPI anxiety severity admission
pi_npi_elat_f	NPI elation frequency admission
pi_npi_elat_s	NPI elation severity admission
pi_npi_apathy_f	NPI apathy frequency admission
pi_npi_apathy_s	NPI apathy severity admission
pi_npi_disinh_f	NPI disinhibition frequency admission
pi_npi_disinh_s	NPI disinhibition severity admission
pi_npi_irr_f	NPI irritability frequency admission
pi_npi_irr_s	NPI irritability severity admission
pi_npi_motorb_f	NPI motor behaviour frequency admission
pi_npi_motorb_s	NPI motor behaviour severity admission
pi_npi_sleep_f	NPI difficulty sleeping frequency admission
pi_npi_sleep_s	NPI difficulty sleeping severity admission
pi_npi_app_f	NPI appetite frequency admission
pi_npi_app_s	NPI appetite severity admission

Follow-up

Patient status data

po_status	Patient status at 90 days (string)	A: alive D: dead
Days_in_study	Days in study until death	

Cognition

po_mmse1 – po_mmse11	MMSE items at follow-up
----------------------	-------------------------

Patient DEMQoL

	<i>Numbers in variable names (e.g.demqol_1) correspond to question number on questionnaire</i>
po_demqol_1	Patient DEMQOL cheerful
po_demqol_2	Patient DEMQOL worried
po_demqol_3	Patient DEMQOL enjoying life
po_demqol_4	Patient DEMQOL frustrated
po_demqol_5	Patient DEMQOL confident
po_demqol_6	Patient DEMQOL full of energy

po_demqol_7	Patient DEMQOL sad
po_demqol_8	Patient DEMQOL lonely
po_demqol_9	Patient DEMQOL distressed
po_demqol_10	Patient DEMQOL lively
po_demqol_11	Patient DEMQOL irritable
po_demqol_12	Patient DEMQOL fed-up
po_demqol_13	Patient DEMQOL things unable to do
po_demqol_14	Patient DEMQOL forgetting recent things
po_demqol_15	Patient DEMQOL forgetting people
po_demqol_16	Patient DEMQOL forgetting what day
po_demqol_17	Patient DEMQOL thoughts muddled
po_demqol_18	Patient DEMQOL difficulty making decisions
po_demqol_19	Patient DEMQOL poor concentration
po_demqol_21	Patient DEMQOL getting on with people
po_demqol_22	Patient DEMQOL affection wanted
po_demqol_23	Patient DEMQOL people not listening
po_demqol_24	Patient DEMQOL making self understood
po_demqol_25	Patient DEMQOL getting help when needed
po_demqol_26	Patient DEMQOL getting to the toilet in time
po_demqol_27	Patient DEMQOL feel in self
po_demqol_28	Patient DEMQOL health overall
po_not_enough_money	Not enough money - included in DEMQoL in error
po_demqol_qol	Patient DEMQOL quality of life overall

EQ-5D

po_eq5d_mob	Patient EQ-5D mobility follow-up	1: No problems 2: Some problems 3: Confined to bed
po_eq5d_sc	Patient EQ-5D self care follow-up	1: No problems 2: Some problems 3: Unable
po_eq5d_act	Patient EQ-5D usual activities follow-up	1: No problems 2: Some problems 3: Confined to bed
po_eq5d_pain	Patient EQ-5D pain follow-up	1: None 2: Moderate 3: Extreme

po_eq5d_anx	Patient EQ-5D anxiety follow-up	1: None 2: Moderate 3: Extreme
po_eq5d_mob_p	Proxy EQ-5D mobility follow-up	<i>See above for value coding</i>
po_eq5d_sc_p	Proxy EQ-5D self care follow-up	
po_eq5d_act_p	Proxy EQ-5D usual activities follow-up	
po_eq5d_pain_p	Proxy EQ-5D pain follow-up	
po_eq5d_anx_p	Proxy EQ-5D anxiety follow-up	
<i>Activities of daily living at follow-up</i>		
po_feed_b	BADL feeding follow-up	Response set 0 1 2, see POQ for coding
po_groom_b	BADL grooming follow-up	Response set 0 1
po_dress_b	BADL dressing follow-up	Response set 0 1 2
po_bath_b	BADL bathing follow-up	Response set 0 1
po_toilet_b	BADL toilet use follow-up	Response set 0 1 2
po_bladd_b	BADL bladder follow-up	Response set 0 1 2
po_bowel_b	BADL bowels follow-up	Response set 0 1 2
po_trans_b	BADL transfer follow-up	Response set 0 1 2 3
po_mob_b	BADL mobility follow-up	Response set 0 1 2 3
po_stair_b	BADL stairs follow-up	Response set 0 1 2
<i>Neuropsychiatric inventory</i>		
po_npi_del_f	NPI delusions frequency follow-up	0: None 1: Occasionally 2: Often 3: Frequent 4: V frequent
po_npi_del_s	NPI delusions severity follow-up	0: None 1: Mild 2: Moderate 3: Marked
po_npi_hal_f	NPI hallucinations frequency follow-up	
po_npi_hal_s	NPI hallucinations severity follow-up	
po_npi_agit_f	NPI agitation frequency follow-up	
po_npi_agit_s	NPI agitation severity follow-up	
po_npi_depress_f	NPI depression frequency follow-up	
po_npi_depress_s	NPI depression severity follow-up	
po_npi_nerv_f	NPI anxiety frequency follow-up	
po_npi_nerv_s	NPI anxiety severity follow-up	
po_npi_elat_f	NPI elation frequency	

po_npi_elat_s	follow-up NPI elation severity
po_npi_apathy_f	follow-up NPI apathy frequency
po_npi_apathy_s	follow-up NPI apathy severity
po_npi_disinh_f	follow-up NPI disinhibition
po_npi_disinh_s	frequency follow-up NPI disinhibition severity
po_npi_irr_f	follow-up NPI irritability frequency
po_npi_irr_s	follow-up NPI irritability severity
po_npi_motorb_f	follow-up NPI motor behaviour
po_npi_motorb_s	frequency follow-up NPI motor behaviour
po_npi_sleep_f	severity follow-up NPI difficulty sleeping
po_npi_sleep_s	frequency follow-up NPI difficulty sleeping
po_npi_app_f	severity follow-up NPI appetite frequency
po_npi_app_s	follow-up NPI appetite severity
	follow-up

Carer proxy DEMQoL

<i>Numbers in variable names (e.g. cpdemqol_1) correspond to question number on questionnaire</i>	
po_cpdemqol_1	Carer proxy DEMQOL cheerful <i>Range 1 to 4 for all questions</i>
po_cpdemqol_2	Carer proxy DEMQOL worried
po_cpdemqol_3	Carer proxy DEMQOL frustrated
po_cpdemqol_4	Carer proxy DEMQOL full of energy
po_cpdemqol_5	Carer proxy DEMQOL sad
po_cpdemqol_6	Carer proxy DEMQOL content
po_cpdemqol_7	Carer proxy DEMQOL distressed
po_cpdemqol_8	Carer proxy DEMQOL lively
po_cpdemqol_9	Carer proxy DEMQOL irritable
po_cpdemqol_10	Carer proxy DEMQOL fed-up
po_cpdemqol_11	Carer proxy DEMQOL things to look forward
po_cpdemqol_12	Carer proxy DEMQOL memory
po_cpdemqol_13	Carer proxy DEMQOL forgetting things in past
po_cpdemqol_14	Carer proxy DEMQOL

po_cpdemqol_15	forgetting recent things Carer proxy DEMQOL
po_cpdemqol_16	forgetting people Carer proxy DEMQOL
po_cpdemqol_17	forgetting place Carer proxy DEMQOL
po_cpdemqol_18	forgetting day Carer proxy DEMQOL
po_cpdemqol_19	thoughts muddled Carer proxy DEMQOL
po_cpdemqol_20	difficulty making decisions Carer proxy DEMQOL
po_cpdemqol_21	making self understood Carer proxy DEMQOL
po_cpdemqol_22	keeping clean Carer proxy DEMQOL
po_cpdemqol_23	looking nice Carer proxy DEMQOL
po_cpdemqol_24	shopping Carer proxy DEMQOL
po_cpdemqol_25	using money Carer proxy DEMQOL
po_cpdemqol_26	looking after finances Carer proxy DEMQOL
po_cpdemqol_27	things taking longer Carer proxy DEMQOL
po_cpdemqol_28	getting in touch Carer proxy DEMQOL not
po_cpdemqol_29	enough company Carer proxy DEMQOL not
po_cpdemqol_30	being helpful Carer proxy DEMQOL not
po_cpdemqol_31	being useful Carer proxy DEMQOL
	physical health