

Investigation of the Effects of Critical Process Parameters such as Temperature, pH and pCO₂ on Charge Variants of mAbdi2

ABDiBIO



Öznur Özasan¹, Fatma Avcı-Köseer¹, Ümit Yiğiteli¹, Dila Kumaş¹, Büşra Demircan¹,
PDEng. Necla Sena Korkmaz¹, Dr. Ravi Kumar Lella¹

¹Abdi İbrahim Production Facility Orhan Gazi District Tunç Street No:3 Esenyurt/İstanbul
oznur.ozaslan@abdiibrahim.com.tr



1. INTRODUCTION

Charge variants in recombinant monoclonal antibodies (mAbs) are critical quality attributes (CQAs), which can affect function of antibodies in vitro and in vivo. Moreover, charge variants occur as cumulative effects of various post-translational modifications (PTMs) and chemical degradations on antibodies. These modifications can be occurred during both upstream and downstream processes. In the upstream process, critical process parameters (CPPs) such as temperature, pH and pCO₂ can be altered to ensure the desired product quality.

2. AIM

The subject of this poster is to investigate the effects of critical process parameters on the charge variants of mAbdi2, a biosimilar being developed at our biotechnology facility AbdiBio.

3. METHODS

- Recombinant Chinese Hamster Ovary (rCHO) cell line expressing the target monoclonal antibody was used in this study.




Figure 1. Process Flow Diagram

- In the first set of experiment, two parallel runs were performed to evaluate shift temperature effect: on the sixth day of production, operating temperatures at 37°C were reduced to 32°C and 34°C.
- In the second study, productions were carried out at pH set points of 7.2 and 7.4 to understand the effect of pH on charge variants.
- In the third set of experiment, effects of partial CO₂ pressure on charge variants were observed. pCO₂ accumulation exceeding 120 mmHg (high) and controlling under 120 mmHg (low) throughout the production were compared.

4. RESULTS & CONCLUSION




Figure 2. First study: Investigation of temperature shift degree on charge variants.




Figure 3. Second study: Investigation of pH on charge variants.




Figure 4. Third study: Investigation of pCO₂ on charge variants.

- In the first study where incubation temperature was shifted to 32°C and 34°C on the 6th day of production, lower percentage of acidic variants was observed in the product at lower shift temperature.
- When pH set points of 7.2 and 7.4 were compared, it was seen that as pH increased, the percentage of basic variants decreased whereas acidic variants increased in the product.
- Our last study showed that production at higher pCO₂ concentration resulted in an increase in the level of basic variants in the product.

5. DISCUSSION

- Lowering shift temperature decreased acidic and increased basic charge variant percentages in our candidate biosimilar product. Our findings align with those of Kishishita et al. (2015) and Sissolak et al. (2019).
- Irfan K. (2017) showed in his doctoral thesis that when pCO₂ accumulation in cell culture increases, the formation of basic variants also increases, consistent with our results.
- In accordance with Brunner et al.'s study (2016), our research demonstrated a decrease in acidic variants and a corresponding increase in basic variants at low pH values.

6. REFERENCES

- Brunner, M. Et al., (2016). Investigation of the interactions of critical scale-up parameters (pH, pO₂ and pCO₂) on CHO batch performance and critical quality attributes. *Bioprocess and Biosystems Engineering*, 40(2), 251–263. doi:10.1007/s00449-016-1693-7.
- Irfan, K. (2017). Carbon dioxide control in bioreactors and the application of principal component analysis to cell culture process data. (Doctoral Thesis, Newcastle University, England). Access adress: <http://theses.ncl.ac.uk/jspui/handle/10443/3776>.
- Kishishita, S. et al., (2015). Effect of temperature shift on levels of acidic charge variants in IgG monoclonal antibodies in Chinese hamster ovary cell culture. *Journal of Bioscience and Bioengineering*, 119(6), 700–705. doi:10.1016/j.jbiosc.2014.10.028.
- Sissolak, B. et al., (2019). Impact of mammalian cell culture conditions on monoclonal antibody charge heterogeneity: an accessory monitoring tool for process development. *Journal of Industrial Microbiology & Biotechnology*. doi:10.1007/s10295-019-02202-5.

7. INFO

Öznur ÖZASLAN, MSc.
AbdiBio – R&D Department
Process Development Project Leader



www.linkedin.com/in/oznurozaslan