

# Bio 1M: Evolutionary processes

## Evolution by natural selection

- Is something missing from the story I told last chapter?
  - Heritable **variation** in traits
  - **Selection** (i.e., differential reproductive success) *based on* these traits
- 

## Some genetics

- Our basic traits are determined by **genes**
- A location where a gene can occur is called a **locus** (pl. **loci**)
- A particular version of a gene is called an **allele**
- Complex organisms usually have two alleles at each locus
  - These can be the same, or different

## Loci

- Complex organisms usually have two alleles at each locus
  - These can be the same, or different
- An organism with different alleles at a particular locus is referred to as **heterozygous** (adj., n. form heterozygote)
- An organism with two copies of the same allele at a particular locus is referred to as **homozygous** (adj., n. form homozygote)

## Two definitions of evolution

- *Lecture*: heritable changes in species traits over time
- *Book*: changes in allele frequencies
- These definitions are consistent; use the one which helps you think clearly

## 1 Analyzing genotype frequencies — S23.1 (2ndEd S25.2)

## Genotypes and phenotypes

- A **genotype** is the collection of an individual's genes
- A **phenotype** is the collection of an individual's physiological and physical traits
  - What we can observe about an individual
  - Phenotype is largely (but by no means entirely) determined by genotype

## Example: peppered moths

- Check “peppered moths” or “Kettlewell’s experiment” on wikipedia
- Two different alleles possible at the wing color gene:  $A_1$  and  $A_2$ .
  - Individuals with  $A_1A_1$  *genotype* have light-winged *phenotype*
  - Individuals with  $A_2A_2$  *genotype* have dark-winged *phenotype*.
  - Individuals with  $A_1A_2$  *genotype* ???
- If individuals with genotype  $XY$  have the same phenotype (on average) as those with  $XX$ , we say that  $X$  is a **dominant** allele and  $Y$  is a **recessive** allele.
- If  $XY$  individuals have an intermediate phenotype (between  $XX$  and  $YY$ , we say  $X$  and  $Y$  are **co-dominant**.

## Analyzing genotype frequencies

- We analyze genotype frequencies as follows:
  - Make simple assumptions about how frequencies work
  - Calculate **expected frequencies** under our assumptions
  - Measure **observed frequencies** in the population
  - Look for evidence of systematic (not random) difference between expected and observed frequencies

## Making simple assumptions

- Expected frequencies are usually calculated by assuming that alleles assort randomly and independently, like flipping two coins, or rolling two dice

## Activity: Coin flipping

- I flip two fair coins (ie., each coin will land heads with probability  $1/2$ ).
- What is the probability of:
  - Two heads
  - Two tails?
  - One of each?
-

## Activity: Professional coin flipping

- A professional gambler can flip a coin so that it lands heads 70 the time. She flips two coins.
- What is the probability of:
  - Two heads
  - Two tails?
  - One of each?
- 

## Hardy-Weinberg distribution

- The Hardy-Weinberg distribution is the distribution expected if alleles work like coins (random and independent).
- If  $p$  is frequency of allele  $A_1$  and  $q$  is frequency of allele  $A_2$ , then:
  - Frequency of genotype  $A_1A_1$  is  $p^2$ .
  - Frequency of genotype  $A_2A_2$  is  $q^2$ .
  - Frequency of genotype  $A_1A_2$  is  $2pq$ .
- Why the 2?
  -

## Example: calculating allele frequencies

- I collect 20 peppered moths from a particular place, and find that 4 have genotype  $A_1A_1$ , 8 have genotype  $A_1A_2$ , and 8 have genotype  $A_2A_2$ .
- What is the observed frequency of each allele?
- What is the expected frequency of each genotype under the Hardy-Weinberg assumptions?
- Is this population in Hardy-Weinberg equilibrium?
  - 
  -

## What do we mean by expected?

- If we flip a fair coin 100 times, what is the expected number of heads?
  - What if we flip it 25 times?
- We don't expect to get exactly the expected value.
- The 'expected value' is an average of what is expected under our assumptions

## How do you know a coin is perfectly fair?

- You can never be sure that a coin is perfectly fair, you can only evaluate your evidence that it's more or less close to fair.
- Similarly, we never have evidence that a population is exactly in Hardy-Weinberg equilibrium
- We can only evaluate our evidence that it is far from (or close to) equilibrium

## Hardy-Weinberg equilibrium

- When do we expect genotype frequencies to behave like coins?
- Alleles selected at random from the previous generation:
  - 
  -
- If these assumptions hold exactly, we expect **Hardy-Weinberg equilibrium**
  - Hardy-Weinberg distribution, with no change in allele frequencies from generation to generation.

## Differences from equilibrium

- If we observe large differences from the Hardy-Weinberg equilibrium, this is usually a sign that mating is not random, or that natural selection is operating
- The analysis tells us that something is going on, but not what
- Hardy-Weinberg is a **null model**: it tells us what to expect if complicating effects are absent
- Without a null model, we couldn't start interpreting.

## Example: Human blood groups

- MN blood groups in different human populations are very close to Hardy-Weinberg equilibrium — Table 23.1 (2ndEd 25.1)
  - No evidence for non-random mating, or for fitness differences.

## Activity Human blood groups at the global level

- 
- 
-

## Example: Human HLA genes — Table 23.2 (2ndEd 25.2)

- HLA genes are used by the immune system to recognize disease-causing organisms
- Researchers hypothesized that heterozygous individuals may recognize more bacteria and viruses
- Data shows that more people are heterozygous for HLA genes than would be expected under the Hardy-Weinberg assumption

## Heterozygous HLA genes

- Why might more people be heterozygous for HLA genes than predicted by Hardy-Weinberg?
  - 
  - 
  - \*
    - \*
  - 
  - \*

## 2 Types of natural selection — S23.2 (2ndEd S25.3)

### 2.1 Directional selection

- **Directional selection** tends to move a population in a particular direction
  - Giraffe necks
  - Human brains

### Multi-directional selection

- Directional selection can change through time with the environment
  - Swallows may get larger during extreme cold spells, smaller again during normal weather
    - \* But we need to know whether the changes we saw were heritable
  - Finch beaks get thicker when food is scarce, and smaller when food is abundant
  - Why might small-beaked finches have advantages?
    - \*

## 2.2 Stabilizing selection

- **Stabilizing selection** tends to keep the population where it is
  - Example: human birthweights — Fig 23.4 (2ndEd 25.4)

## Connections between selection types

- What happens if the target of directional selection stays the same for a long time?
  -
- Examples?
  - 
  - 
  - 
  - \*

## 2.3 Disruptive selection

- Disruptive selection favors phenotypes *different* from the average value — Fig 23.5 (2ndEd 25.5)
  - Black-bellied seedcrackers
  - Animals that get eaten a lot may want to look different from their peers
- Disruptive selection may lead to **speciation** – the formation of new species.

## 2.4 Balancing selection

- Balancing selection tends to maintain allele diversity
  - When there is no single best allele
- **Heterozygote advantage**: when heterozygotes have higher fitness
- **Frequency dependence**: when rare types have higher fitness

## Example: The sickle cell phenotype

- Blood cells that can lose their shape and squash malaria parasites!
  - People heterozygous for this trait get less sick with malaria
  - People homozygous for this trait have too much instability and severe anemia
- This is an example of:
  -

## Example: seedcrackers

- What would happen if almost all of the seedcrackers had large bills?

—  
\*  
—  
—

## 3 Other evolutionary mechanisms

### 3.1 Genetic drift

- **Genetic drift** is change in allele frequencies due to random sampling:
  - Some individuals have more offspring than others due to chance events
  - Offspring receive certain parental alleles, and not others
- These factors will lead to an accumulation of random changes in allele frequencies

### Thought experiment

- Imagine flipping a fair coin 100 times
  - Repeat
- Now imagine choosing 100 alleles at random (with replacement) from a population of 50 *A* and 50 *B* alleles
  - Repeat, using new population as a starting point — Fig 23.6 (2ndEd 25.6)

### Small populations

- Drift is much stronger in small populations than in large ones (law of averages).
- Even if a population is big now, it may have been small in the past
  - **Founder effects** occur when a new population is started by a small number of individuals
  - **Bottlenecks** occur when a population becomes small, then large again
    - \* ... or, when a new genetic mutation takes over a population

## Fixation and loss

- An allele may drift to a frequency of 0 (it's **lost**) or of 1 (it's **fixed**)
- Advantageous alleles are often (not always) fixed
- Disadvantageous alleles are usually (not always) lost
- Alleles with **neutral** differences (no selective difference) will be fixed or lost at random
  - This is also true for alleles with small effects
- Drift tends to reduce genetic variation

### 3.2 Gene flow — S23.4 (2ndEd S25.4)

- **Gene flow** is the movement of alleles from one population to another
  - This happens when individuals move from one population to another and breed
- How we think about gene flow depends on how we choose to define a ‘population’
- Gene flow can be an obstacle to speciation; it helps keep populations similar

### 3.3 Mutation — S23.5 (2ndEd S25.5)

- **Mutations** are heritable errors in copying DNA
- Mutations are rare; by themselves they don't cause much evolution
- Mutations are extremely important to evolution, however:
  - 
  -

## Types of mutations

- Mutations can occur at many scales:
  - a single DNA base might change
  - chunks of DNA can be added or subtracted
  - whole genes (or whole chromosomes) can be duplicated
- New genetic sequence can come from:
  - copying errors
  - other organisms! **lateral gene transfer**



## Mutations are random

- Most mutations are **deleterious** – bad for fitness
- Very rarely mutations are **beneficial** – good for fitness
  - Such mutations are favored by natural selection

## Complex organisms

- Can complex organisms arise through random mutations?
  - A central question of biology
  - Large-scale evolution takes a *long* time
  - Beneficial changes can accumulate gradually
  - Much evidence of intermediate forms
- Check out videos of “evolution of the eye”

## What about sex?

- Sex does not *directly* change allele frequencies
  - Your book may say this means it’s not an “evolutionary process” – don’t worry about that
- It does act to bring alleles together (and to split them apart)
- Sex is not a source of new alleles
  - But it is a source of new combinations
- There is still active debate on the advantages and disadvantages of sex in evolution

## 4 Mating patterns

### 4.1 Inbreeding

- **Inbreeding** refers to mating between close relatives
- Since relatives will tend to share similar alleles, inbred populations will tend to differ from Hardy-Weinberg equilibrium in what way?
  -

## Inbreeding depression

- In many populations, it is observed that inbred individuals have lower fitness:
  - They are more likely to be homozygous for rare genetic defects
  - They are less likely to be heterozygous for immune-system genes
- Inbreeding depression is a serious concern for conservation
  - As populations get smaller, inbreeding becomes more common
- Wildlife studies show that panthers with both parents from Florida (small population) do not survive well — Fig 23.14
- Human demographic studies show strikingly lower survival for children of first cousins

## 4.2 Sexual selection

- **Sexual selection** is a form of natural selection
- Occurs when there is heritable variation in traits related to success in obtaining mates

### Example: zebra finches

- Males but not females have colorful orange beaks
- Hypothesis: these beaks make males more attractive to females — Fig 23.18 (2ndEd 25.17)

### Activity: Zebra finch experiments

- How would you test this hypothesis?
  - 
  -

### Results: Zebra finch experiments

- - 
  -
- Why not simply find and use birds whose beaks are naturally more or less orange?
  -
- What is a possible problem with the conclusion that females prefer birds with orange beaks?
  -

## Why the males?

- Males often have striking traits that females lack, used in courtship, or in battles for mates
  - Sexual **dimorphism** refers to trait differences between males and females
- Why do males more often have these traits than females?
  - Investment in reproduction
  - Variation in reproductive success

## Investment in reproduction

- In many species, females invest much more in each offspring than males do
  - Eggs are expensive, sperm are cheap
  - Females are often more involved in caring for offspring
- If females invest a lot in each offspring, they can maximize fitness by *being choosy about mates*
- If males invest little in each offspring, they can maximize fitness by *mating as much as possible*

## Testing the theory of sexual selection

- How might we test the theory that males compete sexually because females invest more in offspring?
  - - \*
      - - \*

## Variation in reproductive success

- Males often have greater variation in reproductive success than females do
- This is a side-effect of the fact that females usually invest a lot in each offspring
  - Reduces potential total number of offspring
  - Makes females desirable to males
- Greater variation in reproductive success means that winning contests is more important to male than female fitness

## Example: elephant seals

- Male elephant seals compete for control of breeding beaches
- Huge variation in reproductive success
- Huge size difference between males and females (strong sexual dimorphism) — Fig 23.19 (2ndEd 25.19)

## What about people?

- Men and women have pretty clear size differences
- How unequal is child-rearing in people?
  - 
  - 
  - \*
- To what extent do these principles even apply to people?
  - 
  - 
  -

## Conclusions

- **Mutation** (mistakes!) is the source of new variation
- **Natural selection** drives adaptation: selects variation that allows organisms to thrive in diverse settings
- **Sex** facilitates new combinations, but **sexual selection** can work against adaptation to the environment
- **Genetic drift** and **gene flow** are also non-adaptive drivers of evolution
- The adaptation we see is the result of all of these processes:
  - adaptive, non-adaptive, previously adaptive