1. Two populations of bugs live on two different species of trees in the same forest. The adult bugs breed together sometimes, but the hybrid offspring are rarely able to survive to adulthood, because they are not well adapted to either species of tree. This is an example of _______ isolation that will likely lead to ______.

- **A.** pre-zygotic; fusion
- **B.** post-zygotic; fusion
- C. pre-zygotic; natural selection for post-zygotic isolation
- **D.** post-zygotic; natural selection for pre-zygotic isolation

Use the following information for the next two questions. Biologists find a sub-population of plants that have a mutation that means that they can live only on acidic soils. The main population can't live on acidic soils at all. The patches of soil are bigger than the plants but close enough together that plants from different patches can still mate.

- 2. If these populations are to continue to diverge, they must overcome the obstacle of
 - A. natural selection
 - **B.** gene flow
 - C. competition
 - D. genetic drift
 - E. mutation
- **3.** If there are patches of acidic soil and patches of regular soil mixed together, the combined population is likely under
 - A. stabilizing selection
 - **B.** directional selection
 - C. disruptive selection
 - **D.** negative selection
 - E. positive selection
- **4.** A biologist is trying to classify ancient, extinct, shellfish into species based on the shapes of their shells, with no genetic information. They are most likely to use which applicable species concept?
 - A. Biological
 - **B.** Morphological
 - C. Ecological
 - **D.** Phylogenetic

5. Two populations of pine trees live in the same area, but open their flowers at different times, so that they rarely mate with each other. If a scientist argues that this fact makes them different species, which species concept are they applying?

- A. Biological
- B. Morphological
- C. Ecological
- **D.** Phylogenetic
- **6.** Which of the following definition best reflects an advantage in evolutionary fitness?
 - A. Organisms that tend to live longer
 - B. Organisms that are better at competing for resources
 - C. Organisms that tend to have more success at finding mates
 - D. Organisms that tend to produce more viable offspring
- 7. A population of maize plants has a recessive allele at a locus with simple dominance that heritably causes them to grow extra tall. The population genotype frequencies are near the Hardy-Weinberg expectation. A windstorm kills all of the extra tall plants, but not others. If we measure the population after the windstorm, we expect to find:
- **A.** genotype frequencies are near the new Hardy-Weinberg expectation (based on the new allele frequencies)
 - **B.** more heterozygotes than the expectation
 - C. more homozygotes than the expectation
 - **D.** more tall alleles than the expectation
 - E. more short alleles than the expectation
- 8. A species of frog was separated into two populations in the Andean mountains several thousand years ago by a cold spell that prevented them from living on hills separating two valleys. Offspring from the two populations are now observed to survive, grow and reproduce well on the hills and in the valleys. These two populations are most likely undergoing
 - A. Fusion
 - **B.** Divergence through reinforcement
 - C. Speciation through hybridization
 - **D.** Competitive exclusion
 - **E.** Allopolyploidy

9.	Inbre	eding	depres	ssion	is a	concer	n in	species	conser	vation,	because	in small	popu-
lat	ions,	indivi	duals	are n	nore	likely	to fi	nd mate	es who	are		, leading	to an
ino	crease	ed num	iber of	dele	teric	ous		gen	otypes.				

- A. close relatives; homozygous
- B. chosen from a small pool of unrelated individuals; homozygous
- C. close relatives; heterozygous
- D. chosen from a small pool of unrelated individuals; heterozygous
- 10. All else being equal, we might expect a large population to evolve more slowly than a small population because of differences in
 - **A.** gene flow
 - B. genetic drift
 - C. stabilizing selection
 - **D.** balancing selection
- 11. Xenopus tropicalis and Xenopus laevis are two closely related species of frog. X. laevis has two times the number of chromosomes as X. tropicalis. X. laevis most likely arose through:
 - A. allopatric speciation
 - **B.** polyploidy
 - C. reinforcement
 - **D.** coevolution
- **12.** If a population is currently subject to inbreeding, we would generally expect to see more ______than _____.
 - A. heterozygotes; homozygotes
 - **B.** homozygotes; heterozygotes
 - C. heterozygotes; than expected under Hardy-Weinberg
 - **D.** homozygotes; than expected under Hardy-Weinberg
- 13. All madrigals can sublimate. Alice is a madrigal. Bob can sublimate. Which of the following must be true?
 - A. Alice can sublimate
 - **B.** Bob is a madrigal
 - C. Both
 - **D.** Neither

14. All cows can fly. Dushoff can fly. Therefore, Dushoff is a cow. What is wrong with this logic (assuming we accept the assumptions)?

- **A.** nothing; the logic is correct
- B. The first assumption would need to say "Only cows can fly"
- C. The order of the assumptions would need to be reversed
- **D.** The two assumptions contradict each other
- **15.** Which of the following is the *strongest* correct statement we can make about life arising from non-life?
 - **A.** Life cannot arise from non-life
 - B. Life must have arisen from non-life at least once
- C. Life must have arisen from non-life at least five times, because there are five kingdoms of life
- **D.** Life must have arisen from non-life many times, because there is a huge diversity of different families of organisms

Use this information for the next three questions. At a certain allele, humans with the genotype NN are vulnerable to tuberculosis, those with the genotype CN have normal lung function and not vulnerable to tuberculosis, and those with genotype CC don't have normal lung function.

- **16.** This is an example of
 - **A.** polyploidy
 - **B.** inbreeding
 - C. simple dominance
 - **D.** complex dominance
- 17. In an environment with high rates of tuberculosis, the genotype CN has the highest fitness. This is an example of
 - A. directional selection
 - **B.** balancing selection
 - C. disruptive selection
 - **D.** negative selection on allele N
 - **E.** positive selection on allele N

18. In an environment with low rates of tuberculosis, the genotype NN has the highest fitness. This is an example of

- A. directional selection
- **B.** balancing selection
- C. disruptive selection
- **D.** negative selection on allele N
- E. positive selection on allele N
- **19.** Disruptive selection at the trait level is *most* likely to cause what kind of selection at the allele level?
 - A. Balancing selection
 - **B.** Stabilizing selection
 - C. Negative selection on rare alleles
 - **D.** Directional selection
- 20. Stabilizing selection at the trait level is often associated with
 - **A.** Positive selection on rare alleles
 - **B.** Negative selection on rare alleles
 - C. Frequency-dependent selection
 - **D.** Genetic drift
- 21. Yeasts grown in unstable environments develop the ability to quickly shut down their metabolism and survive in a resting state for long periods of time. The ability is itself an ______. The fact that the yeast can develop the ability is probably due to _____.
 - A. adaptation; adaptation
 - **B.** adaptation; acclimation
 - C. acclimation; adaptation
 - **D.** acclimation; acclimation
- **22.** E. coli bacteria grown for many generations in the lab with limited food sources developed the ability to use glucose as a food source. Their ancestors did not have this ability, and neither did control strains grown under similar conditions with ample food. These experiments provide evidence that:
 - **A.** Evolution can occur in the laboratory
 - **B.** Natural selection can occur in the laboratory
 - C. Bacteria can evolve the ability to use glucose for food
 - **D.** All of the above

- 23. When we say an allele is "fixed" in a population, it means:
 - A. It is the most fit allele for that locus under current conditions
 - **B.** A defect in that allele has been repaired
 - C. It is the only allele present at that locus in the population
 - **D.** The allele is not subject to mutation

Use the following information for the next three questions. A population of plants has one allele (R) for red flowers and one (W) for white flowers. The two alleles have simple dominance, with white being the dominant allele. There are 35 RR plants, 40 RW plants, and 25 WW plants.

- **24.** What is the allele frequency of W?
 - **A.** 0.25
 - **B.** 0.4
 - **C.** 0.45
 - **D.** 0.55
 - **E.** 0.65
- **25.** What is the genotype frequency of heterozygotes?
 - **A.** 0.25
 - **B.** 0.4
 - **C.** 0.45
 - **D.** 0.55
 - **E.** 0.65
- **26.** What is the expected phenotype frequency of plants with white flowers?
 - **A.** 0.25
 - **B.** 0.4
 - **C.** 0.45
 - **D.** 0.55
 - **E.** 0.65

Use the following information for the next two questions. Researchers raised 10 lines of bacteria starting from genetically identical individuals. After 100 generations, bacteria in 7 of these lines developed the ability to use one of the starches on the laboratory dishes as a new food sources. Subsequent experiments showed that these bacteria could out-compete ancestral strains, and the bacteria from the other three lines, under the laboratory conditions.

27. What is the most likely explanation for the observed change in the 7 lines?

- A. Random mutation
- **B.** Genetic drift
- C. Natural selection
- **D.** Random mutation combined with natural selection
- E. Genetic drift combined with natural selection
- **28.** What is the most likely explanation for why the other 3 lines did not show the same change?
 - A. The laboratory conditions were different for these lines
 - B. These lines were subject to sexual selection
 - C. These lines had a different set of random mutations
 - **D.** These lines were subject to more gene flow
- **29.** Whales have fairly complicated hip structures that seem to have little use. This is most likely because
- **A.** Whales are evolving to become more amphibious (better able to use both land and water environments)
 - **B.** These structures arose through random mutation
 - C. These structures arose through genetic drift
 - **D.** These structures are homologous to more useful structures in other mammals
- **30.** The biological species concept does not apply at all to
 - A. strictly asexual populations
 - **B.** extinct populations
- C. populations with many different "morphotypes" that all look different from each other
 - **D.** populations that can survive in a wide variety of habitats and conditions

Bio 1M03	Student #	_ Version 3
Short-answer	questions	
_	pen. Briefly show necessary work a information, even when the correct is	
microbes. They place was covered with ch	the 1700s did an experiment designed nutrient broth in two jars. One was eese cloth that was meant to allow a grew in the cloth-covered jar, but n	vas sealed completely, and one air to move in and out. They
/ \ - /	s a major problem with the experiment? G	
to the first experime	ne the researchers redid the experiment. What can you conclude? What century researchers who would have	t remains uncertain (from the
early flowering (and	f maize plants has a locus with two thus early breeding) and L is associ at flower early, 40 that flower late, a	iated with late flowering. You
/ (- /	e that these observations are driven of the <i>genotype</i> frequencies in this	v v

b) (1 point) What is your estimate of the allele frequencies?
c) (1 point) Based only on the information above, and <i>before</i> doing the calculations, how might you expect this locus to differ from the Hardy-Weinberg expectation and why?
d) (1 point) What is the calculated Hardy-Weinberg expectation based on these allele frequencies?
e) (1 point) How does the expectation compare to your observations? Be clear which numbers you are comparing to reach this conclusion.

Student # _____

 ${\bf Bio}~{\bf 1M03}$

Version 3