

UNIT 8A: Infectious disease

1 Introduction

Infectious disease

- Extremely common
- Huge impacts on ecological interactions
- A form of exploitation, but doesn't fit well into our previous modeling framework
 - How many people are there?
 - How many influenza viruses are there?
 - How do they find each other?

Disease agents

- Poll: Name an infectious agent that causes disease in humans.
- Disease agents vary tremendously:
 - Most **viruses** have just a handful of genes that allow them to hijack a cell and get it to make virus copies
 - *
 - **Bacteria** are independent, free-living cells with hundreds or thousands of chemical pathways
 - *
 - **Eukaryotic** pathogens are nucleated cells who are more closely related to you than they are to bacteria
 - *

Microparasites

- For infections with small pathogens (viruses and bacteria), we don't attempt to count pathogens, but instead divide disease into stages
 - Latently infected
 - Productively infected
 - Recovered

Microparasite models

- We model microparasites by counting the number of hosts in various **states**:
 - **Susceptible** individuals can become infected
 - **Infectious** individuals are infected and can infect others
 - **Resistant** individuals are not infected and cannot become infected
- More complicated models might include other states, such as latently infected hosts who are infected with the pathogen but cannot yet infect others

Models as tools

- Models are the tools that we use to connect scales:
 - individuals to populations
 - single actions to trends through time



2 Rate of spread

- Poll: For many diseases, especially new diseases, we can *observe* and *estimate* r .
 - - *
 - *
- Poll: Want to know what factors contribute to that, and how it relates to \mathcal{R} .
 -
 -



Basic reproductive number

- People in the disease field love to talk specifically about \mathcal{R}_0
- But they don't always mean the same thing when they say \mathcal{R}_0 :
 - Actual value of \mathcal{R} before an epidemic
 - Hypothetical value assuming no immunity
 - Hypothetical value assuming no immunity and no control efforts whatsoever
- Often easier to talk simply about \mathcal{R} .

Example: the West African Ebola epidemic



COVID in Ontario



Scales

- Which scale should we look at?

—

—

Population biology

- What quantities do we want to look at?

-
-
- *
-

Instantaneous rate of growth r

- What are the components?

-
- *
- *
-
- *
- *

- How do you think we estimate?

-
- *
-

 * Individuals \leftrightarrow Populations

Finite rate of growth λ

- Why do we want this?

-
-

- How do we calculate it?

-
-

Example

- $r \approx 0.14/\text{day}$ for early COVID spread
- What is λ ?
 - At a time scale of a day?
 - At a time scale of a week?

Reproductive number \mathcal{R}

- What is it?
 -
- Why do we want this?
 -
- How do we calculate it?
 -



Example

- $r \approx 0.14/\text{day}$
- What is our estimate of \mathcal{R} ?
 - When average length of infection $L = 5$ day?
 - When average length of infection $L = 10$ day?

Generation intervals

- Researchers try to estimate the *proportion* of transmission that happens for different **ages of infection**
- How long from the time you are *infected* to the time you *infect someone else*?
- Analogous to a life table
- The effective generation time \hat{G} has units of time

Generation intervals





Speed and risk

- Which is more dangerous, a fast disease, or a slow disease?
 - How are we measuring speed?
 - How are we measuring danger?
 - *What do we already know?*

Generation time and risk

- If we know \mathcal{R} , what does the generation time tell us about r ?
 -
- If we know r , what does the generation time tell us about \mathcal{R} ?
 -
- $\mathcal{R} = \exp(r\hat{G})$

Generation time and risk



Generation time and risk


- $\mathcal{R} = \exp(r\hat{G})$
- An intuitive view:
 - Epidemic speed = Generation strength \times Generation speed
- If we know generation speed, then a faster epidemic speed means:

-
- If we know epidemic speed, a faster generation speed means
-

3 Single-epidemic model

- Susceptible \rightarrow Infectious \rightarrow Recovered
- We also use N to mean the total population

Transition rates



- What factors govern movement through the boxes?
 - People get better independently
 - People get infected by infectious people

Conceptual modeling

- Poll: What happens in the long term if we introduce an infectious individual?

–
–
–

*

Interpreting

- Why might there not be an epidemic?

—
*
—

- Why doesn't everyone get infected?

—

Implementing the model

- The simplest way to implement this conceptual model is with differential equations:

—
$$\frac{dS}{dt} = -\beta \frac{SI}{N}$$

—
$$\frac{dI}{dt} = \beta \frac{SI}{N} - \gamma I$$

—
$$\frac{dR}{dt} = \gamma I$$

—
$$N = S + I + R$$

Quantities



State variables

- S, I, R, N : [people] or [people/ha]

Parameters

- Susceptible people have **potentially effective** contacts at rate β (units [1/time])
 - These are contacts that would lead to infection if the person contacted is infectious
 - Total infection rate is $\beta I/N$, because I/N is the proportion of the population infectious
- Infectious people recover at *per capita* rate γ (units [1/time])
 - Total recovery rate is γI
 - Mean time infectious is $D = 1/\gamma$ (units [time])

Simulating the model



Basic reproductive number

- Poll: What *unitless* parameter can you make from the model above?

—
—
—
*
—

Basic reproductive number implications

- Poll: What happens early in the epidemic if $\mathcal{R}_0 > 1$?

—

- What happens early in the epidemic if $\mathcal{R}_0 < 1$?

—

Effective reproductive number

- The effective reproductive number gives the number of new infections per infectious individual in a partially susceptible population:

—

- Is the disease increasing or decreasing?

—

-
- Why doesn't everyone get infected?

-
-

3.1 Epidemic size

- In this model, the epidemic always burns out
 - No source of new susceptibles
- Epidemic size is determined by:

-
-
- *
 - *
-
- *

Overshoot

- Why does more susceptibles at the beginning mean fewer susceptibles at the end?

-
-
-
-
-
-

Ebola example

- In September, the US CDC predicted “as many as” 1.5 million Ebola cases in Liberia by January
- In fact, their model predicted many *more* cases than that by April
- What happened?

What limits epidemics?

- Poll: What limits epidemics in our simple models?

—

- Poll: What else limits epidemics in real life?

—

—

—

4 Recurrent epidemic models


- Poll: If epidemics tend to burn out, why do we often see repeated epidemics?

—


—

Recurrent epidemics

Measles reports from England and Wales




Closing the circle



•

Closing the circle



•

—

Births and deaths

•

$$\frac{dS}{dt} = bN - \beta \frac{SI}{N} - dS$$

•

$$\frac{dI}{dt} = \beta \frac{SI}{N} - \gamma I - dI$$

-

$$\frac{dR}{dt} = \gamma I - dR$$

- We often assume $b = d$
 - \implies population is constant

4.1 Dynamics

Equilibrium

- At equilibrium, we know that $\mathcal{R}_{\text{eff}} = 1$
 - One case per case
 - Number of susceptibles at equilibrium determined by the number required to keep infection in balance
 - * $S/N = 1/\mathcal{R}_0$
- What does this remind you of?
 -
- Number of infectious individuals determined by number required to keep susceptibles in balance.
- As susceptibles go up, what happens?
 - Per capita replenishment goes down
 - Infections required goes down

Reciprocal control



- What happens to *equilibrium* if we protect susceptibles (move them to R class)?
 -
 -
 -
 -
- What else could happen?
 -
 -
 - *
 -

Reciprocal control

- Poll: What happens if we remove infectious individuals at a constant rate (find them and cure them or isolate them)?

—
—
*
—

Tendency to oscillate



Tendency to oscillate

- “Closed-loop” SIR models (ie., with births or loss of immunity):
 - Tend to oscillate
 - Oscillations tend to be damped
 - * System reaches an **endemic** equilibrium – disease persists

Source of oscillations

- Similar to predator-prey systems
- What happens if we start with too many susceptibles?

—
—
—
—



Persistent oscillations

- Poll: If oscillations tend to be damped in simple models, why do they persist in real life?

—
*
*
—
—
—
—
*

5 Reproductive numbers and risk

- At equilibrium, the proportion of people who are susceptible to disease should be approximately $S/N = 1/\mathcal{R}_0$
- Proportion “affected” (infectious or immune) should be approximately $V/N = 1 - 1/\mathcal{R}_0$
- If you have a single, fast epidemic, the size is also predicted by \mathcal{R}_0 .



Examples

- Ronald Ross predicted 100 years ago that reducing mosquito densities by a factor of 5 or so would *eliminate* malaria
- Gradual disappearance of polio, typhoid, etc., without risk factors going to zero
- Eradication of smallpox!

Threshold for elimination

- What proportion of the population should be vaccinated to eliminate a disease?

—

Examples:

- Polio has an \mathcal{R}_0 of about 5.
 - Poll: What proportion of the population should be vaccinated to eliminate polio?
-
- Measles has an \mathcal{R}_0 of about 20. What proportion of the population should be vaccinated to eliminate measles?
-
- If gonorrhea has an \mathcal{R}_0 of about 2, what proportion of unprotected sexual encounters should be protected to eliminate gonorrhea?
-

Persistence of infectious disease

- Why have infectious diseases persisted?
 - The pathogens *evolve*
 - Human populations are **heterogeneous**
 - * People differ in: nutrition, exposure, access to care
 - Information and misinformation
 - * Vaccine scares, trust in health care in general

Heterogeneity and persistence

- Heterogeneity *increases* \mathcal{R}_0
 - When disease is rare, it is concentrated in the most vulnerable populations
 - * Cases per case is high
 - * Elimination is harder
- Marginal populations
 - Heterogeneity could make it easier to concentrate on the most vulnerable populations and eliminate disease
 - Humans rarely do this, however: the populations that need the most support typically have the least access