UNIT 7: Infectious disease

Outline

Introduction

Rate of spread

Single-epidemic model Epidemic size

Recurrent epidemic models

Reproductive numbers and risk

Extremely common

- Extremely common
- ► Huge impacts on ecological interactions

- Extremely common
- Huge impacts on ecological interactions
- ► A form of exploitation, but doesn't fit well into our previous modeling framework

- Extremely common
- Huge impacts on ecological interactions
- A form of exploitation, but doesn't fit well into our previous modeling framework
 - How many people are there?

- Extremely common
- Huge impacts on ecological interactions
- A form of exploitation, but doesn't fit well into our previous modeling framework
 - How many people are there?
 - ► How many influenza viruses are there?

- Extremely common
- Huge impacts on ecological interactions
- A form of exploitation, but doesn't fit well into our previous modeling framework
 - How many people are there?
 - ▶ How many influenza viruses are there?
 - ► How do they find each other?

- Extremely common
- Huge impacts on ecological interactions
- A form of exploitation, but doesn't fit well into our previous modeling framework
 - How many people are there?
 - How many influenza viruses are there?
 - How do they find each other?

▶ Poll: Can you name an infectious agent that causes disease in humans?

- Poll: Can you name an infectious agent that causes disease in humans?
- ► Disease agents vary tremendously:

- Poll: Can you name an infectious agent that causes disease in humans?
- Disease agents vary tremendously:
 - Most viruses have just a handful of genes that allow them to hijack a cell and get it to make virus copies

- Poll: Can you name an infectious agent that causes disease in humans?
- Disease agents vary tremendously:
 - Most viruses have just a handful of genes that allow them to hijack a cell and get it to make virus copies
 - *****

- Poll: Can you name an infectious agent that causes disease in humans?
- Disease agents vary tremendously:
 - Most viruses have just a handful of genes that allow them to hijack a cell and get it to make virus copies
 - * influenza virus, Ebola virus, HIV, measles

- Poll: Can you name an infectious agent that causes disease in humans?
- Disease agents vary tremendously:
 - Most viruses have just a handful of genes that allow them to hijack a cell and get it to make virus copies
 - * influenza virus, Ebola virus, HIV, measles
 - Bacteria are independent, free-living cells with hundreds or thousands of chemical pathways

- Poll: Can you name an infectious agent that causes disease in humans?
- Disease agents vary tremendously:
 - Most viruses have just a handful of genes that allow them to hijack a cell and get it to make virus copies
 - * influenza virus, Ebola virus, HIV, measles
 - Bacteria are independent, free-living cells with hundreds or thousands of chemical pathways

, ,

- Poll: Can you name an infectious agent that causes disease in humans?
- Disease agents vary tremendously:
 - Most viruses have just a handful of genes that allow them to hijack a cell and get it to make virus copies
 - * influenza virus, Ebola virus, HIV, measles
 - Bacteria are independent, free-living cells with hundreds or thousands of chemical pathways
 - * Tuberculosis, anthrax, pertussis

- Poll: Can you name an infectious agent that causes disease in humans?
- Disease agents vary tremendously:
 - Most viruses have just a handful of genes that allow them to hijack a cell and get it to make virus copies
 - * influenza virus, Ebola virus, HIV, measles
 - Bacteria are independent, free-living cells with hundreds or thousands of chemical pathways
 - * Tuberculosis, anthrax, pertussis
 - Eukaryotic pathogens are nucleated cells who are more closely related to you than they are to bacteria


- Poll: Can you name an infectious agent that causes disease in humans?
- Disease agents vary tremendously:
 - Most viruses have just a handful of genes that allow them to hijack a cell and get it to make virus copies
 - * influenza virus, Ebola virus, HIV, measles
 - Bacteria are independent, free-living cells with hundreds or thousands of chemical pathways
 - * Tuberculosis, anthrax, pertussis
 - Eukaryotic pathogens are nucleated cells who are more closely related to you than they are to bacteria

•


- Poll: Can you name an infectious agent that causes disease in humans?
- Disease agents vary tremendously:
 - Most viruses have just a handful of genes that allow them to hijack a cell and get it to make virus copies
 - * influenza virus, Ebola virus, HIV, measles
 - Bacteria are independent, free-living cells with hundreds or thousands of chemical pathways
 - * Tuberculosis, anthrax, pertussis
 - Eukaryotic pathogens are nucleated cells who are more closely related to you than they are to bacteria
 - * Malaria, various worms

- Poll: Can you name an infectious agent that causes disease in humans?
- Disease agents vary tremendously:
 - Most viruses have just a handful of genes that allow them to hijack a cell and get it to make virus copies
 - * influenza virus, Ebola virus, HIV, measles
 - Bacteria are independent, free-living cells with hundreds or thousands of chemical pathways
 - * Tuberculosis, anthrax, pertussis
 - Eukaryotic pathogens are nucleated cells who are more closely related to you than they are to bacteria
 - * Malaria, various worms

Influenza virus



Tuberculosis bacilli



Malaria sporozoite



 For infections with small pathogens (viruses and bacteria), we don't attempt to count pathogens, but instead divide disease into stages

- For infections with small pathogens (viruses and bacteria), we don't attempt to count pathogens, but instead divide disease into stages
 - ► Latently infected

- For infections with small pathogens (viruses and bacteria), we don't attempt to count pathogens, but instead divide disease into stages
 - Latently infected
 - Productively infected

- For infections with small pathogens (viruses and bacteria), we don't attempt to count pathogens, but instead divide disease into stages
 - Latently infected
 - Productively infected
 - Recovered

- For infections with small pathogens (viruses and bacteria), we don't attempt to count pathogens, but instead divide disease into stages
 - Latently infected
 - Productively infected
 - Recovered

We model microparasites by counting the number of hosts in various states:

- We model microparasites by counting the number of hosts in various states:
 - Susceptible individuals can become infected

- We model microparasites by counting the number of hosts in various states:
 - Susceptible individuals can become infected
 - Infectious individuals are infected and can infect others

- We model microparasites by counting the number of hosts in various states:
 - Susceptible individuals can become infected
 - ▶ Infectious individuals are infected and can infect others
 - Resistant individuals are not infected and cannot become infected

- We model microparasites by counting the number of hosts in various states:
 - Susceptible individuals can become infected
 - ▶ Infectious individuals are infected and can infect others
 - Resistant individuals are not infected and cannot become infected
- More complicated models might include other states, such as latently infected hosts who are infected with the pathogen but cannot yet infect others


- We model microparasites by counting the number of hosts in various states:
 - Susceptible individuals can become infected
 - ▶ Infectious individuals are infected and can infect others
 - Resistant individuals are not infected and cannot become infected
- More complicated models might include other states, such as latently infected hosts who are infected with the pathogen but cannot yet infect others

Models as tools

► Models are the tools that we use to connect scales:



Measles reports from England and Wales




Models as tools

- Models are the tools that we use to connect scales:
 - ► individuals to populations



Measles reports from England and Wales




Models as tools

- Models are the tools that we use to connect scales:
 - individuals to populations
 - single actions to trends through time



Measles reports from England and Wales




Models as tools

- Models are the tools that we use to connect scales:
 - individuals to populations
 - single actions to trends through time



Measles reports from England and Wales



Outline

Introduction

Rate of spread

Single-epidemic model Epidemic size

Recurrent epidemic models

Reproductive numbers and risk

► Poll: For many diseases, especially new diseases, we can observe and estimate r.

► Poll: For many diseases, especially new diseases, we can observe and estimate r.

▶ 3

- ► Poll: For many diseases, especially new diseases, we can observe and estimate r.
 - * the exponential rate of spread

- ► Poll: For many diseases, especially new diseases, we can observe and estimate r.
 - * the exponential rate of spread
- ▶ Poll: Want to know what factors contribute to that, and how it relates to R.

- ▶ Poll: For many diseases, especially new diseases, we can observe and estimate r.
 - * the exponential rate of spread
- Poll: Want to know what factors contribute to that, and how it relates to R.

•

- Poll: For many diseases, especially new diseases, we can observe and estimate r.
 - * the exponential rate of spread
- Poll: Want to know what factors contribute to that, and how it relates to R.
 - ▶ * number of new cases per case

- Poll: For many diseases, especially new diseases, we can observe and estimate r.
 - * the exponential rate of spread
- Poll: Want to know what factors contribute to that, and how it relates to R.
 - ▶ * number of new cases per case

 \blacktriangleright People in the disease field love to talk specifically about \mathcal{R}_0

- lacktriangle People in the disease field love to talk specifically about \mathcal{R}_0
- ▶ But they don't always mean the same thing:

- lacktriangle People in the disease field love to talk specifically about \mathcal{R}_0
- But they don't always mean the same thing:
 - Actual value of R before an epidemic


- lacktriangle People in the disease field love to talk specifically about \mathcal{R}_0
- But they don't always mean the same thing:
 - ightharpoonup Actual value of $\mathcal R$ before an epidemic
 - Hypothetical value assuming no immunity

- People in the disease field love to talk specifically about R₀
- But they don't always mean the same thing:
 - ightharpoonup Actual value of $\mathcal R$ before an epidemic
 - Hypothetical value assuming no immunity
 - Hypothetical value assuming no control efforts whatsoever


- lacktriangle People in the disease field love to talk specifically about \mathcal{R}_0
- But they don't always mean the same thing:
 - Actual value of R before an epidemic
 - Hypothetical value assuming no immunity
 - Hypothetical value assuming no control efforts whatsoever
- ▶ Often easier to talk simply about \mathcal{R} .

- People in the disease field love to talk specifically about R₀
- But they don't always mean the same thing:
 - Actual value of R before an epidemic
 - Hypothetical value assuming no immunity
 - Hypothetical value assuming no control efforts whatsoever
- ▶ Often easier to talk simply about \mathcal{R} .


Example: the West African Ebola epidemic




 Researchers try to estimate the proportion of transmission that happens for different ages of infection



- Researchers try to estimate the proportion of transmission that happens for different ages of infection
- How long from the time you are infected to the time you infect someone else?





- Researchers try to estimate the proportion of transmission that happens for different ages of infection
- How long from the time you are infected to the time you infect someone else?
- ► Analogous to a life table



- Researchers try to estimate the proportion of transmission that happens for different ages of infection
- How long from the time you are infected to the time you infect someone else?
- Analogous to a life table
- ► The effective generation time \hat{G} has units of time

- Researchers try to estimate the proportion of transmission that happens for different ages of infection
- How long from the time you are infected to the time you infect someone else?
- Analogous to a life table
- ► The effective generation time \hat{G} has units of time







► Which is more dangerous, a fast disease, or a slow disease?

Generation interval (days)


- Which is more dangerous, a fast disease, or a slow disease?
 - How are we measuring speed?




- Which is more dangerous, a fast disease, or a slow disease?
 - How are we measuring speed?
 - How are we measuring danger?



- Which is more dangerous, a fast disease, or a slow disease?
 - How are we measuring speed?
 - How are we measuring danger?
 - What do we already know?



- Which is more dangerous, a fast disease, or a slow disease?
 - How are we measuring speed?
 - How are we measuring danger?
 - What do we already know?



Fighting Ebola



▶ If we know \mathcal{R} , what does the generation time tell us about r?

If we know R, what does the generation time tell us about r?

▶ '

- If we know R, what does the generation time tell us about r?
 - ▶ * The faster the generations (small \hat{G}), the faster the exponential growth (large r)


- If we know \mathcal{R} , what does the generation time tell us about r?
 - ▶ * The faster the generations (small \hat{G}), the faster the exponential growth (large r)
- ▶ If we know r, what does the generation time tell us about \mathcal{R} ?


- If we know R, what does the generation time tell us about r?
 - ▶ * The faster the generations (small \hat{G}), the faster the exponential growth (large r)
- If we know r, what does the generation time tell us about R?
 - .


- If we know \mathcal{R} , what does the generation time tell us about r?
 - ▶ * The faster the generations (small \hat{G}), the faster the exponential growth (large r)
- If we know r, what does the generation time tell us about R?
 - ▶ * The faster the generations (small \hat{G}), the the *smaller* the strength of the epidemic (small reproductive number \mathcal{R})

- If we know \mathcal{R} , what does the generation time tell us about r?
 - ▶ * The faster the generations (small \hat{G}), the faster the exponential growth (large r)
- If we know r, what does the generation time tell us about \mathcal{R} ?
 - ▶ * The faster the generations (small \hat{G}), the the *smaller* the strength of the epidemic (small reproductive number \mathcal{R})
- $\mathcal{R} = \exp(r\hat{G})$

- If we know \mathcal{R} , what does the generation time tell us about r?
 - ▶ * The faster the generations (small \hat{G}), the faster the exponential growth (large r)
- If we know r, what does the generation time tell us about \mathcal{R} ?
 - ▶ * The faster the generations (small \hat{G}), the the *smaller* the strength of the epidemic (small reproductive number \mathcal{R})
- $ightharpoonup \mathcal{R} = \exp(r\hat{G})$







► An intuitive view:

- An intuitive view:
 - ► Epidemic speed = Generation speed × Generation strength

- An intuitive view:
 - ► Epidemic speed = Generation speed × Generation strength
- If we know generation speed, then a faster epidemic speed means:

- An intuitive view:
 - ► Epidemic speed = Generation speed × Generation strength
- If we know generation speed, then a faster epidemic speed means:

*

- An intuitive view:
 - ► Epidemic speed = Generation speed × Generation strength
- If we know generation speed, then a faster epidemic speed means:
 - ▶ * More strength required (greater R)

- An intuitive view:
 - ► Epidemic speed = Generation speed × Generation strength
- If we know generation speed, then a faster epidemic speed means:
 - ★ * More strength required (greater R)
- If we know epidemic speed, a faster generation speed means

- An intuitive view:
 - ► Epidemic speed = Generation speed × Generation strength
- If we know generation speed, then a faster epidemic speed means:
 - ★ * More strength required (greater R)
- If we know epidemic speed, a faster generation speed means
 - ×

- An intuitive view:
 - ► Epidemic speed = Generation speed × Generation strength
- If we know generation speed, then a faster epidemic speed means:
 - ★ * More strength required (greater R)
- If we know epidemic speed, a faster generation speed means
 - ▶ * Less strength required (smaller R)

- An intuitive view:
 - ► Epidemic speed = Generation speed × Generation strength
- If we know generation speed, then a faster epidemic speed means:
 - ★ * More strength required (greater R)
- If we know epidemic speed, a faster generation speed means
 - ▶ * Less strength required (smaller R)

Outline

Introduction


Rate of spread

Single-epidemic model Epidemic size

Recurrent epidemic models


Reproductive numbers and risk

Single-epidemic model




 $\blacktriangleright \ \, \text{Susceptible} \to \text{Infectious} \to \text{Recovered}$

Single-epidemic model




- ightharpoonup Susceptible ightarrow Infectious ightarrow Recovered
- ▶ We also use *N* to mean the total population


Single-epidemic model




- ightharpoonup Susceptible ightarrow Infectious ightarrow Recovered
- ▶ We also use *N* to mean the total population




► What factors govern movement through the boxes?



- What factors govern movement through the boxes?
 - ► People get better independently



- What factors govern movement through the boxes?
 - People get better independently
 - ► People get infected by infectious people



- What factors govern movement through the boxes?
 - People get better independently
 - People get infected by infectious people

Poll: What happens in the long term if we introduce an infectious individual?




▶ Poll: What happens in the long term if we introduce an infectious individual?

,



- ▶ Poll: What happens in the long term if we introduce an infectious individual?
 - ▶ * The *may be* an **epidemic**
 - an outbreak of disease



- Poll: What happens in the long term if we introduce an infectious individual?
 - ▶ * The *may be* an **epidemic**
 - an outbreak of disease

. 9



- Poll: What happens in the long term if we introduce an infectious individual?
 - ▶ * The *may be* an **epidemic**
 - an outbreak of disease
 - * Disease burns out



- ▶ Poll: What happens in the long term if we introduce an infectious individual?
 - * The may be an epidemic– an outbreak of disease
 - an outbreak of disease
 - * Disease burns out

.



- Poll: What happens in the long term if we introduce an infectious individual?
 - * The may be an epidemic– an outbreak of disease
 - * Disease burns out
 - * Everyone winds up either recovered or susceptible



- ▶ Poll: What happens in the long term if we introduce an infectious individual?
 - * The may be an epidemic– an outbreak of disease
 - * Disease burns out
 - * Everyone winds up either recovered or susceptible

•



- ▶ Poll: What happens in the long term if we introduce an infectious individual?
 - * The may be an epidemic– an outbreak of disease
 - * Disease burns out
 - * Everyone winds up either recovered or susceptible
 - * Not everyone gets infected!



- ▶ Poll: What happens in the long term if we introduce an infectious individual?
 - * The may be an epidemic– an outbreak of disease
 - * Disease burns out
 - * Everyone winds up either recovered or susceptible
 - * Not everyone gets infected!



Interpreting

► Why might there not be an epidemic?

Interpreting

Why might there not be an epidemic?

Interpreting

- Why might there not be an epidemic?
 - * If the disease can't spread well enough in the population

- Why might there not be an epidemic?
 - * If the disease can't spread well enough in the population
 - •

- Why might there not be an epidemic?
 - * If the disease can't spread well enough in the population
 - ► * Demographic stochasticity: if we only start with one individual, we expect an element of chance

- Why might there not be an epidemic?
 - * If the disease can't spread well enough in the population
 - ► * Demographic stochasticity: if we only start with one individual, we expect an element of chance
- ▶ Why doesn't everyone get infected?

- Why might there not be an epidemic?
 - * If the disease can't spread well enough in the population
 - ► * Demographic stochasticity: if we only start with one individual, we expect an element of chance
- Why doesn't everyone get infected?

► The simplest way to implement this conceptual model is with differential equations:



The simplest way to implement this conceptual model is with differential equations:

$$\frac{dS}{dt} = -\beta \frac{SI}{N}$$



The simplest way to implement this conceptual model is with differential equations:

$$\frac{dS}{dt} = -\beta \frac{SI}{N}$$

•

$$\frac{dI}{dt} = \beta \frac{SI}{N} - \gamma I$$



The simplest way to implement this conceptual model is with differential equations:

$$\frac{dS}{dt} = -\beta \frac{SI}{N}$$

$$\frac{dI}{dt} = \beta \frac{SI}{N} - \gamma I$$

$$\frac{dR}{dt} = \gamma I$$





The simplest way to implement this conceptual model is with differential equations:

$$\frac{dS}{dt} = -\beta \frac{SI}{N}$$

$$\frac{dI}{dt} = \beta \frac{SI}{N} - \gamma I$$


$$\frac{dR}{dt} = \gamma I$$





State variables

► S, I, R, N: [people] or [people/ha]



State variables

► S, I, R, N: [people] or [people/ha]

Parameters

 Susceptible people have potentially effective contacts at rate β (units [1/time])

- Susceptible people have potentially effective contacts at rate β (units [1/time])
 - These are contacts that would lead to infection if the person contacted is infectious

- Susceptible people have potentially effective contacts at rate β (units [1/time])
 - These are contacts that would lead to infection if the person contacted is infectious
 - Total infection rate is βI/N, because I/N is the proportion of the population infectious


- Susceptible people have potentially effective contacts at rate β (units [1/time])
 - These are contacts that would lead to infection if the person contacted is infectious
 - ▶ Total infection rate is $\beta I/N$, because I/N is the proportion of the population infectious
- ► Infectious people recover at *per capita* rate γ (units [1/time])

- Susceptible people have potentially effective contacts at rate β (units [1/time])
 - These are contacts that would lead to infection if the person contacted is infectious
 - ▶ Total infection rate is $\beta I/N$, because I/N is the proportion of the population infectious
- Infectious people recover at *per capita* rate γ (units [1/time])
 - ► Total recovery rate is γI


- Susceptible people have potentially effective contacts at rate β (units [1/time])
 - These are contacts that would lead to infection if the person contacted is infectious
 - ▶ Total infection rate is $\beta I/N$, because I/N is the proportion of the population infectious
- Infectious people recover at *per capita* rate γ (units [1/time])
 - ▶ Total recovery rate is ~I
 - ▶ Mean time infectious is $D = 1/\gamma$ (units [time])


- Susceptible people have potentially effective contacts at rate β (units [1/time])
 - These are contacts that would lead to infection if the person contacted is infectious
 - ▶ Total infection rate is $\beta I/N$, because I/N is the proportion of the population infectious
- Infectious people recover at *per capita* rate γ (units [1/time])
 - ▶ Total recovery rate is ~I
 - ▶ Mean time infectious is $D = 1/\gamma$ (units [time])

Simulating the model



Simulating the model





► Poll: What *unitless* parameter can you make from the model above?

Poll: What unitless parameter can you make from the model above?

•

- Poll: What unitless parameter can you make from the model above?
 - * $\mathcal{R}_0 = \beta D = \beta/\gamma$ is the basic reproductive number

- Poll: What unitless parameter can you make from the model above?
 - * $\mathcal{R}_0 = \beta D = \beta/\gamma$ is the basic reproductive number

- Poll: What unitless parameter can you make from the model above?
 - * $\mathcal{R}_0 = \beta D = \beta/\gamma$ is the basic reproductive number
 - * The potential number of infections caused by an average infectious individual

- Poll: What unitless parameter can you make from the model above?
 - * $\mathcal{R}_0 = \beta D = \beta/\gamma$ is the basic reproductive number
 - * The potential number of infections caused by an average infectious individual



- Poll: What unitless parameter can you make from the model above?
 - * $\mathcal{R}_0 = \beta D = \beta/\gamma$ is the basic reproductive number
 - * The potential number of infections caused by an average infectious individual
 - * That is: the number they would cause on average if everyone else were susceptible

- Poll: What unitless parameter can you make from the model above?
 - * $\mathcal{R}_0 = \beta D = \beta/\gamma$ is the basic reproductive number
 - * The potential number of infections caused by an average infectious individual
 - * That is: the number they would cause on average if everyone else were susceptible

▶ Poll: What happens early in the epidemic if $\mathcal{R}_0 > 1$?

▶ Poll: What happens early in the epidemic if $\mathcal{R}_0 > 1$?

- ▶ Poll: What happens early in the epidemic if $\mathcal{R}_0 > 1$?
 - * Number of infected individuals grows exponentially

- ▶ Poll: What happens early in the epidemic if $\mathcal{R}_0 > 1$?
 - * Number of infected individuals grows exponentially
- ▶ What happens early in the epidemic if $\mathcal{R}_0 < 1$?

- ▶ Poll: What happens early in the epidemic if $\mathcal{R}_0 > 1$?
 - * Number of infected individuals grows exponentially
- ▶ What happens early in the epidemic if $\mathcal{R}_0 < 1$?

- ▶ Poll: What happens early in the epidemic if $\mathcal{R}_0 > 1$?
 - * Number of infected individuals grows exponentially
- ▶ What happens early in the epidemic if $\mathcal{R}_0 < 1$?
 - * Number of infected individuals cannot grow (disease cannot invade)

- ▶ Poll: What happens early in the epidemic if $\mathcal{R}_0 > 1$?
 - * Number of infected individuals grows exponentially
- ▶ What happens early in the epidemic if $\mathcal{R}_0 < 1$?
 - * Number of infected individuals cannot grow (disease cannot invade)

Effective reproductive number

► The effective reproductive number gives the number of new infections per infectious individual in a partially susceptible population:

The effective reproductive number gives the number of new infections per infectious individual in a partially susceptible population:

→ ³

• *
$$\mathcal{R}_e = \mathcal{R}_0 S/N$$

The effective reproductive number gives the number of new infections per infectious individual in a partially susceptible population:

• *
$$\mathcal{R}_e = \mathcal{R}_0 S/N$$

Is the disease increasing or decreasing?

The effective reproductive number gives the number of new infections per infectious individual in a partially susceptible population:

• *
$$\mathcal{R}_e = \mathcal{R}_0 S/N$$

Is the disease increasing or decreasing?

• *
$$\mathcal{R}_e = \mathcal{R}_0 S/N$$

- Is the disease increasing or decreasing?
 - ▶ * It will increase when $\mathcal{R}_e > 1$ (more than one case ber case)

• *
$$\mathcal{R}_e = \mathcal{R}_0 S/N$$

- Is the disease increasing or decreasing?
 - ▶ * It will increase when $\mathcal{R}_e > 1$ (more than one case ber case)
 - *

• *
$$\mathcal{R}_e = \mathcal{R}_0 S/N$$

- Is the disease increasing or decreasing?
 - ▶ * It will increase when $R_e > 1$ (more than one case ber case)
 - ▶ * This happens when $S/N > 1/\mathcal{R}_0$

• *
$$\mathcal{R}_e = \mathcal{R}_0 S/N$$

- Is the disease increasing or decreasing?
 - ▶ * It will increase when $R_e > 1$ (more than one case ber case)
 - ▶ * This happens when $S/N > 1/\mathcal{R}_0$
- Why doesn't everyone get infected?

• *
$$\mathcal{R}_e = \mathcal{R}_0 S/N$$

- Is the disease increasing or decreasing?
 - ▶ * It will increase when $R_e > 1$ (more than one case ber case)
 - ▶ * This happens when $S/N > 1/\mathcal{R}_0$
- Why doesn't everyone get infected?



• *
$$\mathcal{R}_e = \mathcal{R}_0 S/N$$

- Is the disease increasing or decreasing?
 - ▶ * It will increase when $R_e > 1$ (more than one case ber case)
 - ▶ * This happens when $S/N > 1/\mathcal{R}_0$
- Why doesn't everyone get infected?
 - ▶ * When susceptibles are low enough R_e < 1

• *
$$\mathcal{R}_e = \mathcal{R}_0 S/N$$

- Is the disease increasing or decreasing?
 - ▶ * It will increase when $R_e > 1$ (more than one case ber case)
 - ▶ * This happens when $S/N > 1/R_0$
- Why doesn't everyone get infected?
 - ▶ * When susceptibles are low enough R_e < 1
 - •

• *
$$\mathcal{R}_e = \mathcal{R}_0 S/N$$

- Is the disease increasing or decreasing?
 - ▶ * It will increase when $R_e > 1$ (more than one case ber case)
 - ▶ * This happens when $S/N > 1/R_0$
- Why doesn't everyone get infected?
 - ▶ * When susceptibles are low enough R_e < 1
 - $\,\blacktriangleright\,$ * When $\mathcal{R}_e <$ 1, the disease dies out on its own (less than one case ber case)

• *
$$\mathcal{R}_e = \mathcal{R}_0 S/N$$

- Is the disease increasing or decreasing?
 - ▶ * It will increase when $R_e > 1$ (more than one case ber case)
 - ▶ * This happens when $S/N > 1/R_0$
- Why doesn't everyone get infected?
 - ▶ * When susceptibles are low enough R_e < 1
 - $\,\blacktriangleright\,$ * When $\mathcal{R}_e <$ 1, the disease dies out on its own (less than one case ber case)

Subsection 1

► In this model, the epidemic always burns out

- In this model, the epidemic always burns out
 - ► No source of new susceptibles

- In this model, the epidemic always burns out
 - ▶ No source of new susceptibles
- ► Epidemic size is determined by:

- In this model, the epidemic always burns out
 - No source of new susceptibles
- Epidemic size is determined by:

•


- In this model, the epidemic always burns out
 - No source of new susceptibles
- Epidemic size is determined by:
 - ▶ * \mathcal{R}_0 larger \mathcal{R}_0 leads to a bigger epidemic

- In this model, the epidemic always burns out
 - No source of new susceptibles
- Epidemic size is determined by:
 - ▶ * \mathcal{R}_0 larger \mathcal{R}_0 leads to a bigger epidemic
 - •

- In this model, the epidemic always burns out
 - No source of new susceptibles
- Epidemic size is determined by:
 - ▶ * \mathcal{R}_0 larger \mathcal{R}_0 leads to a bigger epidemic
 - * The number of susceptibles at the beginning of the epidemic


- In this model, the epidemic always burns out
 - No source of new susceptibles
- Epidemic size is determined by:
 - ▶ * \mathcal{R}_0 larger \mathcal{R}_0 leads to a bigger epidemic
 - * The number of susceptibles at the beginning of the epidemic

Why does more susceptibles at the beginning mean fewer susceptibles at the end?




Why does more susceptibles at the beginning mean fewer susceptibles at the end?

, ·




- Why does more susceptibles at the beginning mean fewer susceptibles at the end?
 - ▶ * Bigger epidemic ⇒⇒




Why does more susceptibles at the beginning mean fewer susceptibles at the end?


* Bigger epidemic ⇒



- Why does more susceptibles at the beginning mean fewer susceptibles at the end?
 - ▶ * Bigger epidemic ⇒⇒
 - ★ More infections at peak (same number of susceptibles) ⇒




- Why does more susceptibles at the beginning mean fewer susceptibles at the end?
 - ▶ * Bigger epidemic ⇒⇒
 - * More infections at peak (same number of susceptibles)
 - .




- Why does more susceptibles at the beginning mean fewer susceptibles at the end?
 - ▶ * Bigger epidemic ⇒
 - * More infections at peak (same number of susceptibles)
 - * More infections after peak

. . .




- Why does more susceptibles at the beginning mean fewer susceptibles at the end?
 - ▶ * Bigger epidemic ⇒
 - * More infections at peak (same number of susceptibles)
 - * More infections after peak


. . .




► In September, the US CDC predicted "as many as" 1.5 million Ebola cases in Liberia by January




- In September, the US CDC predicted "as many as" 1.5 million Ebola cases in Liberia by January
- In fact, their model predicted many more cases than that by April



- In September, the US CDC predicted "as many as" 1.5 million Ebola cases in Liberia by January
- In fact, their model predicted many more cases than that by April
- ► What happened?



- In September, the US CDC predicted "as many as" 1.5 million Ebola cases in Liberia by January
- In fact, their model predicted many more cases than that by April
- ▶ What happened?



What limits epidemics?

▶ Poll: What limits epidemics in our simple models?

What limits epidemics?

► Poll: What limits epidemics in our simple models?

- ▶ Poll: What limits epidemics in our simple models?
 - * Depletion of susceptibles

- ▶ Poll: What limits epidemics in our simple models?
 - * Depletion of susceptibles
- ▶ Poll: What else limits epidemics in real life?

- Poll: What limits epidemics in our simple models?
 - * Depletion of susceptibles
- ▶ Poll: What else limits epidemics in real life?

▶ `

- Poll: What limits epidemics in our simple models?
 - * Depletion of susceptibles
- ▶ Poll: What else limits epidemics in real life?
 - * Interventions

- Poll: What limits epidemics in our simple models?
 - * Depletion of susceptibles
- ▶ Poll: What else limits epidemics in real life?
 - * Interventions
 - •

- Poll: What limits epidemics in our simple models?
 - * Depletion of susceptibles
- Poll: What else limits epidemics in real life?
 - * Interventions
 - * Behaviour change

- Poll: What limits epidemics in our simple models?
 - * Depletion of susceptibles
- Poll: What else limits epidemics in real life?
 - * Interventions
 - * Behaviour change

- ▶ Poll: What limits epidemics in our simple models?
 - * Depletion of susceptibles
- ▶ Poll: What else limits epidemics in real life?
 - * Interventions
 - ▶ * Behaviour change
 - * Heterogeneity (differences between hosts, locations, etc.)

- ▶ Poll: What limits epidemics in our simple models?
 - * Depletion of susceptibles
- ▶ Poll: What else limits epidemics in real life?
 - * Interventions
 - ▶ * Behaviour change
 - * Heterogeneity (differences between hosts, locations, etc.)

Outline

Introduction


Rate of spread

Single-epidemic model Epidemic size

Recurrent epidemic models


Reproductive numbers and risk

Poll: If epidemics tend to burn out, why do we often see repeated epidemics?




Poll: If epidemics tend to burn out, why do we often see repeated epidemics?

,




- Poll: If epidemics tend to burn out, why do we often see repeated epidemics?
 - * People might lose immunity




- Poll: If epidemics tend to burn out, why do we often see repeated epidemics?
 - * People might lose immunity


,




- Poll: If epidemics tend to burn out, why do we often see repeated epidemics?
 - * People might lose immunity
 - * Births and deaths





- Poll: If epidemics tend to burn out, why do we often see repeated epidemics?
 - * People might lose immunity
 - * Births and deaths




Recurrent epidemics









* Loss of immunity




* Loss of immunity







▶ * Births and deaths







- * Births and deaths
 - * Effect on dynamics is similar to loss of immunity



- * Births and deaths
 - * Effect on dynamics is similar to loss of immunity

$$\frac{dS}{dt} = bN - \beta \frac{SI}{N} - dS$$



$$\frac{dS}{dt} = bN - \beta \frac{SI}{N} - dS$$

$$\frac{dI}{dt} = \beta \frac{SI}{N} - \gamma I - dI$$



$$\frac{dS}{dt} = bN - \beta \frac{SI}{N} - dS$$

$$\frac{dI}{dt} = \beta \frac{SI}{N} - \gamma I - dI$$

$$\frac{dR}{dt} = \gamma I - dR$$



$$\frac{dS}{dt} = bN - \beta \frac{SI}{N} - dS$$

$$\frac{dI}{dt} = \beta \frac{SI}{N} - \gamma I - dI$$



$$\frac{dR}{dt} = \gamma I - dR$$

▶ We often assume b = d



$$\frac{dS}{dt} = bN - \beta \frac{SI}{N} - dS$$

$$\frac{dI}{dt} = \beta \frac{SI}{N} - \gamma I - dI$$

 $\begin{array}{c|c} \downarrow & & \\ \hline s & & \\ \downarrow & & \\ \end{array}$

$$\frac{dR}{dt} = \gamma I - dR$$

- ▶ We often assume b = d
 - population is constant

$$\frac{dS}{dt} = bN - \beta \frac{SI}{N} - dS$$

$$\frac{dI}{dt} = \beta \frac{SI}{N} - \gamma I - dI$$

$$S \longrightarrow I \longrightarrow R$$

$$\frac{dR}{dt} = \gamma I - dR$$

- ▶ We often assume b = d
 - population is constant

▶ At equilibrium, we know that $R_e = 1$

- ▶ At equilibrium, we know that $\mathcal{R}_e = 1$
 - ▶ One case per case

- ▶ At equilibrium, we know that $\mathcal{R}_e = 1$
 - ► One case per case
 - Number of susceptibles at equilibrium determined by the number required to keep infection in balance

- ▶ At equilibrium, we know that $\mathcal{R}_e = 1$
 - One case per case
 - Number of susceptibles at equilibrium determined by the number required to keep infection in balance
 - ► $S/N = 1/\mathcal{R}_0$

- ▶ At equilibrium, we know that $\mathcal{R}_e = 1$
 - One case per case
 - Number of susceptibles at equilibrium determined by the number required to keep infection in balance
 - ▶ $S/N = 1/R_0$

.

- ▶ At equilibrium, we know that $\mathcal{R}_e = 1$
 - One case per case
 - Number of susceptibles at equilibrium determined by the number required to keep infection in balance
 - ▶ $S/N = 1/\mathcal{R}_0$
 - * Reciprocal control!

- ▶ At equilibrium, we know that $\mathcal{R}_e = 1$
 - One case per case
 - Number of susceptibles at equilibrium determined by the number required to keep infection in balance
 - ▶ $S/N = 1/\mathcal{R}_0$
 - * Reciprocal control!

Number of infectious individuals determined by number required to keep susceptibles in balance.

- Number of infectious individuals determined by number required to keep susceptibles in balance.
- As susceptibles go up, what happens?

- Number of infectious individuals determined by number required to keep susceptibles in balance.
- As susceptibles go up, what happens?
 - Per capita replenishment goes down

- Number of infectious individuals determined by number required to keep susceptibles in balance.
- As susceptibles go up, what happens?
 - Per capita replenishment goes down
 - Infections required goes down

- Number of infectious individuals determined by number required to keep susceptibles in balance.
- As susceptibles go up, what happens?
 - Per capita replenishment goes down
 - Infections required goes down

► What happens if we protect susceptibles (move them to *R* class)?

What happens if we protect susceptibles (move them to R class)?

4□ > 4□ > 4 = > 4 = > = 900

- What happens if we protect susceptibles (move them to R class)?
 - ► * Equation for *dI/dt* does not change

- What happens if we protect susceptibles (move them to R class)?
 - ▶ * Equation for *dI/dt* does not change
 - •

- What happens if we protect susceptibles (move them to R class)?
 - * Equation for dI/dt does not change
 - * Number of susceptibles does not change

- What happens if we protect susceptibles (move them to R class)?
 - * Equation for dI/dt does not change
 - * Number of susceptibles does not change
 - → ³

- What happens if we protect susceptibles (move them to R class)?
 - * Equation for dI/dt does not change
 - * Number of susceptibles does not change
 - * Fewer susceptibles need to be removed by infection (some are removed by us)

- What happens if we protect susceptibles (move them to R class)?
 - * Equation for dI/dt does not change
 - * Number of susceptibles does not change
 - * Fewer susceptibles need to be removed by infection (some are removed by us)
 - •

- What happens if we protect susceptibles (move them to R class)?
 - * Equation for dI/dt does not change
 - * Number of susceptibles does not change
 - * Fewer susceptibles need to be removed by infection (some are removed by us)
 - * Number of infectious individuals goes down

- What happens if we protect susceptibles (move them to R class)?
 - ▶ * Equation for dI/dt does not change
 - * Number of susceptibles does not change
 - * Fewer susceptibles need to be removed by infection (some are removed by us)
 - * Number of infectious individuals goes down
- What else could happen?

- What happens if we protect susceptibles (move them to R class)?
 - ▶ * Equation for dI/dt does not change
 - * Number of susceptibles does not change
 - * Fewer susceptibles need to be removed by infection (some are removed by us)
 - * Number of infectious individuals goes down
- What else could happen?

- What happens if we protect susceptibles (move them to R class)?
 - * Equation for dI/dt does not change
 - * Number of susceptibles does not change
 - * Fewer susceptibles need to be removed by infection (some are removed by us)
 - * Number of infectious individuals goes down
- What else could happen?
 - * If we remove susceptibles fast enough, infection could go extinct

- What happens if we protect susceptibles (move them to R class)?
 - * Equation for dI/dt does not change
 - * Number of susceptibles does not change
 - * Fewer susceptibles need to be removed by infection (some are removed by us)
 - * Number of infectious individuals goes down
- What else could happen?
 - * If we remove susceptibles fast enough, infection could go extinct
 - *

- What happens if we protect susceptibles (move them to R class)?
 - * Equation for dI/dt does not change
 - * Number of susceptibles does not change
 - * Fewer susceptibles need to be removed by infection (some are removed by us)
 - * Number of infectious individuals goes down
- What else could happen?
 - * If we remove susceptibles fast enough, infection could go extinct
 - * If we keep increasing the rate . . .

- What happens if we protect susceptibles (move them to R class)?
 - * Equation for dI/dt does not change
 - * Number of susceptibles does not change
 - * Fewer susceptibles need to be removed by infection (some are removed by us)
 - * Number of infectious individuals goes down
- What else could happen?
 - * If we remove susceptibles fast enough, infection could go extinct
 - * If we keep increasing the rate . . .

- What happens if we protect susceptibles (move them to R class)?
 - * Equation for dI/dt does not change
 - * Number of susceptibles does not change
 - * Fewer susceptibles need to be removed by infection (some are removed by us)
 - * Number of infectious individuals goes down
- What else could happen?
 - * If we remove susceptibles fast enough, infection could go extinct
 - * If we keep increasing the rate . . .
 - * Number of susceptibles goes down

- What happens if we protect susceptibles (move them to R class)?
 - * Equation for dI/dt does not change
 - * Number of susceptibles does not change
 - * Fewer susceptibles need to be removed by infection (some are removed by us)
 - * Number of infectious individuals goes down
- What else could happen?
 - * If we remove susceptibles fast enough, infection could go extinct
 - * If we keep increasing the rate . . .
 - * Number of susceptibles goes down

► Poll: What happens if we remove infectious individuals at a constant rate (find them and cure them or isolate them)?

► Poll: What happens if we remove infectious individuals at a constant rate (find them and cure them or isolate them)?

▶ ′

- ► Poll: What happens if we remove infectious individuals at a constant rate (find them and cure them or isolate them)?
 - ▶ * We need more susceptibles to balance dI/dt

- ► Poll: What happens if we remove infectious individuals at a constant rate (find them and cure them or isolate them)?
 - ▶ * We need more susceptibles to balance dI/dt

4D > 4B > 4E > 4E > 9Q0

- ▶ Poll: What happens if we remove infectious individuals at a constant rate (find them and cure them or isolate them)?
 - * We need more susceptibles to balance dI/dt
 - * If we have more susceptibles, then per capita replenishment goes down

- ▶ Poll: What happens if we remove infectious individuals at a constant rate (find them and cure them or isolate them)?
 - ▶ * We need more susceptibles to balance dI/dt
 - * If we have more susceptibles, then per capita replenishment goes down

•


- ▶ Poll: What happens if we remove infectious individuals at a constant rate (find them and cure them or isolate them)?
 - * We need more susceptibles to balance dI/dt
 - * If we have more susceptibles, then per capita replenishment goes down
 - * So the number of infectious individuals goes down


- ▶ Poll: What happens if we remove infectious individuals at a constant rate (find them and cure them or isolate them)?
 - ▶ * We need more susceptibles to balance dI/dt
 - * If we have more susceptibles, then per capita replenishment goes down
 - * So the number of infectious individuals goes down

- ▶ Poll: What happens if we remove infectious individuals at a constant rate (find them and cure them or isolate them)?
 - * We need more susceptibles to balance dI/dt
 - * If we have more susceptibles, then per capita replenishment goes down
 - * So the number of infectious individuals goes down
 - * If we remove infectious individuals fast enough, the infection could go extinct

- ▶ Poll: What happens if we remove infectious individuals at a constant rate (find them and cure them or isolate them)?
 - * We need more susceptibles to balance dI/dt
 - * If we have more susceptibles, then per capita replenishment goes down
 - * So the number of infectious individuals goes down
 - * If we remove infectious individuals fast enough, the infection could go extinct

Tendency to oscillate





Tendency to oscillate

"Closed-loop" SIR models (ie., with births or loss of immunity):

- "Closed-loop" SIR models (ie., with births or loss of immunity):
 - ► Tend to oscillate

- "Closed-loop" SIR models (ie., with births or loss of immunity):
 - ► Tend to oscillate
 - Oscillations tend to be damped

- "Closed-loop" SIR models (ie., with births or loss of immunity):
 - Tend to oscillate
 - Oscillations tend to be damped
 - System reaches an endemic equilibrium disease persists

- "Closed-loop" SIR models (ie., with births or loss of immunity):
 - Tend to oscillate
 - Oscillations tend to be damped
 - System reaches an endemic equilibrium disease persists

Similar to predator-prey systems

- Similar to predator-prey systems
- ► What happens if we start with too many susceptibles?

- Similar to predator-prey systems
- What happens if we start with too many susceptibles?

- Similar to predator-prey systems
- What happens if we start with too many susceptibles?
 - * There will be a big epidemic

- Similar to predator-prey systems
- What happens if we start with too many susceptibles?
 - * There will be a big epidemic
 - •

- Similar to predator-prey systems
- What happens if we start with too many susceptibles?
 - ▶ * There will be a big epidemic
 - * . . . then a very low number of susceptibles

- Similar to predator-prey systems
- What happens if we start with too many susceptibles?
 - * There will be a big epidemic
 - * . . . then a very low number of susceptibles
 - •


- Similar to predator-prey systems
- What happens if we start with too many susceptibles?
 - * There will be a big epidemic
 - * . . . then a very low number of susceptibles
 - * ...then a very low level of disease

- Similar to predator-prey systems
- What happens if we start with too many susceptibles?
 - * There will be a big epidemic
 - * . . . then a very low number of susceptibles
 - ▶ * ... then a very low level of disease
 - •

- Similar to predator-prey systems
- What happens if we start with too many susceptibles?
 - ▶ * There will be a big epidemic
 - * ... then a very low number of susceptibles
 - * ... then a very low level of disease
 - * ... then an increase in the number of susceptibles


- Similar to predator-prey systems
- What happens if we start with too many susceptibles?
 - ▶ * There will be a big epidemic
 - * ... then a very low number of susceptibles
 - * ... then a very low level of disease
 - * ... then an increase in the number of susceptibles

► Poll: If oscillations tend to be damped in simple models, why do they persist in real life?




Poll: If oscillations tend to be damped in simple models, why do they persist in real life?


*




- Poll: If oscillations tend to be damped in simple models, why do they persist in real life?
 - ▶ * Weather




- Poll: If oscillations tend to be damped in simple models, why do they persist in real life?
 - * Weather
 - *




- Poll: If oscillations tend to be damped in simple models, why do they persist in real life?
 - * Weather
 - * School terms




- Poll: If oscillations tend to be damped in simple models, why do they persist in real life?
 - * Weather
 - * School terms
 - *




- Poll: If oscillations tend to be damped in simple models, why do they persist in real life?
 - * Weather
 - * School terms
 - * Demographic stochasticity



- Poll: If oscillations tend to be damped in simple models, why do they persist in real life?
 - * Weather
 - * School terms
 - * Demographic stochasticity




- Poll: If oscillations tend to be damped in simple models, why do they persist in real life?
 - * Weather
 - * School terms
 - * Demographic stochasticity
 - * Changes in Behaviour




- Poll: If oscillations tend to be damped in simple models, why do they persist in real life?
 - * Weather
 - * School terms
 - * Demographic stochasticity
 - * Changes in Behaviour


,



- Poll: If oscillations tend to be damped in simple models, why do they persist in real life?
 - * Weather
 - * School terms
 - * Demographic stochasticity
 - * Changes in Behaviour
 - * People are more careful when disease levels are high



- Poll: If oscillations tend to be damped in simple models, why do they persist in real life?
 - * Weather
 - * School terms
 - * Demographic stochasticity
 - * Changes in Behaviour
 - * People are more careful when disease levels are high



Outline

Introduction

Rate of spread

Single-epidemic model Epidemic size


Recurrent epidemic models

▶ At equilibrium, the proportion of people who are susceptible to disease should be approximately $S/N = 1/\mathcal{R}_0$

- At equilibrium, the proportion of people who are susceptible to disease should be approximately $S/N = 1/\mathcal{R}_0$
- ▶ Proportion "affected" (infectious or immune) should be approximately $V/N = 1 1/\mathcal{R}_0$

- ▶ At equilibrium, the proportion of people who are susceptible to disease should be approximately $S/N = 1/R_0$
- ▶ Proportion "affected" (infectious or immune) should be approximately $V/N = 1 1/R_0$
- ▶ If you have a single, fast epidemic, the size is also predicted by \mathcal{R}_0 .

- ▶ At equilibrium, the proportion of people who are susceptible to disease should be approximately $S/N = 1/R_0$
- ▶ Proportion "affected" (infectious or immune) should be approximately $V/N = 1 1/R_0$
- If you have a single, fast epidemic, the size is also predicted by \mathcal{R}_0 .



Examples

 Ronald Ross predicted 100 years ago that reducing mosquito densities by a factor of 5 or so would *eliminate* malaria

Examples

- Ronald Ross predicted 100 years ago that reducing mosquito densities by a factor of 5 or so would *eliminate* malaria
- Gradual disappearance of polio, typhoid, etc., without risk factors going to zero

Examples

- Ronald Ross predicted 100 years ago that reducing mosquito densities by a factor of 5 or so would *eliminate* malaria
- Gradual disappearance of polio, typhoid, etc., without risk factors going to zero
- Eradication of smallpox!

- Ronald Ross predicted 100 years ago that reducing mosquito densities by a factor of 5 or so would *eliminate* malaria
- Gradual disappearance of polio, typhoid, etc., without risk factors going to zero
- Eradication of smallpox!

► What proportion of the population should be vaccinated to eliminate a disease?

What proportion of the population should be vaccinated to eliminate a disease?

▶ '

- What proportion of the population should be vaccinated to eliminate a disease?
 - ▶ * Transmission should be reduced by a factor of \mathcal{R} , so a fraction $1 1/\mathcal{R}$ should be vaccinated

- What proportion of the population should be vaccinated to eliminate a disease?
 - ▶ * Transmission should be reduced by a factor of \mathcal{R} , so a fraction $1 1/\mathcal{R}$ should be vaccinated

▶ Polio has an \mathcal{R}_0 of about 5.

- ▶ Polio has an \mathcal{R}_0 of about 5.
- ► Poll: What proportion of the population should be vaccinated to eliminate polio?

- ▶ Polio has an \mathcal{R}_0 of about 5.
- Poll: What proportion of the population should be vaccinated to eliminate polio?
 - •

- ▶ Polio has an \mathcal{R}_0 of about 5.
- Poll: What proportion of the population should be vaccinated to eliminate polio?
 - ► * At least 1-1/5 = 80%

- Polio has an R₀ of about 5.
- Poll: What proportion of the population should be vaccinated to eliminate polio?
 - ► * At least 1-1/5 = 80%
- ► Measles has an R₀ of about 20. What proportion of the population should be vaccinated to eliminate measles?

- ▶ Polio has an \mathcal{R}_0 of about 5.
- Poll: What proportion of the population should be vaccinated to eliminate polio?
 - ► * At least 1-1/5 = 80%
- ► Measles has an R₀ of about 20. What proportion of the population should be vaccinated to eliminate measles?

- Polio has an R₀ of about 5.
- Poll: What proportion of the population should be vaccinated to eliminate polio?
 - ► * At least 1-1/5 = 80%
- ▶ Measles has an R₀ of about 20. What proportion of the population should be vaccinated to eliminate measles?
 - ► * At least 1-1/20 = 95%

- Polio has an R₀ of about 5.
- Poll: What proportion of the population should be vaccinated to eliminate polio?
 - ► * At least 1-1/5 = 80%
- Measles has an R₀ of about 20. What proportion of the population should be vaccinated to eliminate measles?
 - ► * At least 1-1/20 = 95%

Why have infectious diseases persisted?

- Why have infectious diseases persisted?
 - ► The pathogens evolve

- Why have infectious diseases persisted?
 - ► The pathogens evolve
 - ► Human populations are heterogeneous

- Why have infectious diseases persisted?
 - The pathogens evolve
 - Human populations are heterogeneous
 - People differ in: nutrition, exposure, access to care

- Why have infectious diseases persisted?
 - ▶ The pathogens evolve
 - Human populations are heterogeneous
 - People differ in: nutrition, exposure, access to care
 - Information and misinformation

- Why have infectious diseases persisted?
 - ▶ The pathogens evolve
 - Human populations are heterogeneous
 - People differ in: nutrition, exposure, access to care
 - Information and misinformation
 - Vaccine scares, trust in health care in general

- Why have infectious diseases persisted?
 - ▶ The pathogens evolve
 - Human populations are heterogeneous
 - People differ in: nutrition, exposure, access to care
 - Information and misinformation
 - Vaccine scares, trust in health care in general

▶ Heterogeneity *increases* \mathcal{R}_0

- ▶ Heterogeneity increases R₀
 - When disease is rare, it is concentrated in the most vulnerable populations

- Heterogeneity increases R₀
 - When disease is rare, it is concentrated in the most vulnerable populations
 - Cases per case is high

- ▶ Heterogeneity increases R₀
 - When disease is rare, it is concentrated in the most vulnerable populations
 - Cases per case is high
 - ► Elimination is harder

- Heterogeneity increases R₀
 - When disease is rare, it is concentrated in the most vulnerable populations
 - Cases per case is high
 - Elimination is harder
- Marginal populations

- ▶ Heterogeneity increases R₀
 - When disease is rare, it is concentrated in the most vulnerable populations
 - Cases per case is high
 - Elimination is harder
- Marginal populations
 - Heterogeneity could make it easier to concentrate on the most vulnerable populations and eliminate disease

- Heterogeneity increases R₀
 - When disease is rare, it is concentrated in the most vulnerable populations
 - Cases per case is high
 - ► Elimination is harder
- Marginal populations
 - Heterogeneity could make it easier to concentrate on the most vulnerable populations and eliminate disease
 - Humans rarely do this, however: the populations that need the most support typically have the least access

- Heterogeneity increases R₀
 - When disease is rare, it is concentrated in the most vulnerable populations
 - Cases per case is high
 - ► Elimination is harder
- Marginal populations
 - Heterogeneity could make it easier to concentrate on the most vulnerable populations and eliminate disease
 - Humans rarely do this, however: the populations that need the most support typically have the least access