

## FORUM REVIEW ARTICLE

**RNA-mediated control of protein translation under stress**Matthew J. Payea\*, Showkat A. Dar\*, Sulochan Malla, Manolis Maragkakis<sup>§</sup>

Laboratory of Genetics and Genomics, National Institute on Aging, Intramural Research  
Program, National Institutes of Health, Baltimore, MD 21224, USA

\* Equal contribution

<sup>§</sup> To whom correspondence should be addressed.**Contact:** maragkakis@nih.gov

## Abstract

The need of cells to constantly respond to endogenous and exogenous stress has necessitated the evolution of pathways to counter the deleterious effects of stress and to restore cellular homeostasis. The inability to activate a timely and adequate response can lead to disease and is a hallmark of aging. Besides protein-coding genes, cells contain a plethora of non-coding regulatory elements that allow cells to respond rapidly and efficiently to external stimuli by activating highly specific and tightly controlled mechanisms. Many of these programs converge on the regulation of translation, one of the most energy-consuming processes in cells. The non-coding dimension of translational regulation includes short and long non-coding RNAs, as well as mRNA features, such as the sequence and modification status of the 5' and 3' untranslated regions (UTRs), that do not change the amino acid sequence of the produced protein. In this review, we discuss the regulatory role of the non-protein-coding components of translation under stress, particularly oxidative stress. We conclude that the regulation of translation through non-coding RNAs, UTRs and nucleotide modifications are emerging as critical components of the stress response and suggest further areas of study that will improve understanding of this growing field.

## Introduction

During oxidative stress, cells accumulate high levels of reactive oxygen species (ROS) that can originate from endogenous or external sources such as the mitochondria (Larosa and Remacle, 2018), cadmium, arsenite and pollutants (Archer, 2011, Nemmiche, 2017). While ROS have several established roles in signaling (Archer, 2011, Nemmiche, 2017), their highly reactive nature and unregulated increase can cause oxidation of proteins (Holmstrom and Finkel, 2014, Reczek and Chandel, 2015, Schieber and Chandel, 2014), DNA (Dizdaroglu and Jaruga, 2012), RNA (Li et al., 2014, Wilkinson et al., 2021), and lipids (Angelova et al., 2021). Oxidation of these fundamental molecules can lead to permanent damage and cellular dysfunction and, if unchecked, cell death. Cells have evolved with mechanisms to defend against oxidative stress by producing detoxifying enzymes, amino acids, and other molecules (Gorrini et al., 2013), as well as by limiting the amounts of endogenously produced ROS. As the most energy-intensive process in the cell, translation requires high levels of mitochondrial respiration that generates a substantial amount of ROS (Leibovitch and Topisirovic, 2018), and is thus tightly regulated as part of an oxidative stress response.

While the factors that regulate translation are most readily associated with the coding region of mRNA, as well as the protein components of the translation machinery, non-coding RNA (ncRNA) and the untranslated regions (UTRs) of mRNA are equally essential regulatory elements. ncRNAs are a diverse class of structural and functional RNAs with roles across all areas of biology (Goodall and Wickramasinghe, 2021). In translation, ncRNAs such as rRNAs are responsible for the function and structure of the ribosome, the delivery of amino acids to the ribosome by tRNA, and the modulation of translation efficiency through long ncRNA (lncRNA) sponging of miRNAs or direct binding to mRNA. In contrast, UTRs are present on the 5' and 3' ends of coding mRNAs, and while they are not actively translated themselves, they are targets of regulatory pathways that govern translation. The UTR regions are also frequent substrates for nucleotide modifications that can alter their function in translation. In this review, we will discuss the mechanisms by which oxidative stress alters translation from the perspective of non-coding elements such as ncRNAs, UTRs, and RNA modifications. We will describe how these non-coding elements

participate in translation and serve as both critical modulators of translation efficiency and regions of vulnerability to oxidative stress. Finally, we will discuss future outlooks for the study of non-coding elements in translation and advancement of new technologies applicable to the field.

### Non-coding RNA elements in translation

Protein translation, the synthesis of protein from the mRNA template, can be divided into three phases of ribosomal action: initiation, during which ribosomes are loaded onto the mRNA; elongation, during which the polypeptide chain is synthesized according to the mRNA coding region; and termination, during which the nascent peptide chain is released and ribosomal subunits are recycled. While the subject of translation is the open reading frame (ORF) corresponding to the mRNA region coding for the protein, the act of translation requires numerous non-coding RNA elements, including some within the ribosome itself (**Fig. 1**).

Translation begins with a mature mRNA that has completed splicing and that contains a modified guanine nucleotide N7-methylguanosine ( $m^7G$ ) cap at the 5' end. Capped mRNAs are recognized in the cytoplasm by the eukaryotic initiation factor 4F (eIF4F) complex (eIF4E, eIF4G, and eIF4A), which serves both scaffolding and helicase functions, and eIF4B, which enhances the helicase function of eIF4F (Jackson et al., 2010). Part of the function of eIF4F is to facilitate ribosome loading by unwinding the secondary structure in the mRNA 5' UTR (Pestova and Kolupaeva, 2002). After the mRNA is loaded with the eIF4F complex and the 5' UTR is unwound, the 43S pre-initiation complex, composed of the 40S small ribosomal subunit, the ternary complex (eIF2 - GTP -  $tRNA^{iMet}$ ), and other initiation factors (Jackson et al., 2010) is recruited. This process is commonly referred to as cap-dependent or 5'-dependent translation and represents most translation events in cells. A separate cap-independent form of translation also exists and uses Internal Ribosome Entry Sites (IRESs) to recruit the 43S complex, rather than an eIF4F-cap structure (Yang and Wang, 2019); this mechanism of translation is discussed later in this review. Even though mRNAs largely begin translation using a 5' dependent mechanism, the efficiency by which ribosomes engage with mRNAs is not equal among all transcripts.

Initiation efficiency is determined by sequence elements and structures in the 5'UTR (Leppek et al., 2018), which can also act as a mechanism for selective translation during stress, as is the case for the mammalian target of rapamycin complex 1 (mTORC1) pathway discussed further below.

Once loaded, the 43S pre-initiation complex begins “scanning” through the 5' UTR in search of an AUG start site that the tRNA<sup>iMet</sup> of the loaded ternary complex can pair with. To ensure fidelity of start site selection, the intended start site that defines the expected ORF is often in an optimal context defined by the “Kozak Sequence” (GCCRCCAUGG in vertebrates), wherein the most important nucleotides are a purine at -3 and a G at +4, relative to the A in the AUG codon (Hernandez et al., 2019). However, start sites are not exclusive to the expected ORF, and can appear elsewhere in the 5' UTR where they can also act as alternative translation initiation sites and define upstream ORFs (uORFs). A growing number of mRNAs have been shown to contain uORFs (McGillivray et al., 2018), which can regulate translation through multiple mechanisms (Young and Wek, 2016). Regardless, after start site selection, the GTP of the ternary complex is consumed to GDP and a free 60S ribosomal subunit binds to the committed 40S subunit, displacing the remaining initiation factors (Jackson et al., 2010) and forming an 80S monosome.

Elongation begins with a formed 80S monosome loaded with a tRNA<sup>iMet</sup> in the P-site that is paired to the AUG start site in a codon-anticodon interaction. tRNAs are highly structured ncRNAs that are essential for elongation, as they are the source of amino acids during protein synthesis. Amino acids are covalently attached to tRNA by aminoacyl-tRNA synthetases and recognized by the elongation factor eEF1A, that then brings the tRNA to the A-site of the ribosome (Dever and Green, 2012). In the A-site, the anticodon loop of the tRNA is paired to the codon of the mRNA, assisted by rRNA bases in the small subunit. tRNA pairing in the A-site is a rate-limiting step during elongation and is dependent on factors such as tRNA abundance, the strength of pairing between the codon and anticodon, and even neighboring codons (Brule and Grayhack, 2017, Hanson and Coller, 2018). tRNAs are also regulated extensively during stress, and can be altered in abundance depending on cellular conditions (Gingold et al., 2014), modification status (Chan et al., 2010, Endres et al., 2015), and other types of regulation discussed below. After the tRNA is

accommodated in the A-site and eEF1A is released, peptide bond formation occurs between the amino acid of the A-site tRNA and the P-site peptidyl chain in the peptidyl transfer center of the ribosome; the ribosome then translocates one codon, assisted by the elongation factor eEF2, and new tRNAs are accommodated into the A-site (Dever and Green, 2012).

The process of elongation continues until a stop codon (UAG, UGA, UAA) occupies the A-site and is recognized by the release factors eRF1 and eRF3 (Gerovac and Tampe, 2019). Recognition of the stop codon by eRF1 triggers the release of the nascent peptide and the splitting of the ribosomal subunits by ABCE1 into a 60S subunit and an mRNA-tRNA-bound 40S subunit that is recycled before translation initiation can begin again (Pisarev et al., 2010). As with start codons, the context of stop codons determines the efficiency of the stop, which is important during pre-mature stop codon insertions and in ribosome readthrough (Lombardi et al., 2022).

### **rRNA susceptibility to oxidative stress**

The ribosome is a ribonucleoprotein complex with a ncRNA core composed of rRNA for both the 40S subunit (18S rRNA) and the 60S subunit (28S, 5.8S, 5S rRNAs), around which a multitude of ribosomal proteins (33 for the 40S and 47 proteins for the 60S) are arranged (Henras et al., 2015). The maturation of rRNAs is a highly complex process that varies even among eukaryotes. Briefly, a single precursor rRNA is transcribed by Pol I and extensively processed and modified to yield the 18S, 28S, and 5.8S segments, while the 5S rRNA is transcribed separately by Pol III (Henras et al., 2015); the resulting mature rRNAs, along with the ribosomal proteins, are assembled into the pre-40S and pre-60S subunits (Panse and Johnson, 2010).

rRNA is the most abundant RNA species in the cell, accounting for ~80% of cellular RNA content (Milo et al., 2010), while ribosomes themselves exist at an order of magnitude of  $10^7$  copies per cell (Wisniewski et al., 2014). The high abundance of these components make ribosome biogenesis a highly energy-consuming process (Warner, 1999), and thus a major contributor to the generation of endogenous ROS. Much of ribosome biogenesis, including rDNA transcription, rRNA processing and maturation, and

subunit assembly, occurs within the nucleolus, a membraneless region in the nucleus (Boisvert et al., 2007) (**Fig. 2**). Since the nucleolus functions as the factory for ribosomes, it is also a sensor for stress (Boulon et al., 2010). During oxidative stress, ROS can induce nucleolar stress and negatively affect rRNA processing, thus decreasing the number of ribosomes produced (Chou and Lo, 2019). It has recently been shown that the first stages of rRNA processing are inhibited during oxidative stress (Szaflarski et al., 2022) and induce accumulation of precursor rRNAs (Sapio et al., 2021).

Besides inhibition of rRNA biogenesis, ROS can also directly damage rRNA and thus inhibit translation (Shcherbik and Pestov, 2019). rRNA has extensive secondary structure that is essential to the formation of the ribosome. rRNA bases form critical components of the decoding and peptidyl transferase center of the ribosome, where codon-anticodon pairing and amino acids transfer to the nascent peptide chain occur, respectively (Khatter et al., 2015). ROS-mediated damage to rRNA is unevenly distributed across the rRNA. In bacterial ribosomes, oxidative stress is more pervasive in damaging the catalytic center of the large subunit than the decoding center (Willi et al., 2018). In yeast, endonucleolytic cleavage of a specific site in rRNA of the 60S subunit has been proposed to be a marker of oxidative stress (Shedlovskiy et al., 2017). Furthermore, as with tRNA discussed later, oxidative stress can also induce rRNA cleavage (Thompson et al., 2008). Direct rRNA oxidation is associated with several neurodegenerative diseases, likely because of significant oxidative stress produced in the brain (Cobley et al., 2018). Increased rRNA oxidation is associated with early stages of Alzheimer's disease, and the resulting decrease in protein synthesis may be a critical stage in the development of the disease (Ding et al., 2006, Ding et al., 2012). Similarly, oxidative stress-induced dysfunctional ribosome biogenesis is also associated with Parkinson's disease (Parlato and Liss, 2014).

### **tRNA are regulatory targets of oxidative stress**

tRNAs are short, stable, structured, and highly modified ncRNAs that are essential for translation (Phizicky and Hopper, 2010). Amino acids are covalently attached to the 3' end of tRNAs, while the distal anticodon loop is used to decode mRNA codons during translation. While there are 20 canonical amino acids and 21 corresponding aminoacyl-

tRNA synthetases (Rubio Gomez and Ibba, 2020), there are many more tRNA species owing to the degeneracy of the genetic code (61 codons for 20 amino acids), and an even greater number of tRNA genes, since a given tRNA species often exists with multiple DNA copies. In fact, 415 individual tRNA genes have been predicted in human cells (Chan and Lowe, 2016). This diversity of tRNA genes arises from the presence of tRNA “isoacceptors” and “isodecoders” (Schimmel, 2018). tRNA species are considered isodecoders when the anticodon is identical but the tRNA body contains differences; for example, tRNA<sup>Ala(AGC)</sup> has approximately 15 variants (Chan and Lowe, 2016), though all of them decode the same alanine codon. The role of isodecoders in human biology and their effects on translation are poorly understood, although isodecoders have tissue specificity (Ishimura et al., 2014), indicating that this diversity may be important for cellular function.

In contrast, isoacceptors accept the same amino acid but recognize different mRNA codons owing to different anticodons; for example, the tRNA<sup>Ser</sup> isoacceptor family has four anticodons (AGA, CGA, UGA, GCU) that decode 6 serine codons in the mRNA (UCU, UCC, UCA, UCG, AGU, AGC). An important feature of isoacceptors is that codons and anticodons are not always exact matches, and both RNA modifications and non-Watson-Crick base pairing (termed wobble pairing) regulate the anticodon-codon pairing (Kubyshkin et al., 2018). As accommodation into the A-site is the rate-limiting step of translation elongation, this feature creates a substantial element of regulation referred to as codon optimality (Brule and Grayhack, 2017).

Aside from serving as critical components of translation, tRNA have also been shown to perform a regulatory role during oxidative stress in the form of tRNA-derived stress-induced RNAs (tiRNAs) (**Fig. 3**). tiRNA species are produced from endonucleolytic cleavage of tRNA in the anticodon loop by angiogenin (Thompson et al., 2008, Yamasaki et al., 2009), which is typically a pro-growth protein involved in rRNA transcription when localized to the nucleus (Li and Hu, 2010, Tsuji et al., 2005). While the cleavage of tRNAs to produce tiRNAs might seem to regulate translation through tRNA reduction, this appears unlikely given that the amount of tRNA converted to tiRNAs is miniscule compared to the pool available for protein synthesis (Thompson et al., 2008). Instead, tiRNAs have been



shown to have numerous other effects on biology including gene expression, mRNA stability, and translation (George et al., 2022, Wang et al., 2022b).

Binding of tiRNAs to the ORF and 3' UTR of mRNA encoding ribosomal proteins can enhance their translation, which has further downstream effects on ribosome biogenesis (Kim et al., 2017, Kim et al., 2019). tiRNAs can also act to decrease translation by displacing the eIF4G component of eIF4F from the 5' UTR of mRNA, thus inhibiting translation initiation (Ivanov et al., 2011). Recently, it was shown that the potency of tiRNAs to displace eIF4G can be predicted by stretches of guanines in the tiRNA since eIF4G has a propensity to bind G-quadruplexes (Lyons et al., 2020). The function of tiRNAs themselves as biologically active regulators of translation is supported by experiments showing that transfection of tiRNAs into unstressed cells is sufficient to induce the ISR (Yamasaki et al., 2009).

### **circRNAs are substrates for translational regulation under stress**

Circular RNAs (circRNAs) are a large class of RNAs produced by the back-splicing of exons of precursor RNAs, typically ncRNA, (Kristensen et al., 2019) in a wide range of cell types (Hansen et al., 2013, Hsu and Coca-Prados, 1979, Wu et al., 2012). circRNAs are generally less abundant than linear RNAs, but the absence of a 5' and 3' end makes them resistant to decay by exonucleases. Although many of the functions of circRNAs remain unknown, certain abundant circRNAs can sequester miRNAs and RNA-binding proteins, in a mechanism known as "sponging", thus modulating their biological function (Zheng et al., 2016). Disruption of circRNA expression programs has been found to be associated with aging and many human diseases, such as Alzheimer's disease, diabetes, and cancer (Cortés-López et al., 2018, Gruner et al., 2016, Long et al., 2021, Verduci et al., 2021, Vo et al., 2019). Emerging evidence suggests that circRNAs play a role in the regulation of translation of linear RNAs, although direct interaction between circRNAs and the translation machinery has not been discovered thus far. It is therefore probable that their influence on translation is indirectly mediated through miRNA "sponging" (Chekulaeva and Rajewsky, 2019, Lin et al., 2019, Prats et al., 2020). Typically, the RNA-induced silencing complex (RISC) is guided by miRNAs to trigger deadenylation followed by mRNA

degradation or translational repression of target mRNAs (Bartel, 2018). Disruption of the miRNA-mRNA interaction through miRNA sponging by an abundant circRNA reduces miRNA availability and limits its ability to regulate translation (Hansen et al., 2013, Yang et al., 2018, Zhang et al., 2019, Zheng et al., 2016). Similarly, abundant circRNA can act as sponge for translation-associated RNA binding proteins (**Fig. 4**). For example, circPABPN1 interacts with HuR, to inhibit its binding and function as a translational activator of the poly(A)-binding protein nuclear 1 (PABPN1) mRNA (Abdelmohsen et al., 2017, Li et al., 2020).

Recent studies have revealed that circRNAs can also be direct substrates for translation themselves, mediated via Internal Ribosome Entry Sites (IRES). Following ISR activation, cap-dependent translation is reduced under stress (Pakos-Zebrucka et al., 2016). It is therefore reasonable to hypothesize that circRNA translation, mediated via IRES, is likely favored in such conditions. In fact, Chen et al. showed that a circRNA in the fibroblast growth factor receptor 1 (FGFR1) gene produces a protein that partially overlaps the protein product of the linear counterpart (Chen et al., 2021). Interestingly, under stress conditions, the ratio of circular over linear protein production increases, thus enabling the protein product of circFGFR1 to act as a dominant-negative regulator of FGFR1 and induce cell proliferation (Chen et al., 2021). Combined with other recent evidence, these findings highlight a tight connection between the stress response and translation of circRNAs (Chen et al., 2019, Cheng et al., 2019, Feng et al., 2020, Hanan et al., 2020, Li et al., 2021, Zhang and Sui, 2020).

The increase in usage of IRES translation under stress implies a distinct regulatory mechanism for IRES-dependent translation of linear versus circular RNAs. Past studies have identified that the m6A reader YTHDF3 can recognize N6-methyladenosine in circRNA to directly recruit translation initiation factors (Yang et al., 2017). A recent study used a library of random 10-nt sequences inserted before the start codon of GFP coded by a circRNA to identify sequences that can drive circRNA translation (Fan et al., 2022). In total, 97 6-nt long sequences that drive circRNA translation were identified. Most of these sequences were found to be A-U rich and preferentially enriched in endogenous circular rather than linear RNAs. Similarly, Chen et al. used a high-throughput reporter assay to

systematically screen and quantify IRES elements that can facilitate circRNA translation (Chen et al., 2021). Interestingly complementarity between the 18S rRNA and short stem-loops on the IRES was found to facilitate cap-independent translation, specifically for circRNAs. In contrast, the number of AUG codons, the Kozak sequence, and DRACH motif did not have a similar effect (Chen et al., 2021). These results provide evidence of a bipartite mechanism that regulates circular and linear RNA translation via IRES but the exact details of the mechanism remain unknown. An intriguing hypothesis arises regarding the utilization of various components of the translation machinery by cells in response to stress. Typically, cap-dependent translation surpasses IRES-dependent translation in terms of protein output under normal conditions. However, during times of stress, cells have the ability to modify their proteome by downregulating cap-dependent translation and, simultaneously, sustaining or even enhancing IRES-dependent translation specifically for circRNAs (**Fig. 4**).

### **The 5' UTR: translation selectivity during stress**

The critical role of the 5' UTR in ribosome recruitment and start site selection allows it to introduce selectivity to translation, adding an extra layer of regulatory control beyond mRNA abundance. Mechanisms of selective translation are most often activated in response to acute stressors and are notably involved in the two major pathways of translation regulation: the mTORC1 and the Integrated Stress Response (ISR) pathway.

The mTORC1 pathway is a sensor for numerous growth signals, such as the presence of amino acids, energy availability, and growth factors, that regulates translation using the mTORC1 serine/threonine protein kinase (Sengupta et al., 2010). During optimal growth conditions, the mTORC1 pathway acts to increase protein synthesis through phosphorylation of two proteins with divergent regulatory roles: the inhibitory protein eIF4EBP1 and the protein kinase S6K. eIF4EBP1 negatively regulates translation by binding and sequestering eIF4E from the eIF4F cap-binding complex, thereby reducing translation initiation; in contrast, S6K is a pro-translational kinase that phosphorylates components of the translation machinery to increase protein synthesis (Saxton and Sabatini, 2017). mTORC1 phosphorylation of eIF4EBP1 suppresses its binding to eIF4E (Gingras et al.,

2001), while phosphorylation of S6K enhances its function as a kinase (Holz et al., 2005), thus both roles coordinate to increase protein synthesis (**Fig. 5A**).

Although the mTORC1 pathway affects translation globally by activating S6K and suppressing eIF4EBP1, certain mRNAs are affected more than others. Experiments probing selective translation after mTORC1 inhibition identified a common 5' terminal oligopyrimidine (5' TOP) motif among the most sensitive transcripts; this selectivity is largely due to the inhibitory action of eIF4EBP1, since 5' TOP motif RNAs are more sensitive to disruptions of the eIF4E-eIF4G1 components of the eIF4F cap-binding complex (Thoreen et al., 2012). Interestingly, 5' TOP motifs are overwhelmingly found on mRNAs encoding proteins that function in translation, strongly suggesting they are part of a coordinated mechanism to reduce protein synthesis during non-optimal growth conditions that suppress mTORC1 activity (Hsieh et al., 2012). This mechanism is utilized as a survival mechanism in cancer, wherein increased oxidative stress broadly reduces the translation of ribosomal proteins, since their transcripts are sensitive to mTORC1 inhibition (Tang et al., 2016).

In contrast to the mTOR pathway, the ISR pathway primarily functions to decrease translation in response to a diverse array of stressors including viral invasion, endoplasmic reticulum (ER) stress, amino acid starvation, oxidative stress, and others (Pakos-Zebrucka et al., 2016). The ISR pathway regulates translation initiation at the formation of the ternary complex (eIF2 - GTP - tRNA<sup>Met</sup>) necessary for start site selection by phosphorylating the eIF2 $\alpha$  subunit of the eIF2 initiation factor. eIF2 $\alpha$  phosphorylation is mediated through the activity of one of four kinases (eIF2AK1-4), commonly referred to as HRI, PKR, PERK, and GCN2 (Donnelly et al., 2013) (**Fig. 5B**). Due to their highly reactive nature, ROS can induce several different types of damage that will be recognized by the ISR. In fact, HRI (Ill-Raga et al., 2015, Koncha et al., 2021, Suragani et al., 2012, Szwed et al., 2019), PERK (Farrukh et al., 2014, Harding et al., 2003, Verfaillie et al., 2013), GCN2 (Baker et al., 2012, Zhu et al., 2021), and PKR (Li et al., 2010, Zeng et al., 2022) have all been associated with activation of the ISR in response to oxidative stress. Phosphorylation of eIF2 $\alpha$  globally reduces translation initiation, thus limiting protein expression. This reduction in protein expression also produces a selective increase in translation for certain

transcripts with uORFs in their 5' UTR, the most important of which is the multi-purpose transcription factor ATF4. Mechanistically, the decrease in ternary complex due to eIF2 $\alpha$  phosphorylation increases the scanning time of ribosomes, allowing them to bypass an inhibitory uORF in the 5' UTR of ATF4 (Young and Wek, 2016). The use of translation selectivity by 5' uORFs during oxidative stress extends beyond ATF4 as well (Andreev et al., 2015). Activation of the ISR during oxidative stress is also very important, as experiments show that disruption of the ISR increases sensitivity to ROS (Krishnamoorthy et al., 2018, Malin et al., 2021, McEwen et al., 2005, Rajesh et al., 2015).

### **The 3' UTR: Selenocysteine Insertion Under Stress**

Selenoproteins serve antioxidant roles and require non-canonical insertion of a selenocysteine residue at UGA stop codons. This event is mediated by a structured RNA element in the 3' UTR region called the selenocysteine insertion sequence (SECIS) that is expected to scaffold recoding factors, which ultimately enable insertion of selenocysteine at UGA stop codons (Vindry et al., 2018). The SECIS site is necessary for UGA recoding of selenoproteins, as well as sufficient to induce recoding when inserted into non-selenocysteine mRNAs (Shen et al., 1993). During oxidative stress, the recoding factors SBP2, L30, and EFsec are imported into the nucleus where they are recruited to the SECIS sequence in selenoprotein mRNAs before their export into the cytoplasm for translation (Papp et al., 2006, Zahia et al., 2014). Thus, despite its non-coding nature, the 3' UTR can function to dramatically alter translation during oxidative stress.

### **Epitranscriptomic control of translation under stress**

Chemical modifications of RNA nucleotides were first discovered in 1957 (Davis and Allen, 1957) and currently, more than 330 modifications have been described (Boccaletto et al., 2022). Most of them localize on ncRNAs such as tRNAs, rRNAs, snRNAs, snoRNAs and to a lesser extent on mRNAs (Boccaletto et al., 2022). Addition, removal, and sensing of chemical-moieties on RNA bases is mediated by proteins known as writers, erasers, and readers, respectively (Esteve-Puig et al., 2020, Kumar and Mohapatra, 2021, Shi et al., 2019, Xuan et al., 2018). RNA modifications can be reversible (e.g., m<sup>6</sup>A, 5-methylcytosine) or irreversible (e.g., A to I or C to U), add functional complexity in cellular regulation

(Destefanis et al., 2021, Mathlin et al., 2020) and affect a range of cellular functions such as the immune response, cell death, DNA damage repair, stress response and protein translation (Frye et al., 2018, Moradian et al., 2022, Wilkinson et al., 2021, Wilkinson et al., 2022).

N6-methyladenosine ( $m^6A$ ) is among the most common reversible modification in mRNAs and has been involved in cellular response to oxidative stress and pathologies such as cancer, ischemic stroke, neurodegeneration, and aging (Chang et al., 2022, Condic et al., 2022, Desrosiers et al., 1974, Nunomura et al., 2012, Sun et al., 2022, Wu et al., 2022, Yang and Chen, 2021, Zhang et al., 2022a, Zhao et al., 2019). Recent reports have shown that  $m^6A$  deposition on 5' UTRs in response to oxidative stress provides a selective mechanism for sequestering mRNAs to stress granules, membraneless organelles formed upon stress, mediated by YTHDF3 (Anders et al., 2018). In addition, under hypoxic conditions  $m^6A$  content increases for a subset of mRNAs and results in increased mRNA stability (Fry et al., 2017).

N4-acetylcytidine ( $ac^4C$ ), a reversible post-transcriptional RNA modification, is catalyzed by the N-acetyltransferase (NAT10) and has recently been shown to be involved in position-dependent regulation of mRNA translation. While  $ac^4C$  is primarily found in the 5'UTR, in close proximity to the start codon,  $ac^4C$  deposited within the coding region can promote translational efficiency by facilitating decoding of wobble sites and thus preventing mRNA decay (Arango et al., 2018). In contrast,  $ac^4C$  deposited in the 5' UTR impacts translation by promoting initiation at upstream sequences, thus competitively reducing initiation at canonical ORFs. Furthermore,  $ac^4C$  within the Kozak sequence and immediately upstream of optimal start codons can disrupt an interaction between the modified C and the initiator tRNA, thus further inhibiting initiation (Arango et al., 2022). These studies highlight the role of  $ac^4C$  in modulating translation and RNA stability and raise the possibility that these processes are closely regulated in cells and can be hijacked by pathogens or be deregulated in disease. In fact, it was recently shown that HIV-1 RNA is modified with  $ac^4C$  at multiple discrete sites that result in increased stability and thus viral gene expression (Tsai et al., 2020). Similarly,  $ac^4C$  deposited within the 5' UTR of enterovirus 71 (EV71) enhances viral RNA translation via selective recruitment of PCBP2 to

the viral IRES and boosts its genome stability. ac<sup>4</sup>C has also been shown to stabilize and increase translation of oncogenes, thus linking the modification to cancer (Feng et al., 2022, Wang et al., 2022a, Zhang et al., 2021).

Pseudouridine ( $\Psi$ ) is typically deposited in rRNAs and tRNAs and to a lesser extent mRNAs. (Cui et al., 2021, Guzzi et al., 2018, Jack et al., 2011, McMahon et al., 2019) Nevertheless, recent evidence show that deposition of  $\Psi$  on mRNAs is tightly regulated during stress and is implicated with human disease (Barbieri and Kouzarides, 2020, Cerneckis et al., 2022). Interestingly, while oxidative and heat shock stresses increase pseudouridylation, starvation modestly decreases it (Begik et al., 2021, Li et al., 2015). The exact regulatory role of  $\Psi$  under stress remain unknown, but deletion of PUS7, a pseudouridine synthase, in yeast resulted in reduced mRNA stability under stress suggesting a possible role of  $\Psi$  in RNA protection (Schwartz et al., 2014). Bisulfite-induced deletion sequencing (BID-seq) in human cells confirmed that TRUB1-deposited  $\Psi$  sites led to transcript stabilization and ribosome readthrough at  $\Psi$  sites within stop codons (Dai et al., 2022, Fernandez et al., 2013).

N1-methyladenosine ( $m^1A$ ), besides its role in tRNAs, is also found in the 5'UTR of mRNAs close to the transcription start site. It is dynamically regulated under different stress conditions and has been associated with translation efficiency and initiation (Li et al., 2017, Li et al., 2016, Safra et al., 2017). Despite recent progress, more research is required to discover molecular mechanisms whereby mRNA modifications regulate cell processes under stress (**Fig. 6**).

Besides mRNA, RNA modifications are found in ncRNAs such as rRNA and tRNA (Roundtree et al., 2017). The rRNA epitranscriptome is reprogrammed in response to the cellular environment as well as in development and disease (Sloan et al., 2017, Xue and Barna, 2012). 2'-O-methylation of rRNA defines the ribosome conformational status and is essential for its biogenesis and function (Khoshnevis et al., 2022, Natchiar et al., 2018). Also, several rRNA modifications are clustered in the peptidyl transferase center and other functional sites of the ribosome to promote translational fidelity (Sloan et al., 2017). In fact, single-molecule RNA modification profiling revealed that the eukaryotic rRNA is

modified at more than 100 sites, particularly at highly conserved and functionally relevant nucleotides (Bailey et al., 2022). Since cap-independent translation becomes more prominent under stress (Spriggs et al., 2008), the efficiency of these sites for ribosome recruitment and initiation is substantially influenced by modifications both on rRNA and on translation substrates such as mRNAs, circRNAs or lincRNAs (Coots et al., 2017, Meyer et al., 2015, Qin et al., 2022, Yoon et al., 2006). Oxidative stress in particular, results in the creation of abasic sites, strand breaks, and 8-oxo-7,8-dihydroguanine (8-oxoG) modifications on rRNAs and mRNAs that substantially alter their function (Shcherbik and Pestov, 2019).

Modifications on tRNA are also dynamically regulated under stress. Recent evidence shows that wobble modifications in tRNAs lead to selective translation of stress response proteins from codon-biased genes (Endres et al., 2015, Huber et al., 2022). Reprogramming of the epitranscriptome is reflected in changes in translation and its deregulation is involved in disease (Bednarova et al., 2017, Chujo and Tomizawa, 2021, Suzuki, 2021, Torres et al., 2014).

The role of the epitranscriptome in translation has not been fully appreciated until recently, partly due to the lack of scalable methods to probe modified nucleotides. Established technologies for the detection of modifications can generally be divided into quantification methods, locus-specific detection methods and next-generation sequencing-based detection methods (reviewed in (Zhang et al., 2022b)). While the latter constitute a major improvement, they are limited by read length and the inability to probe multiple modifications simultaneously at single RNA molecules. Nanopore direct RNA sequencing is emerging as a technology that can bridge that gap and provide valuable modification information for distinct RNA classes including mRNAs, tRNAs and rRNAs (Gu et al., 2012, Wang et al., 2021, Zhang et al., 2020). However, while the information for modified nucleotides is likely contained in the raw sequencer output, the software to extract the information is currently under development (**Table 1**).



## Conclusion and Perspectives

In this review, we have discussed the role of non-coding RNA elements in translation during cellular stress. While translation has been studied extensively, much remains to be deciphered, particularly regarding the role of ncRNAs. Ribosome biogenesis and the maturation of rRNA continues to be a highly complex area of study with numerous idiosyncrasies between organisms and has now been further shown to change in a directed way during stress. Recent research has shown extensive contributions of the 5' UTR to translation efficiency, in particular the realization that uORFs are far more abundant than previously thought and may mediate selective translational mechanisms not previously known. Similarly, tiRNAs generated during stress from tRNAs have gained increased recognition as important signaling molecules during stress, and the diversity of these molecules is further enhanced by the enormous diversity within tRNA sequences. The identification of circRNAs has also opened new possibilities of decay-resistant RNAs that can be induced by stress to either tune the translation machinery by binding specific RNAs and proteins or be translated themselves through IRES sequences.

The epitranscriptome offers an added level of complexity to translation regulation with RNA modifications having increased utility on mRNAs, rRNAs, and tRNAs. In particular, ac<sup>4</sup>C and m<sup>6</sup>A have recently been found to play roles in the regulation of translation for both mRNAs and circRNAs. The study of RNA modifications at scale has only recently become feasible with the development of new high-throughput, small-read sequencing approaches that reveal genome- and transcriptome-wide maps of modified nucleotides. Nevertheless, while these approaches provide an unprecedented view into the world of RNA modifications, they cannot capture long-range interactions nor the combinatorial effect of different modifications on the same RNA molecule. Intriguingly, new technologies using nanopore sequencing are emerging as potentially revolutionary approaches to sequence RNA directly and deconvolve modified nucleotides and other RNA features at the single-molecule level. Future studies will need to address current technology and software shortcomings to profile the complete RNome and expand upon the currently limited subset of identifiable modifications. This information will allow a transcriptome-

wide exploration of the combinatorial role of RNA features, including modifications and 5'UTR regulatory elements, on translation under stress.

### **Acknowledgements**

We thank Dr. Myriam Gorospe for editing our manuscript and providing feedback. We would also like to thank Yolanda L. Jones, NIH Library, for manuscript editing assistance. Figures were created with BioRender.com.

### **Funding**

This work was supported by the Intramural Research Program of the National Institute on Aging, National Institutes of Health (ZIAAG000696).

### **Conflict of Interest**

None

## Abbreviations Used

8-oG: 8-Oxoguanine

ABCE1: ATP Binding Cassette Subfamily E Member 1

ac<sup>4</sup>C: N4-Acetylcytidine

ATF4: Activating Transcription Factor 4

BID-seq: Bisulfite-Induced Deletion sequencing

circRNA: circular ribonucleic acid

DNA: Deoxyribonucleic Acid

eEF1A: eukaryotic translation Elongation Factor 1A

eEF2: eukaryotic translation Elongation Factor 2

eIF2: eukaryotic translation Initiation Factor 2

eIF2AK1: eukaryotic translation Initiation Factor 2 $\alpha$  Kinase 1

eIF2AK2: eukaryotic translation Initiation Factor 2 $\alpha$  Kinase 2

eIF2AK3: eukaryotic translation Initiation Factor 2 $\alpha$  Kinase 3

eIF2AK4: eukaryotic translation Initiation Factor 2 $\alpha$  Kinase 4

eIF2 $\alpha$ : eukaryotic translation Initiation Factor 2 $\alpha$

eIF4A: eukaryotic translation Initiation factor 4A

eIF4B: eukaryotic translation Initiation factor 4B

eIF4E: eukaryotic translation Initiation factor 4E

eIF4E: eukaryotic translation Initiation factor 4E

eIF4EBP1: eukaryotic translation Initiation Factor 4E Binding Protein 1

eIF4F: eukaryotic translation Initiation Factor 4F

eIF4G: eukaryotic translation Initiation Factor 4G

eRF1: eukaryotic translation Recycling Factor 1

GCN2: General Control Nonderepressible 2

GDP: Guanosine Diphosphate

GFP: Green Fluorescent Protein

GTP: Guanosine Triphosphate

HRI: Heme-Regulated eukaryotic translation Initiation Factor 2 $\alpha$  Kinase

Ikk $\beta$ : I-kappaB kinase beta

IRES: Internal Ribosome Entry Site

ISR: Integrated Stress Response

LncRNA: Long noncoding RNA

m<sup>1</sup>A: N1-methyladenosine

m<sup>5</sup>C: N5-methylcytosine

m<sup>6</sup>A: N6-methyladenosine

m<sup>7</sup>G: N7-methylguanosine

miRNA: micro Ribonucleic Acid

mRNA: messenger Ribonucleic Acid

mTORC1: mammalian Target of Rapamycin Complex 1

ORF: Open Reading Frame

PERK: Protein kinase RNA-like ER Kinase

PKR: Protein Kinase RNA-activated

Pol I: DNA Polymerase I

Pol III: DNA Polymerase III

rDNA: ribosomal Deoxynucleic Acid

RISC: RNA-induced Silencing Complex

RNA: Ribonucleic Acid

ROS: Reactive Oxygen Species

rRNA: ribosomal Ribonucleic Acid

tiRNA: tRNA-derived stress-induced Ribonucleic Acid

TOP motif: 5' Terminal Oligopyrimidine Motif

tRNA: transfer Ribonucleic Acid

tRNA<sup>Ala</sup>(AGC): tRNA Alanine AGC

tRNA<sup>iMet</sup>: tRNA initiator Methionine

tRNA<sup>Ser</sup>: tRNA Serine

TSC: Tuberous Sclerosis Complex

uORF: upstream Open Reading Frame

UTR: Untranslated Region

Ψ: Pseudouridine

## References

- Abdelmohsen K, Panda AC, Munk R, et al. Identification of HuR target circular RNAs uncovers suppression of PABPN1 translation by CircPABPN1. *RNA Biol* 2017;14(3):361-369; doi: 10.1080/15476286.2017.1279788.
- Anders M, Chelysheva I, Goebel I, et al. Dynamic m(6)A methylation facilitates mRNA triaging to stress granules. *Life Sci Alliance* 2018;1(4):e201800113; doi: 10.26508/lsa.201800113.
- Andreev DE, O'Connor PB, Fahey C, et al. Translation of 5' leaders is pervasive in genes resistant to eIF2 repression. *Elife* 2015;4:e03971; doi: 10.7554/eLife.03971.
- Angelova PR, Esteras N, Abramov AY. Mitochondria and lipid peroxidation in the mechanism of neurodegeneration: Finding ways for prevention. *Med Res Rev* 2021;41(2):770-784; doi: 10.1002/med.21712.
- Arango D, Sturgill D, Alhusaini N, et al. Acetylation of Cytidine in mRNA Promotes Translation Efficiency. *Cell* 2018;175(7):1872-1886 e1824; doi: 10.1016/j.cell.2018.10.030.
- Arango D, Sturgill D, Yang R, et al. Direct epitranscriptomic regulation of mammalian translation initiation through N4-acetylcytidine. *Mol Cell* 2022;82(15):2797-2814 e2711; doi: 10.1016/j.molcel.2022.05.016.
- Archer T. Effects of exogenous agents on brain development: stress, abuse and therapeutic compounds. *CNS Neurosci Ther* 2011;17(5):470-489; doi: 10.1111/j.1755-5949.2010.00171.x.
- Bailey AD, Talkish J, Ding H, et al. Concerted modification of nucleotides at functional centers of the ribosome revealed by single-molecule RNA modification profiling. *Elife* 2022;11; doi: 10.7554/eLife.76562.
- Baker BM, Nargund AM, Sun T, et al. Protective coupling of mitochondrial function and protein synthesis via the eIF2alpha kinase GCN-2. *PLoS Genet* 2012;8(6):e1002760; doi: 10.1371/journal.pgen.1002760.

Barbieri I, Kouzarides T. Role of RNA modifications in cancer. *Nat Rev Cancer* 2020;20(6):303-322; doi: 10.1038/s41568-020-0253-2.

Bartel DP. Metazoan MicroRNAs. *Cell* 2018;173(1):20-51; doi: 10.1016/j.cell.2018.03.006.

Bednarova A, Hanna M, Durham I, et al. Lost in Translation: Defects in Transfer RNA Modifications and Neurological Disorders. *Front Mol Neurosci* 2017;10:135; doi: 10.3389/fnmol.2017.00135.

Begik O, Lucas MC, Pryszcz LP, et al. Quantitative profiling of pseudouridylation dynamics in native RNAs with nanopore sequencing. *Nat Biotechnol* 2021;39(10):1278-1291; doi: 10.1038/s41587-021-00915-6.

Boccaletto P, Stefaniak F, Ray A, et al. MODOMICS: a database of RNA modification pathways. 2021 update. *Nucleic Acids Res* 2022;50(D1):D231-D235; doi: 10.1093/nar/gkab1083.

Boisvert FM, van Koningsbruggen S, Navascues J, et al. The multifunctional nucleolus. *Nat Rev Mol Cell Biol* 2007;8(7):574-585; doi: 10.1038/nrm2184.

Boulon S, Westman BJ, Hutten S, et al. The nucleolus under stress. *Mol Cell* 2010;40(2):216-227; doi: 10.1016/j.molcel.2010.09.024.

Brule CE, Grayhack EJ. Synonymous Codons: Choose Wisely for Expression. *Trends Genet* 2017;33(4):283-297; doi: 10.1016/j.tig.2017.02.001.

Cerneckis J, Cui Q, He C, et al. Decoding pseudouridine: an emerging target for therapeutic development. *Trends Pharmacol Sci* 2022;43(6):522-535; doi: 10.1016/j.tips.2022.03.008.

Chan CT, Dyavaiah M, DeMott MS, et al. A quantitative systems approach reveals dynamic control of tRNA modifications during cellular stress. *PLoS Genet* 2010;6(12):e1001247; doi: 10.1371/journal.pgen.1001247.

Chan PP, Lowe TM. GtRNAdb 2.0: an expanded database of transfer RNA genes identified in complete and draft genomes. *Nucleic Acids Res* 2016;44(D1):D184-189; doi: 10.1093/nar/gkv1309.

Chang H, Yang J, Wang Q, et al. Role of N6-methyladenosine modification in pathogenesis of ischemic stroke. *Expert Rev Mol Diagn* 2022;22(3):295-303; doi: 10.1080/14737159.2022.2049246.

Chekulaeva M, Rajewsky N. Roles of long noncoding RNAs and circular RNAs in translation. *Cold Spring Harb Perspect Biol* 2019;11(6):a032680; doi: 10.1101/cshperspect.a032680.

Chen B, Li Y, Liu Y, et al. circLRP6 regulates high glucose-induced proliferation, oxidative stress, ECM accumulation, and inflammation in mesangial cells. *J Cell Physiol* 2019;234(11):21249-21259; doi: 10.1002/jcp.28730.

Chen C-K, Cheng R, Demeter J, et al. Structured elements drive extensive circular RNA translation. *Mol Cell* 2021;81(20):4300-4318.e4313; doi: 10.1016/j.molcel.2021.07.042.

Cheng Q, Cao X, Xue L, et al. CircPRKCI-miR-545/589-E2F7 axis dysregulation mediates hydrogen peroxide-induced neuronal cell injury. *Biochem Biophys Res Commun* 2019;514(2):428-435; doi: 10.1016/j.bbrc.2019.04.131.

Chou YT, Lo KY. Thallium(I) treatment induces nucleolar stress to stop protein synthesis and cell growth. *Sci Rep* 2019;9(1):6905; doi: 10.1038/s41598-019-43413-1.

Chujo T, Tomizawa K. Human transfer RNA modopathies: diseases caused by aberrations in transfer RNA modifications. *FEBS J* 2021;288(24):7096-7122; doi: 10.1111/febs.15736.

Cobley JN, Fiorello ML, Bailey DM. 13 reasons why the brain is susceptible to oxidative stress. *Redox Biol* 2018;15:490-503; doi: 10.1016/j.redox.2018.01.008.

Condic M, Ralser DJ, Klumper N, et al. Comprehensive Analysis of N6-Methyladenosine (m6A) Writers, Erasers, and Readers in Cervical Cancer. *Int J Mol Sci* 2022;23(13); doi: 10.3390/ijms23137165.

Coots RA, Liu XM, Mao Y, et al. m(6)A Facilitates eIF4F-Independent mRNA Translation. *Mol Cell* 2017;68(3):504-514 e507; doi: 10.1016/j.molcel.2017.10.002.



Cortés-López M, Gruner MR, Cooper DA, et al. Global accumulation of circRNAs during aging in *Caenorhabditis elegans*. *BMC Genomics* 2018;19(1):8; doi: 10.1186/s12864-017-4386-y.

Cui Q, Yin K, Zhang X, et al. Targeting PUS7 suppresses tRNA pseudouridylation and glioblastoma tumorigenesis. *Nat Cancer* 2021;2(9):932-949; doi: 10.1038/s43018-021-00238-0.

Dai Q, Zhang LS, Sun HL, et al. Quantitative sequencing using BID-seq uncovers abundant pseudouridines in mammalian mRNA at base resolution. *Nat Biotechnol* 2022; doi: 10.1038/s41587-022-01505-w.

Davis FF, Allen FW. Ribonucleic acids from yeast which contain a fifth nucleotide. *J Biol Chem* 1957;227(2):907-915.

Desrosiers R, Friderici K, Rottman F. Identification of methylated nucleosides in messenger RNA from Novikoff hepatoma cells. *Proc Natl Acad Sci U S A* 1974;71(10):3971-3975; doi: 10.1073/pnas.71.10.3971.

Destefanis E, Avsar G, Groza P, et al. A mark of disease: how mRNA modifications shape genetic and acquired pathologies. *RNA* 2021;27(4):367-389; doi: 10.1261/rna.077271.120.

Dever TE, Green R. The elongation, termination, and recycling phases of translation in eukaryotes. *Cold Spring Harb Perspect Biol* 2012;4(7):a013706; doi: 10.1101/cshperspect.a013706.

Ding Q, Markesbery WR, Cekarini V, et al. Decreased RNA, and increased RNA oxidation, in ribosomes from early Alzheimer's disease. *Neurochem Res* 2006;31(5):705-710; doi: 10.1007/s11064-006-9071-5.

Ding Q, Zhu H, Zhang B, et al. Increased 5S rRNA oxidation in Alzheimer's disease. *J Alzheimers Dis* 2012;29(1):201-209; doi: 10.3233/JAD-2012-111058.

Dizdaroglu M, Jaruga P. Mechanisms of free radical-induced damage to DNA. *Free Radic Res* 2012;46(4):382-419; doi: 10.3109/10715762.2011.653969.

Donnelly N, Gorman AM, Gupta S, et al. The eIF2 $\alpha$  kinases: their structures and functions. *Cell Mol Life Sci* 2013;70(19):3493-3511; doi: 10.1007/s00018-012-1252-6.

Endres L, Dedon PC, Begley TJ. Codon-biased translation can be regulated by wobble-base tRNA modification systems during cellular stress responses. *RNA Biol* 2015;12(6):603-614; doi: 10.1080/15476286.2015.1031947.

Esteve-Puig R, Bueno-Costa A, Esteller M. Writers, readers and erasers of RNA modifications in cancer. *Cancer Lett* 2020;474:127-137; doi: 10.1016/j.canlet.2020.01.021.

Fan X, Yang Y, Chen C, et al. Pervasive translation of circular RNAs driven by short IRES-like elements. *Nat Commun* 2022;13(1):3751; doi: 10.1038/s41467-022-31327-y.

Farrukh MR, Nissar UA, Afnan Q, et al. Oxidative stress mediated Ca<sup>2+</sup> release manifests endoplasmic reticulum stress leading to unfolded protein response in UV-B irradiated human skin cells. *J Dermatol Sci* 2014;75(1):24-35; doi: 10.1016/j.jdermsci.2014.03.005.

Feng D, Wang Z, Zhao Y, et al. circ-PRKCB acts as a ceRNA to regulate p66Shc-mediated oxidative stress in intestinal ischemia/reperfusion. *Theranostics* 2020;10(23):10680-10696; doi: 10.7150/thno.44250.

Feng Z, Li K, Qin K, et al. The LINC00623/NAT10 signaling axis promotes pancreatic cancer progression by remodeling ac4C modification of mRNA. *J Hematol Oncol* 2022;15(1):112; doi: 10.1186/s13045-022-01338-9.

Fernandez IS, Ng CL, Kelley AC, et al. Unusual base pairing during the decoding of a stop codon by the ribosome. *Nature* 2013;500(7460):107-110; doi: 10.1038/nature12302.

Fry NJ, Law BA, Ilkayeva OR, et al. N(6)-methyladenosine is required for the hypoxic stabilization of specific mRNAs. *RNA* 2017;23(9):1444-1455; doi: 10.1261/rna.061044.117.

Frye M, Harada BT, Behm M, et al. RNA modifications modulate gene expression during development. *Science* 2018;361(6409):1346-1349; doi: 10.1126/science.aau1646.

George S, Rafi M, Aldarmaki M, et al. tRNA derived small RNAs-Small players with big roles. *Front Genet* 2022;13:997780; doi: 10.3389/fgene.2022.997780.

Gerovac M, Tampe R. Control of mRNA Translation by Versatile ATP-Driven Machines. *Trends Biochem Sci* 2019;44(2):167-180; doi: 10.1016/j.tibs.2018.11.003.

Gingold H, Tehler D, Christoffersen NR, et al. A dual program for translation regulation in cellular proliferation and differentiation. *Cell* 2014;158(6):1281-1292; doi: 10.1016/j.cell.2014.08.011.

Gingras AC, Raught B, Gygi SP, et al. Hierarchical phosphorylation of the translation inhibitor 4E-BP1. *Genes Dev* 2001;15(21):2852-2864; doi: 10.1101/gad.912401.

Goodall GJ, Wickramasinghe VO. RNA in cancer. *Nat Rev Cancer* 2021;21(1):22-36; doi: 10.1038/s41568-020-00306-0.

Gorrini C, Harris IS, Mak TW. Modulation of oxidative stress as an anticancer strategy. *Nat Rev Drug Discov* 2013;12(12):931-947; doi: 10.1038/nrd4002.

Gruner H, Cortés-López M, Cooper DA, et al. CircRNA accumulation in the aging mouse brain. *Sci Rep* 2016;6(1); doi: 10.1038/srep38907.

Gu LQ, Wanunu M, Wang MX, et al. Detection of miRNAs with a nanopore single-molecule counter. *Expert Rev Mol Diagn* 2012;12(6):573-584; doi: 10.1586/erm.12.58.

Guzzi N, Ciesla M, Ngoc PCT, et al. Pseudouridylation of tRNA-Derived Fragments Steers Translational Control in Stem Cells. *Cell* 2018;173(5):1204-1216 e1226; doi: 10.1016/j.cell.2018.03.008.

Hanan M, Simchovitz A, Yayon N, et al. A Parkinson's disease CircRNAs Resource reveals a link between circSLC8A1 and oxidative stress. *EMBO Mol Med* 2020;12(9):e11942; doi: 10.15252/emmm.201911942.

Hansen TB, Jensen TI, Clausen BH, et al. Natural RNA circles function as efficient microRNA sponges. *Nature* 2013;495(7441):384-388; doi: 10.1038/nature11993.

Hanson G, Collier J. Codon optimality, bias and usage in translation and mRNA decay. *Nat Rev Mol Cell Biol* 2018;19(1):20-30; doi: 10.1038/nrm.2017.91.

Harding HP, Zhang Y, Zeng H, et al. An integrated stress response regulates amino acid metabolism and resistance to oxidative stress. *Mol Cell* 2003;11(3):619-633; doi: 10.1016/s1097-2765(03)00105-9.

Hendra C, Pratanwanich PN, Wan YK, et al. Detection of m6A from direct RNA sequencing using a multiple instance learning framework. *Nat Methods* 2022;19(12):1590-1598; doi: 10.1038/s41592-022-01666-1.

Henras AK, Plisson-Chastang C, O'Donohue MF, et al. An overview of pre-ribosomal RNA processing in eukaryotes. *Wiley Interdiscip Rev RNA* 2015;6(2):225-242; doi: 10.1002/wrna.1269.

Hernandez G, Osnaya VG, Perez-Martinez X. Conservation and Variability of the AUG Initiation Codon Context in Eukaryotes. *Trends Biochem Sci* 2019;44(12):1009-1021; doi: 10.1016/j.tibs.2019.07.001.

Holmstrom KM, Finkel T. Cellular mechanisms and physiological consequences of redox-dependent signalling. *Nat Rev Mol Cell Biol* 2014;15(6):411-421; doi: 10.1038/nrm3801.

Holz MK, Ballif BA, Gygi SP, et al. mTOR and S6K1 mediate assembly of the translation preinitiation complex through dynamic protein interchange and ordered phosphorylation events. *Cell* 2005;123(4):569-580; doi: 10.1016/j.cell.2005.10.024.

Hsieh AC, Liu Y, Edlind MP, et al. The translational landscape of mTOR signalling steers cancer initiation and metastasis. *Nature* 2012;485(7396):55-61; doi: 10.1038/nature10912.

Hsu MT, Coca-Prados M. Electron microscopic evidence for the circular form of RNA in the cytoplasm of eukaryotic cells. *Nature* 1979;280(5720):339-340; doi: 10.1038/280339a0.

Huber SM, Begley U, Sarkar A, et al. Arsenite toxicity is regulated by queuine availability and oxidation-induced reprogramming of the human tRNA epitranscriptome. *Proc Natl Acad Sci U S A* 2022;119(38):e2123529119; doi: 10.1073/pnas.2123529119.

Ill-Raga G, Tajés M, Busquets-García A, et al. Physiological Control of Nitric Oxide in Neuronal BACE1 Translation by Heme-Regulated eIF2 $\alpha$  Kinase HRI Induces Synaptogenesis. *Antioxid Redox Signal* 2015;22(15):1295-1307; doi: 10.1089/ars.2014.6080.

Ishimura R, Nagy G, Dotu I, et al. RNA function. Ribosome stalling induced by mutation of a CNS-specific tRNA causes neurodegeneration. *Science* 2014;345(6195):455-459; doi: 10.1126/science.1249749.

Ivanov P, Emara MM, Villen J, et al. Angiogenin-induced tRNA fragments inhibit translation initiation. *Mol Cell* 2011;43(4):613-623; doi: 10.1016/j.molcel.2011.06.022.

Jack K, Bellodi C, Landry DM, et al. rRNA pseudouridylation defects affect ribosomal ligand binding and translational fidelity from yeast to human cells. *Mol Cell* 2011;44(4):660-666; doi: 10.1016/j.molcel.2011.09.017.

Jackson RJ, Hellen CU, Pestova TV. The mechanism of eukaryotic translation initiation and principles of its regulation. *Nat Rev Mol Cell Biol* 2010;11(2):113-127; doi: 10.1038/nrm2838.

Jenjaroenpun P, Wongsurawat T, Wadley TD, et al. Decoding the epitranscriptional landscape from native RNA sequences. *Nucleic Acids Res* 2021;49(2):e7; doi: 10.1093/nar/gkaa620.

Khatter H, Myasnikov AG, Natchiar SK, et al. Structure of the human 80S ribosome. *Nature* 2015;520(7549):640-645; doi: 10.1038/nature14427.

Khoshnevis S, Dreggors-Walker RE, Marchand V, et al. Ribosomal RNA 2'-O-methylations regulate translation by impacting ribosome dynamics. *Proc Natl Acad Sci U S A* 2022;119(12):e2117334119; doi: 10.1073/pnas.2117334119.

Kim HK, Fuchs G, Wang S, et al. A transfer-RNA-derived small RNA regulates ribosome biogenesis. *Nature* 2017;552(7683):57-62; doi: 10.1038/nature25005.

Kim HK, Xu J, Chu K, et al. A tRNA-Derived Small RNA Regulates Ribosomal Protein S28 Protein Levels after Translation Initiation in Humans and Mice. *Cell Rep* 2019;29(12):3816-3824 e3814; doi: 10.1016/j.celrep.2019.11.062.

Koncha RR, Ramachandran G, Sepuri NBV, et al. CCCP-induced mitochondrial dysfunction - characterization and analysis of integrated stress response to cellular signaling and homeostasis. *FEBS J* 2021;288(19):5737-5754; doi: 10.1111/febs.15868.

Krishnamoorthy J, Tenkerian C, Gupta J, et al. Downregulation of PERK activity and eIF2alpha serine 51 phosphorylation by mTOR complex 1 elicits pro-oxidant and pro-death effects in tuberous sclerosis-deficient cells. *Cell Death Dis* 2018;9(3):254; doi: 10.1038/s41419-018-0326-2.

Kristensen LS, Andersen MS, Stagsted LVW, et al. The biogenesis, biology and characterization of circular RNAs. *Nat Rev Genet* 2019;20(11):675-691; doi: 10.1038/s41576-019-0158-7.

Kubyshev V, Acevedo-Rocha CG, Budisa N. On universal coding events in protein biogenesis. *Biosystems* 2018;164:16-25; doi: 10.1016/j.biosystems.2017.10.004.

Kumar S, Mohapatra T. Deciphering Epitranscriptome: Modification of mRNA Bases Provides a New Perspective for Post-transcriptional Regulation of Gene Expression. *Front Cell Dev Biol* 2021;9:628415; doi: 10.3389/fcell.2021.628415.

Larosa V, Remacle C. Insights into the respiratory chain and oxidative stress. *Biosci Rep* 2018;38(5); doi: 10.1042/BSR20171492.

Leger A, Amaral PP, Pandolfi L, et al. RNA modifications detection by comparative Nanopore direct RNA sequencing. *Nat Commun* 2021;12(1):7198; doi: 10.1038/s41467-021-27393-3.

Leibovitch M, Topisirovic I. Dysregulation of mRNA translation and energy metabolism in cancer. *Adv Biol Regul* 2018;67:30-39; doi: 10.1016/j.jbior.2017.11.001.

Leppek K, Das R, Barna M. Functional 5' UTR mRNA structures in eukaryotic translation regulation and how to find them. *Nat Rev Mol Cell Biol* 2018;19(3):158-174; doi: 10.1038/nrm.2017.103.

Li G, Scull C, Ozcan L, et al. NADPH oxidase links endoplasmic reticulum stress, oxidative stress, and PKR activation to induce apoptosis. *J Cell Biol* 2010;191(6):1113-1125; doi: 10.1083/jcb.201006121.

Li L, Ni Z, Si X, et al. Emerging clues of regulatory roles of circular RNAs through modulating oxidative stress: Focus on neurological and vascular diseases. *Oxid Med Cell Longev* 2021;2021:6659908; doi: 10.1155/2021/6659908.

Li S, Hu GF. Angiogenin-mediated rRNA transcription in cancer and neurodegeneration. *Int J Biochem Mol Biol* 2010;1(1):26-35.

Li X-X, Xiao L, Chung HK, et al. Interaction between HuR and circPABPN1 modulates autophagy in the intestinal epithelium by altering ATG16L1 translation. *Mol Cell Biol* 2020;40(6); doi: 10.1128/MCB.00492-19.

Li X, Peng J, Yi C. Transcriptome-Wide Mapping of N (1)-Methyladenosine Methylome. *Methods Mol Biol* 2017;1562:245-255; doi: 10.1007/978-1-4939-6807-7\_16.

Li X, Xiong X, Wang K, et al. Transcriptome-wide mapping reveals reversible and dynamic N(1)-methyladenosine methylome. *Nat Chem Biol* 2016;12(5):311-316; doi: 10.1038/nchembio.2040.

Li X, Zhu P, Ma S, et al. Chemical pulldown reveals dynamic pseudouridylation of the mammalian transcriptome. *Nat Chem Biol* 2015;11(8):592-597; doi: 10.1038/nchembio.1836.

Li Z, Malla S, Shin B, et al. Battle against RNA oxidation: molecular mechanisms for reducing oxidized RNA to protect cells. *Wiley Interdiscip Rev RNA* 2014;5(3):335-346; doi: 10.1002/wrna.1214.

Lin Y-C, Lee Y-C, Chang K-L, et al. Analysis of common targets for circular RNAs. *BMC Bioinformatics* 2019;20(1):372; doi: 10.1186/s12859-019-2966-3.

Lombardi S, Testa MF, Pinotti M, et al. Translation termination codons in protein synthesis and disease. *Adv Protein Chem Struct Biol* 2022;132:1-48; doi: 10.1016/bs.apcsb.2022.06.001.

Long F, Lin Z, Li L, et al. Comprehensive landscape and future perspectives of circular RNAs in colorectal cancer. *Mol Cancer* 2021;20(1):26; doi: 10.1186/s12943-021-01318-6.

Lyons SM, Kharel P, Akiyama Y, et al. eIF4G has intrinsic G-quadruplex binding activity that is required for tiRNA function. *Nucleic Acids Res* 2020;48(11):6223-6233; doi: 10.1093/nar/gkaa336.

Malin D, Lee Y, Chepikova O, et al. Methionine restriction exposes a targetable redox vulnerability of triple-negative breast cancer cells by inducing thioredoxin reductase. *Breast Cancer Res Treat* 2021;190(3):373-387; doi: 10.1007/s10549-021-06398-y.

Mathlin J, Le Pera L, Colombo T. A Census and Categorization Method of Epitranscriptomic Marks. *Int J Mol Sci* 2020;21(13); doi: 10.3390/ijms21134684.

McEwen E, Kedersha N, Song B, et al. Heme-regulated inhibitor kinase-mediated phosphorylation of eukaryotic translation initiation factor 2 inhibits translation, induces stress granule formation, and mediates survival upon arsenite exposure. *J Biol Chem* 2005;280(17):16925-16933; doi: 10.1074/jbc.M412882200.

McGillivray P, Ault R, Pawashe M, et al. A comprehensive catalog of predicted functional upstream open reading frames in humans. *Nucleic Acids Res* 2018;46(7):3326-3338; doi: 10.1093/nar/gky188.

McMahon M, Contreras A, Holm M, et al. A single H/ACA small nucleolar RNA mediates tumor suppression downstream of oncogenic RAS. *Elife* 2019;8; doi: 10.7554/eLife.48847.

Meyer KD, Patil DP, Zhou J, et al. 5' UTR m(6)A Promotes Cap-Independent Translation. *Cell* 2015;163(4):999-1010; doi: 10.1016/j.cell.2015.10.012.



Milo R, Jorgensen P, Moran U, et al. BioNumbers--the database of key numbers in molecular and cell biology. *Nucleic Acids Res* 2010;38(Database issue):D750-753; doi: 10.1093/nar/gkp889.

Moradian H, Roch T, Anthofer L, et al. Chemical modification of uridine modulates mRNA-mediated proinflammatory and antiviral response in primary human macrophages. *Mol Ther Nucleic Acids* 2022;27:854-869; doi: 10.1016/j.omtn.2022.01.004.

Natchiar SK, Myasnikov AG, Hazemann I, et al. Visualizing the Role of 2'-OH rRNA Methylations in the Human Ribosome Structure. *Biomolecules* 2018;8(4); doi: 10.3390/biom8040125.

Nemmiche S. Oxidative Signaling Response to Cadmium Exposure. *Toxicol Sci* 2017;156(1):4-10; doi: 10.1093/toxsci/kfw222.

Nunomura A, Moreira PI, Castellani RJ, et al. Oxidative damage to RNA in aging and neurodegenerative disorders. *Neurotox Res* 2012;22(3):231-248; doi: 10.1007/s12640-012-9331-x.

Pakos-Zebrucka K, Koryga I, Mnich K, et al. The integrated stress response. *EMBO Rep* 2016;17(10):1374-1395; doi: 10.15252/embr.201642195.

Panse VG, Johnson AW. Maturation of eukaryotic ribosomes: acquisition of functionality. *Trends Biochem Sci* 2010;35(5):260-266; doi: 10.1016/j.tibs.2010.01.001.

Papp LV, Lu J, Striebel F, et al. The redox state of SECIS binding protein 2 controls its localization and selenocysteine incorporation function. *Mol Cell Biol* 2006;26(13):4895-4910; doi: 10.1128/MCB.02284-05.

Parlato R, Liss B. How Parkinson's disease meets nucleolar stress. *Biochim Biophys Acta* 2014;1842(6):791-797; doi: 10.1016/j.bbadis.2013.12.014.

Pestova TV, Kolupaeva VG. The roles of individual eukaryotic translation initiation factors in ribosomal scanning and initiation codon selection. *Genes Dev* 2002;16(22):2906-2922; doi: 10.1101/gad.1020902.

Phizicky EM, Hopper AK. tRNA biology charges to the front. *Genes Dev* 2010;24(17):1832-1860; doi: 10.1101/gad.1956510.

Pisarev AV, Skabkin MA, Pisareva VP, et al. The role of ABCE1 in eukaryotic posttermination ribosomal recycling. *Mol Cell* 2010;37(2):196-210; doi: 10.1016/j.molcel.2009.12.034.

Pratanwanich PN, Yao F, Chen Y, et al. Identification of differential RNA modifications from nanopore direct RNA sequencing with xPore. *Nat Biotechnol* 2021;39(11):1394-1402; doi: 10.1038/s41587-021-00949-w.

Prats A-C, David F, Diallo LH, et al. Circular RNA, the Key for Translation. *Int J Mol Sci* 2020;21(22); doi: 10.3390/ijms21228591.

Qin S, Zhang Q, Xu Y, et al. m(6)A-modified circRNAs: detections, mechanisms, and prospects in cancers. *Mol Med* 2022;28(1):79; doi: 10.1186/s10020-022-00505-5.

Rajesh K, Krishnamoorthy J, Kazimierczak U, et al. Phosphorylation of the translation initiation factor eIF2alpha at serine 51 determines the cell fate decisions of Akt in response to oxidative stress. *Cell Death Dis* 2015;6:e1591; doi: 10.1038/cddis.2014.554.

Reczek CR, Chandel NS. ROS-dependent signal transduction. *Curr Opin Cell Biol* 2015;33:8-13; doi: 10.1016/j.ceb.2014.09.010.

Roundtree IA, Evans ME, Pan T, et al. Dynamic RNA Modifications in Gene Expression Regulation. *Cell* 2017;169(7):1187-1200; doi: 10.1016/j.cell.2017.05.045.

Rubio Gomez MA, Ibba M. Aminoacyl-tRNA synthetases. *RNA* 2020;26(8):910-936; doi: 10.1261/rna.071720.119.

Safra M, Sas-Chen A, Nir R, et al. The m1A landscape on cytosolic and mitochondrial mRNA at single-base resolution. *Nature* 2017;551(7679):251-255; doi: 10.1038/nature24456.

Sapio RT, Burns CJ, Pestov DG. Effects of Hydrogen Peroxide Stress on the Nucleolar Redox Environment and Pre-rRNA Maturation. *Front Mol Biosci* 2021;8:678488; doi: 10.3389/fmolb.2021.678488.

Saxton RA, Sabatini DM. mTOR Signaling in Growth, Metabolism, and Disease. *Cell* 2017;168(6):960-976; doi: 10.1016/j.cell.2017.02.004.

Schieber M, Chandel NS. ROS function in redox signaling and oxidative stress. *Curr Biol* 2014;24(10):R453-462; doi: 10.1016/j.cub.2014.03.034.

Schimmel P. The emerging complexity of the tRNA world: mammalian tRNAs beyond protein synthesis. *Nat Rev Mol Cell Biol* 2018;19(1):45-58; doi: 10.1038/nrm.2017.77.

Schwartz S, Bernstein DA, Mumbach MR, et al. Transcriptome-wide mapping reveals widespread dynamic-regulated pseudouridylation of ncRNA and mRNA. *Cell* 2014;159(1):148-162; doi: 10.1016/j.cell.2014.08.028.

Sengupta S, Peterson TR, Sabatini DM. Regulation of the mTOR complex 1 pathway by nutrients, growth factors, and stress. *Mol Cell* 2010;40(2):310-322; doi: 10.1016/j.molcel.2010.09.026.

Shcherbik N, Pestov DG. The Impact of Oxidative Stress on Ribosomes: From Injury to Regulation. *Cells* 2019;8(11); doi: 10.3390/cells8111379.

Shedlovskiy D, Zinskie JA, Gardner E, et al. Endonucleolytic cleavage in the expansion segment 7 of 25S rRNA is an early marker of low-level oxidative stress in yeast. *J Biol Chem* 2017;292(45):18469-18485; doi: 10.1074/jbc.M117.800003.

Shen Q, Chu FF, Newburger PE. Sequences in the 3'-untranslated region of the human cellular glutathione peroxidase gene are necessary and sufficient for selenocysteine incorporation at the UGA codon. *J Biol Chem* 1993;268(15):11463-11469.

Shi H, Wei J, He C. Where, When, and How: Context-Dependent Functions of RNA Methylation Writers, Readers, and Erasers. *Mol Cell* 2019;74(4):640-650; doi: 10.1016/j.molcel.2019.04.025.

Sloan KE, Warda AS, Sharma S, et al. Tuning the ribosome: The influence of rRNA modification on eukaryotic ribosome biogenesis and function. *RNA Biol* 2017;14(9):1138-1152; doi: 10.1080/15476286.2016.1259781.

Spriggs KA, Stoneley M, Bushell M, et al. Re-programming of translation following cell stress allows IRES-mediated translation to predominate. *Biol Cell* 2008;100(1):27-38; doi: 10.1042/BC20070098.

Sun J, Cheng B, Su Y, et al. The Potential Role of m6A RNA Methylation in the Aging Process and Aging-Associated Diseases. *Front Genet* 2022;13:869950; doi: 10.3389/fgene.2022.869950.

Suragani RN, Zachariah RS, Velazquez JG, et al. Heme-regulated eIF2alpha kinase activated Atf4 signaling pathway in oxidative stress and erythropoiesis. *Blood* 2012;119(22):5276-5284; doi: 10.1182/blood-2011-10-388132.

Suzuki T. The expanding world of tRNA modifications and their disease relevance. *Nat Rev Mol Cell Biol* 2021;22(6):375-392; doi: 10.1038/s41580-021-00342-0.

Szaflarski W, Lesniczak-Staszak M, Sowinski M, et al. Early rRNA processing is a stress-dependent regulatory event whose inhibition maintains nucleolar integrity. *Nucleic Acids Res* 2022;50(2):1033-1051; doi: 10.1093/nar/gkab1231.

Szwed M, Sonstevold T, Overbye A, et al. Small variations in nanoparticle structure dictate differential cellular stress responses and mode of cell death. *Nanotoxicology* 2019;13(6):761-782; doi: 10.1080/17435390.2019.1576238.

Tang H, Li J, Liu X, et al. Down-regulation of HSP60 Suppresses the Proliferation of Glioblastoma Cells via the ROS/AMPK/mTOR Pathway. *Sci Rep* 2016;6:28388; doi: 10.1038/srep28388.

Thompson DM, Lu C, Green PJ, et al. tRNA cleavage is a conserved response to oxidative stress in eukaryotes. *RNA* 2008;14(10):2095-2103; doi: 10.1261/rna.1232808.

Thoreen CC, Chantranupong L, Keys HR, et al. A unifying model for mTORC1-mediated regulation of mRNA translation. *Nature* 2012;485(7396):109-113; doi: 10.1038/nature11083.

Torres AG, Batlle E, Ribas de Pouplana L. Role of tRNA modifications in human diseases. *Trends Mol Med* 2014;20(6):306-314; doi: 10.1016/j.molmed.2014.01.008.

Tsai K, Jaguva Vasudevan AA, Martinez Campos C, et al. Acetylation of Cytidine Residues Boosts HIV-1 Gene Expression by Increasing Viral RNA Stability. *Cell Host Microbe* 2020;28(2):306-312 e306; doi: 10.1016/j.chom.2020.05.011.

Tsuji T, Sun Y, Kishimoto K, et al. Angiogenin is translocated to the nucleus of HeLa cells and is involved in ribosomal RNA transcription and cell proliferation. *Cancer Res* 2005;65(4):1352-1360; doi: 10.1158/0008-5472.CAN-04-2058.

Verduci L, Tarcitano E, Strano S, et al. CircRNAs: role in human diseases and potential use as biomarkers. *Cell Death Dis* 2021;12(5):468; doi: 10.1038/s41419-021-03743-3.

Verfaillie T, van Vliet A, Garg AD, et al. Pro-apoptotic signaling induced by photo-oxidative ER stress is amplified by Noxa, not Bim. *Biochem Biophys Res Commun* 2013;438(3):500-506; doi: 10.1016/j.bbrc.2013.07.107.

Vindry C, Ohlmann T, Chavatte L. Translation regulation of mammalian selenoproteins. *Biochim Biophys Acta Gen Subj* 2018;1862(11):2480-2492; doi: 10.1016/j.bbagen.2018.05.010.

Vo JN, Cieslik M, Zhang Y, et al. The landscape of circular RNA in cancer. *Cell* 2019;176(4):869-881.e813; doi: 10.1016/j.cell.2018.12.021.

Wang G, Zhang M, Zhang Y, et al. NAT10-mediated mRNA N4-acetylcytidine modification promotes bladder cancer progression. *Clin Transl Med* 2022a;12(5):e738; doi: 10.1002/ctm2.738.

Wang Y, Weng Q, Ge J, et al. tRNA-derived small RNAs: Mechanisms and potential roles in cancers. *Genes Dis* 2022b;9(6):1431-1442; doi: 10.1016/j.gendis.2021.12.009.

Wang Y, Zhao Y, Bollas A, et al. Nanopore sequencing technology, bioinformatics and applications. *Nat Biotechnol* 2021;39(11):1348-1365; doi: 10.1038/s41587-021-01108-x.

Warner JR. The economics of ribosome biosynthesis in yeast. *Trends Biochem Sci* 1999;24(11):437-440; doi: 10.1016/s0968-0004(99)01460-7.

Wilkinson E, Cui YH, He YY. Context-Dependent Roles of RNA Modifications in Stress Responses and Diseases. *Int J Mol Sci* 2021;22(4); doi: 10.3390/ijms22041949.

Wilkinson E, Cui YH, He YY. Roles of RNA Modifications in Diverse Cellular Functions. *Front Cell Dev Biol* 2022;10:828683; doi: 10.3389/fcell.2022.828683.

Willi J, Kupfer P, Evequoz D, et al. Oxidative stress damages rRNA inside the ribosome and differentially affects the catalytic center. *Nucleic Acids Res* 2018;46(4):1945-1957; doi: 10.1093/nar/gkx1308.

Wisniewski JR, Hein MY, Cox J, et al. A "proteomic ruler" for protein copy number and concentration estimation without spike-in standards. *Mol Cell Proteomics* 2014;13(12):3497-3506; doi: 10.1074/mcp.M113.037309.

Wu F, Zhang L, Lai C, et al. Dynamic Alteration Profile and New Role of RNA m6A Methylation in Replicative and H<sub>2</sub>O<sub>2</sub>-Induced Premature Senescence of Human Embryonic Lung Fibroblasts. *Int J Mol Sci* 2022;23(16); doi: 10.3390/ijms23169271.

Wu Q, Wang Y, Cao M, et al. Homology-independent discovery of replicating pathogenic circular RNAs by deep sequencing and a new computational algorithm. *Proc Natl Acad Sci U S A* 2012;109(10):3938-3943; doi: 10.1073/pnas.1117815109.

Xuan JJ, Sun WJ, Lin PH, et al. RMBase v2.0: deciphering the map of RNA modifications from epitranscriptome sequencing data. *Nucleic Acids Res* 2018;46(D1):D327-D334; doi: 10.1093/nar/gkx934.

Xue S, Barna M. Specialized ribosomes: a new frontier in gene regulation and organismal biology. *Nat Rev Mol Cell Biol* 2012;13(6):355-369; doi: 10.1038/nrm3359.

Yamasaki S, Ivanov P, Hu GF, et al. Angiogenin cleaves tRNA and promotes stress-induced translational repression. *J Cell Biol* 2009;185(1):35-42; doi: 10.1083/jcb.200811106.

Yang B, Chen Q. Cross-Talk between Oxidative Stress and m(6)A RNA Methylation in Cancer. *Oxid Med Cell Longev* 2021;2021:6545728; doi: 10.1155/2021/6545728.

Yang C, Yuan W, Yang X, et al. Circular RNA circ-ITCH inhibits bladder cancer progression by sponging miR-17/miR-224 and regulating p21, PTEN expression. *Mol Cancer* 2018;17(1):19; doi: 10.1186/s12943-018-0771-7.

Yang Y, Fan X, Mao M, et al. Extensive translation of circular RNAs driven by N6-methyladenosine. *Cell Res* 2017;27(5):626-641; doi: 10.1038/cr.2017.31.

Yang Y, Wang Z. IRES-mediated cap-independent translation, a path leading to hidden proteome. *J Mol Cell Biol* 2019;11(10):911-919; doi: 10.1093/jmcb/mjz091.

Yoon A, Peng G, Brandenburger Y, et al. Impaired control of IRES-mediated translation in X-linked dyskeratosis congenita. *Science* 2006;312(5775):902-906; doi: 10.1126/science.1123835.

Young SK, Wek RC. Upstream Open Reading Frames Differentially Regulate Gene-specific Translation in the Integrated Stress Response. *J Biol Chem* 2016;291(33):16927-16935; doi: 10.1074/jbc.R116.733899.

Zahia TH, Yona L, Anne-Laure B, et al. Selective up-regulation of human selenoproteins in response to oxidative stress. *Free Radic Biol Med* 2014;75 Suppl 1:S25; doi: 10.1016/j.freeradbiomed.2014.10.745.

Zeng Y, Wang L, Zhou Y, et al. NMDA receptor antagonists engender neuroprotection against gp120-induced cognitive dysfunction in rats through modulation of PKR activation, oxidative stress, ER stress and IRE1alpha signal pathway. *Eur J Neurosci* 2022;56(2):3806-3824; doi: 10.1111/ejn.15688.

Zhang J, Yan S, Chang L, et al. Direct microRNA Sequencing Using Nanopore-Induced Phase-Shift Sequencing. *iScience* 2020;23(3):100916; doi: 10.1016/j.isci.2020.100916.

Zhang R, Zhang Y, Guo F, et al. RNA N6-Methyladenosine Modifications and Its Roles in Alzheimer's Disease. *Front Cell Neurosci* 2022a;16:820378; doi: 10.3389/fncel.2022.820378.

Zhang W, Sui Y. CircBPTF knockdown ameliorates high glucose-induced inflammatory injuries and oxidative stress by targeting the miR-384/LIN28B axis in human umbilical vein endothelial cells. *Mol Cell Biochem* 2020;471(1-2):101-111; doi: 10.1007/s11010-020-03770-2.

Zhang X, Wang S, Wang H, et al. Circular RNA circNRIP1 acts as a microRNA-149-5p sponge to promote gastric cancer progression via the AKT1/mTOR pathway. *Mol Cancer* 2019;18(1):20; doi: 10.1186/s12943-018-0935-5.

Zhang Y, Jing Y, Wang Y, et al. NAT10 promotes gastric cancer metastasis via N4-acetylated COL5A1. *Signal Transduct Target Ther* 2021;6(1):173; doi: 10.1038/s41392-021-00489-4.

Zhang Y, Lu L, Li X. Detection technologies for RNA modifications. *Exp Mol Med* 2022b;54(10):1601-1616; doi: 10.1038/s12276-022-00821-0.

Zhao T, Li X, Sun D, et al. Oxidative stress: One potential factor for arsenite-induced increase of N(6)-methyladenosine in human keratinocytes. *Environ Toxicol Pharmacol* 2019;69:95-103; doi: 10.1016/j.etap.2019.04.005.

Zheng Q, Bao C, Guo W, et al. Circular RNA profiling reveals an abundant circHIPK3 that regulates cell growth by sponging multiple miRNAs. *Nat Commun* 2016;7(1):11215; doi: 10.1038/ncomms11215.


Zhu HL, Shi XT, Xu XF, et al. Melatonin protects against environmental stress-induced fetal growth restriction via suppressing ROS-mediated GCN2/ATF4/BNIP3-dependent mitophagy in placental trophoblasts. *Redox Biol* 2021;40:101854; doi: 10.1016/j.redox.2021.101854.




**Table 1: Software for detection of RNA modifications from nanopore direct RNA sequencing data.**

Tool	Modifications	Sample requirements	Reference
m6Anet	m6A	Single sample	(Hendra et al., 2022)
nanocompore	agnostic	Case-control	(Leger et al., 2021)
xpore	agnostic	Case-control	(Pratanwanich et al., 2021)
nanoRMS	$\Psi$ and agnostic	Single sample (only $\Psi$ ) or case control	(Begik et al., 2021)
eligos2	agnostic	Single sample (with pre-trained model) or case control	(Jenjaroenpun et al., 2021)


## Figure Legends




**Figure 1:** Schematic of the non-coding elements and pathways that regulate translation of mRNAs under stress and that are discussed in this review. All underlined elements are discussed.




**Figure 2:** Schematic of rRNA processing. The 18S, 28S, and 5.8S rRNAs are transcribed by Pol I in the nucleolus, a sub region of the nucleus, as a single precursor transcript and extensively processed to their mature forms; 5S rRNA is transcribed separately and imported into the nucleolus. There, ribosomal proteins and mature rRNAs assemble into the ribosomal subunits that are then utilized in translation. ROS can inhibit rRNA transcription and processing resulting in reduced biogenesis and thus reduced translation.




**Figure 3:** Overview for generation for stress-induced tRNA fragments (tiRNA). During stress, angiogenin translocates from the nucleus to the cytoplasm and cleaves tRNAs in the anticodon loop. The resulting RNA molecules, termed tiRNAs, can then perform functions that include regulation of translation.



**Figure 4:** Schematic of circRNA function in cells. CircRNAs can act as sponges for RNA binding proteins and miRNAs and thereby altering the translation of their targets. CircRNAs can also be translated themselves via an IRES that become increasingly utilized for translation initiation during stress-dependent inhibition of cap-dependent translation.



**Figure 5: A)** Schematic of the mTORC1 pathway. The activity of the serine/threonine kinase mTORC1 is regulated by cellular growth conditions. When nutrients, energy, and growth factors are in abundance, the mTORC1 complex is activated and promotes translation by phosphorylating S6K, which increases its activity, and by phosphorylating eIF4EBP1, which inhibits its activity. Transcripts with a 5' terminal oligopyrimidine (TOP) motif are more sensitive to eIF4EBP1 and thus mTORC1 activity. **B)** Schematic of the integrated stress response (ISR). The ISR responds to a diverse array of stressors including iron deprivation, viral infection, proteotoxic stress and starvation. A set of four kinases respond to their respective stimuli and a response is integrated into phosphorylation of the translation initiation factor eIF2a, which reduces its affinity for a guanine exchange factor. Without GTP, eIF2a does not participate in translation initiation thereby reducing global translation and increasing ribosome scanning, which leads to the selective translation of the critical ISR transcription factor ATF4.



**Figure 6:** Schematic of RNA modifications and their detection by nanopore RNA sequencing. The function of mRNA modifications is in part dependent on the region they are located with consequences ranging from altered mRNA stability to changes in translation efficiency. mRNA modifications can be detected by nanopore RNA sequencing since RNA bases are directly measured through electrical resistance as they are enzymatically forced through the nanopore. In this way, nanopore sequencing sufficiently trained on different mRNA modifications can offer the most authentic view of the epitranscriptome available.