



## **BioGears: Drug Modeling Overview**

A. Baird<sup>1</sup>, J. Carter<sup>1</sup>, L. Marin<sup>1</sup>, M. McDaniel<sup>1</sup>, N.  
Tatum<sup>1</sup>, S. White<sup>1</sup>

*1. Applied Research Associates Inc.*

# Outline

## BioGears Background

- Lumped-parameter approach
- System overview

## Methods: PK / PD Methodology

- Substance definitions
- Pharmacokinetic model
- Pharmacodynamic model


## Results: Complex Treatment Scenarios

- Development
- Results

Technical overview of the BioGears engine

# BACKGROUND

# Background



# Background

## The Cardiovascular Circuit

- Time-varying compliance in heart mimics pumping
- Adapted from Stergiopoulos et al
- Pathways to tissue compartments


Define the change in elastance to be:

$$E_v(E_{\max,v} - E_{\min,v})\left(\frac{f(t)}{f_{\max}}\right) + E_{\min,v}$$

Double hill time-varying relationship:



$$f(t) = \left[ \frac{\left(\frac{t}{\alpha_1 T}\right)^{n_1}}{1 + \left(\frac{t}{\alpha_1 T}\right)^{n_1}} \right] \left[ \frac{1}{1 + \left(\frac{t}{\alpha_2 T}\right)^{n_2}} \right]$$

Set the left and right heart compliance to be the inverse of elastance



# Background



## Pressure volume curve over time



# Background

## The Respiratory Circuit

- Variable pressure source drives air in and out of lungs
- Alveolar transfer of blood gases
- Adopted from model by Albanese et al.
- Driver pressure is coupled to chemoreceptors



$$\dot{D}_x = \frac{D_{cap,O_2} \cdot C_{D,x} \cdot (P_{x,a} - P_{x, cap}) \cdot SA_a}{d}$$

$D_{cap,O_2}$  = Oxygen diffusive capacity

$C_{D,x}$  = Diffusivity of gas x relative to  $O_2$

$SA_a$  = Alveolar surface area

BioGears drug model description

# METHODS




# Methods

## Administration routes in BioGears include

- Intravenous/intraarterial (infusion or bolus)
- Intramuscular
- Oral
- Inhaled

## Current supported/validated drugs in BioGears

Prednisone	Desflurane	Morphine	Midazolam	Fentanyl
Naloxone	Insulin	Succinylcholine	Rocuronium	Albuterol
Epinephrine	Sarin	Propofol	Ketamine	Pralidoxime
Acetaminophen	Tranexamic acid	Vasopressin	Furosemide	




# Methods: Partition Coefficient

**Perfusion-limited diffusion assumed for each organ:**

$$\frac{dm}{dt} = Q_B \left( C_B - \frac{C_T}{K_{TB}} \right)$$

- Q is flow rate,
- C denotes the concentration in blood and tissue,
- K is the given tissues computed partition coefficient



# Methods: Partition Coefficient

Moderate to strong bases (Rogers, Leahy, and Rowland, 2004)

$$K_{TB} = \frac{f_u}{\lambda_{BP}} \left[ f_{EW} + \frac{1 + 10^{pK_a - pH_{IW}}}{1 + 10^{pK_a - pH_p}} f_{IW} + \frac{K_a [AP^-]_T \cdot 10^{pK_a - pH_{IW}}}{1 + 10^{pK_a - pH_p}} + \frac{P \cdot f_{NL} + (0.3P + 0.7)f_{NP}}{1 + 10^{pK_a - pH_p}} \right]$$

Acids, neutral compounds, weak bases (Rogers and Rowland, 2005)

$$K_{TB} = \frac{f_u}{\lambda_{BP}} \left[ f_{EW} + \frac{X \cdot f_{IW}}{Y} + \frac{P \cdot f_{NL} + (0.3P + 0.7)f_{NP}}{Y} + K_{Bind} \left( \frac{1}{f_u} - 1 - \left( \frac{P \cdot f_{NL} + (0.3P + 0.7)f_{NP}}{Y} \right) \right) \right]$$


$$X = \begin{cases} 1 + 10^{pH_{IW} - pK_a} & \text{Acid} \\ 1 + 10^{pK_a - pH_{IW}} & \text{Weak base} \\ 1 & \text{Neutral} \end{cases}$$

$$Y = \begin{cases} 1 + 10^{pH_p - pK_a} & \text{Acid} \\ 1 + 10^{pK_a - pH_p} & \text{Weak base} \\ 1 & \text{Neutral} \end{cases}$$


# Methods: Clearance

$Cl_I$ ,  $Cl_R$ ,  $Cl_S$ , and  $f_u$  defined in substance schema and are:

- Hepatic
- Renal
- Systemic
- Fraction unbound in plasma



# Methods: Pharmacodynamics




# Methods


## Consequences of effect compartment

- Example: More rapid onset/offset of Fentanyl compared to Morphine


### *Fentanyl*




### *Morphine*



# Methods



4 hour Plasma Concentration Curve




## PK

- Perfusion-limited diffusion
- Physiochemical properties used to calculate partition coefficients,  $K_p$
- **Or** partition coefficient can be an input
- Renal, Hepatic, and Systemic Clearance


## PD

- Patient Physiological response is governed by the saturation of the drug at the “effect-site compartment” within the body
  - The rate constant is specific to the drug




Worked with subcontractor UNC Eshelman School of Pharmacy  
Pharmacodynamics also validated through scenario validation  
All drugs validated in this manor

### Fentanyl-Bolus



### Morphine-Bolus





Investigate vasopressin during a severe hemorrhage scenario

# RESULTS

# Substance Test Case

## Vasopressin


- Endogenously produced hormone
- Pharmacological implications for multiple systems
  - Cardiovascular
  - Renal
- Exogenous administration in multiple shock states
  - Distributive shock (e.g. sepsis)
  - Hypovolemic shock

```
<Substance>
  <Name>Vasopressin</Name>
  <MolarMass readOnly="false" unit="g/mol" value="1084.24"/>
  <State>Liquid</State>
  <Clearance>
    <Systemic>
      <FractionExcretedInFeces readOnly="false" value="0"/>
      <FractionUnboundInPlasma readOnly="false" value="0.99"/>
      <IntrinsicClearance readOnly="false" unit="mL/min kg" value="0"/>
      <RenalClearance readOnly="false" unit="mL/min kg" value="1.13"/>
      <SystemicClearance readOnly="false" unit="mL/min kg" value="15.17"/>
    </Systemic>
    <RenalDynamics>
      <Clearance readOnly="false" unit="mL/min kg" value="1.13"/>
    </RenalDynamics>
  </Clearance>
  <Pharmacokinetics>
    <Physicochemicals>
      <AcidDissociationConstant readOnly="false" value="9.8"/>
      <BindingProtein>Albumin</BindingProtein>
      <BloodPlasmaRatio readOnly="false" value="1"/>
      <FractionUnboundInPlasma readOnly="false" value="0.99"/>
      <IonicState>Neutral</IonicState>
      <LogP readOnly="false" value="-4.8"/>
    </Physicochemicals>
  </Pharmacokinetics>
</Substance>
```

# Pharmacokinetic Validation, Part 1

## Scenario Definition

- Modeled after Glanzer, 1982
- Study design
  - Male volunteers
  - Intravenous vasopressin infusion
    - Rate: 160 pmol/min (0.174  $\mu$ g/min)
    - Length: 60 min
  - Post-infusion observation
    - Length: 60 min



# Scenario Background

## Vasopressin as hemorrhagic shock therapeutic

- Rationale
  - Blood pressure support in the case of catecholamine-resistant shock (Beloncle, 2013)
  - Splanchnic vasoconstriction could improve vital organ perfusion (Beloncle, 2013)
    - Decreased fluid requirements during resuscitation
    - Lower risk of tissue edema
  - Some success in animal studies (Voelckel, 2003; Ready et al, 1991)

## Controlled clinical studies provide sparse data

- Vasopressin in Traumatic Shock (VITRIS) Study (Cohn et al, 2010)
  - Exploratory, not enough participants enrolled
- University Hospital in San Antonio (Cohn et al, 2011)
  - Treatment: 0.2 Unit/min vasopressin
  - Placebo: Saline
  - Conclusion: Patients in vasopressin group required fewer fluids but no difference in 30-day mortality found
  - No patient metrics recorded


## Case for BioGears

- Results from *in silico* experiments can provide direction for future studies
- Tool for medical training

# BioGears Vasopressin Treatment Scenario

## Treatment protocols similar to Cohn et al, 2011





- 0.5 mL / min saline (control)
- 1 Unit / min vasopressin (1 U = 2.5  $\mu$ g)
  - Administered at same volumetric rate as control



# Complex Scenario: Results

## Treatment protocols similar to Cohn et al, 2011

- 0.5 mL / min saline (control)
- 1 Unit / min vasopressin (1 U = 2.5 µg)
- Administered at same volumetric rate as control



# Complex Scenario: Comments

## Pre-treatment

- BioGears captures feedback responses to massive blood loss
  - Baroreceptor-mediated increase in heart rate
  - Decrease in glomerular filtration (and urine output)

## Treatment

- Less fluid required to achieve MAP increase
- Tachycardia reversal
- Increased urine output indicative of end-organ perfusion

## Possible improvements

- Baroreceptor decompensation (Foex, 1999)
- Site-of-action pharmacodynamics
  - Vasopressin-mediated vasoconstriction occurs mainly in liver, smooth muscle, and peripheral tissue (Ready, 1991)

# Conclusions: BioGears PK/PD System

## Current Capabilities

- Whole-body, physiologically-based pharmacokinetic model
- Wide range of pharmacologic effects with potential for delayed-onset
- Multiple routes of administration
- Extendable to new drugs
- Integration with BioGears Physiology Engine promotes simulation of complex medical training scenarios
- Graphical User Interface (GUI) removes coding burden from users
- Ability to create simulated clinical studies and treatment protocols



# References

1. Rodgers, Trudy, David Leahy, and Malcolm Rowland. Physiologically based pharmacokinetic modeling 1: predicting the tissue distribution of moderate-to-strong bases. *Journal of pharmaceutical sciences* 94.6 (2005): 1259-1276.
2. Rodgers, Trudy, and Malcolm Rowland. Physiologically based pharmacokinetic modelling 2: predicting the tissue distribution of acids, very weak bases, neutrals and zwitterions. *Journal of pharmaceutical sciences* 95.6 (2006): 1238-1257.
3. Glänzer, K., Prüßing, B., Düsing, R., and Kramer, H. J. Hemodynamic and hormonal responses to 8-arginine-vasopressin in healthy man: effects of indomethacin. *Klinische Wochenschrift*, 60, 1234-1239 (1982).
4. Ebert, T. J., Cowley, A.W. and Skelton, M. Vasopressin reduces cardiac function and augments cardiopulmonary baroreflex resistance increases in man. *The Journal of clinical investigation*, 77, 1136-1142 (1986).
5. Morton, J. J., Padfield, P. L., and FORSLING, M. L. A radioimmunoassay for plasma arginine-vasopressin in man and dog: application to physiological and pathological states. *Journal of Endocrinology*, 65, 411-424 (1975).
6. Andersen, L.J., Andersen, J.L., Schutten, H.J., Warberg, J. and Bie, P. Antidiuretic effect of subnormal levels of arginine vasopressin in normal humans. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, 259(1), R53-R60 (1990).
7. Beloncle, F., Meziani, F., Lerolle, N., Radermacher, P. and Asfar, P. Does vasopressor therapy have an indication in hemorrhagic shock?. *Annals of intensive care*, 3(1), 13 (2013).
8. Voelckel, W. G, et al. Arginine vasopressin, but not epinephrine, improves survival in uncontrolled hemorrhagic shock after liver trauma in pigs. *Critical care medicine*, 31, 1160-1165 (2003).
9. Ready, J.B., Robertson, A.D. and Rector, W.G. Effects of vasopressin on portal pressure during hemorrhage from esophageal varices. *Gastroenterology*, 100, 1411-1416 (1991).
10. Krismer, A. C., et al. Employing vasopressin as an adjunct vasopressor in uncontrolled traumatic hemorrhagic shock. *Der Anaesthetist*, 54, 220-224 (2005).
11. Tsuneyoshi, I., Onomoto, M., Yonetani, A., and Kanmura, Y. Low-dose vasopressin infusion in patients with severe vasodilatory hypotension after prolonged hemorrhage during general anesthesia. *Journal of anesthesia*, 19, 170-173 (2005)
12. Cohn, S. M., Blackburne, L. H., Landry, D. W., Proctor, K. G., Walley, K. R., and Wenzel, V. San Antonio vasopressin in shock symposium report. *Resuscitation*, 81, 1473-1475 (2010).
13. Cohn, S. M., McCarthy, J., Stewart, R. M., Jonas, R. B., Dent, D. L., & Michalek, J. E. Impact of low-dose vasopressin on trauma outcome: prospective randomized study. *World journal of surgery*, 35, 430-439 (2011).
14. Foex, B. A. Systemic responses to trauma. *British medical bulletin*, 55, 726-743 (1999).