

# Monte Carlo methods for expensive models based on repulsive point processes

Rémi Bardenet

CNRS & CRIStAL, Univ. Lille, France





**Figure:** Adrien Hardy, Ayoub Belhadji, Pierre Chainais

Numerical integration and DPPs

Tight interpolation rates in RKHSs

**Numerical integration and DPPs**

Tight interpolation rates in RKHSs

- ▶ The goal is to approximate

$$\int f d\mu = \int f(x)\omega(x)dx \approx \sum_{i=1}^N w_i f(x_i),$$

with nodes  $x_i$  and weights  $w_i$ .

### Monte Carlo integration (importance sampling, MCMC, etc.)

- ▶ Choose the nodes randomly, and the weights  $w_i = w(x_i, x_{-i})$ .
- ▶ Typical error is

$$\forall f \in L^2, \quad \sqrt{\mathbb{E} \left[ \int f d\mu - \sum_{i=1}^N w_i f(x_i) \right]^2} \sim \frac{1}{\sqrt{N}}.$$

- ▶ The goal is to approximate

$$\int f d\mu = \int f(x)\omega(x)dx \approx \sum_{i=1}^N w_i f(x_i),$$

with nodes  $x_i$  and weights  $w_i$ .

## Quasi Monte Carlo integration (Dick and Pillichshammer, 2010)

- ▶ Choose the nodes deterministically, and the weights  $w_i = w(x_i, x_{-i})$ .
- ▶ Typical error is

$$\sup_{f \in \mathcal{F}} \left| \int f d\mu - \sum_{i=1}^N w_i f(x_i) \right| \sim \frac{\log^{s-1} N}{N}.$$

Numerical integration and DPPs

**Tight interpolation rates in RKHSs**

- ▶ Consider the RKHS  $\mathcal{F}$  with kernel  $\kappa$ , i.e. the completion of

$$\left\{ \sum_{i=1}^M \alpha_i \kappa(x_i, \cdot), M \in \mathbb{N}, \alpha_1, \dots, \alpha_n \in \mathbb{R}, x_1, \dots, x_M \in \mathbb{R}^d \right\}.$$

for the inner product defined by  $\langle \kappa(x, \cdot), \kappa(y, \cdot) \rangle_{\mathcal{F}} := \kappa(x, y)$ .

- ▶ Under general assumptions,  $\mathcal{F} \subset L^2(d\mu)$ , is dense, there is an ON basis  $(e_n)$  of  $L^2(d\mu)$  and  $\sigma_n \rightarrow 0$  such that, pointwise,

$$\kappa(x, y) = \sum_{n \geq 1} \sigma_n e_n(x) e_n(y).$$

- ▶ In that case,  $f \in \mathcal{F}$  if and only if  $\sum_n \sigma_n^{-1} |\langle f, e_n \rangle|^2$  converges.



- ▶ Consider the RKHS  $\mathcal{F}$  with kernel  $\kappa$ , i.e. the completion of

$$\left\{ \sum_{i=1}^M \alpha_i \kappa(x_i, \cdot), M \in \mathbb{N}, \alpha_1, \dots, \alpha_n \in \mathbb{R}, x_1, \dots, x_M \in \mathbb{R}^d \right\}.$$

for the inner product defined by  $\langle \kappa(x, \cdot), \kappa(y, \cdot) \rangle_{\mathcal{F}} := \kappa(x, y)$ .

- ▶ Under general assumptions,  $\mathcal{F} \subset L^2(d\mu)$ , is dense, there is an ON basis  $(e_n)$  of  $L^2(d\mu)$  and  $\sigma_n \rightarrow 0$  such that, pointwise,

$$\kappa(x, y) = \sum_{n \geq 1} \sigma_n e_n(x) e_n(y).$$

- ▶ In that case,  $f \in \mathcal{F}$  if and only if  $\sum_n \sigma_n^{-1} |\langle f, e_n \rangle|^2$  converges.

- ▶ Consider the RKHS  $\mathcal{F}$  with kernel  $\kappa$ , i.e. the completion of

$$\left\{ \sum_{i=1}^M \alpha_i \kappa(x_i, \cdot), M \in \mathbb{N}, \alpha_1, \dots, \alpha_n \in \mathbb{R}, x_1, \dots, x_M \in \mathbb{R}^d \right\}.$$

for the inner product defined by  $\langle \kappa(x, \cdot), \kappa(y, \cdot) \rangle_{\mathcal{F}} := \kappa(x, y)$ .

- ▶ Under general assumptions,  $\mathcal{F} \subset L^2(d\mu)$ , is dense, there is an ON basis  $(e_n)$  of  $L^2(d\mu)$  and  $\sigma_n \rightarrow 0$  such that, pointwise,

$$\kappa(x, y) = \sum_{n \geq 1} \sigma_n e_n(x) e_n(y).$$

- ▶ In that case,  $f \in \mathcal{F}$  if and only if  $\sum_n \sigma_n^{-1} |\langle f, e_n \rangle|^2$  converges.

- ▶ Let  $f \in \mathcal{F}$ ,  $g \in L^2(d\mu)$  then

$$\left| \int fg d\mu - \sum_{i=1}^N w_i f(x_i) \right| \leq \|f\|_{\mathcal{F}} \left\| \mu_g - \sum_{i=1}^N w_i \kappa(x_i, \cdot) \right\|_{\mathcal{F}}, \quad (1)$$

where

$$\mu_g = \int g(x) \kappa(x, \cdot) d\mu(x)$$

is the **mean element of  $g$** .

- ▶ Once the nodes  $x_1, \dots, x_N$  are known, minimizing the RHS of (1) in  $w$  boils down to inverting an  $N \times N$  matrix.
- ▶  $\sum_{i=1}^N \hat{w}_i \kappa(x_i, \cdot)$  is the orthogonal projection of  $\mu_g$  onto

$$S = \text{Span}(\kappa(x_1, \cdot), \dots, \kappa(x_N, \cdot)).$$

- ▶ The squared volume in  $S$  of the parallelotope spanned by these functions is  $\det[\kappa(x_i, x_j)]$ .

- ▶ Let  $f \in \mathcal{F}$ ,  $g \in L^2(d\mu)$  then

$$\left| \int fg d\mu - \sum_{i=1}^N w_i f(x_i) \right| \leq \|f\|_{\mathcal{F}} \left\| \mu_g - \sum_{i=1}^N w_i \kappa(x_i, \cdot) \right\|_{\mathcal{F}}, \quad (1)$$

where

$$\mu_g = \int g(x) \kappa(x, \cdot) d\mu(x)$$

is the **mean element of  $g$** .

- ▶ Once the nodes  $x_1, \dots, x_N$  are known, minimizing the RHS of (1) in  $w$  boils down to inverting an  $N \times N$  matrix.
- ▶  $\sum_{i=1}^N \hat{w}_i \kappa(x_i, \cdot)$  is the orthogonal projection of  $\mu_g$  onto

$$S = \text{Span}(\kappa(x_1, \cdot), \dots, \kappa(x_N, \cdot)).$$

- ▶ The squared volume in  $S$  of the parallelotope spanned by these functions is  $\det[\kappa(x_i, x_j)]$ .

- ▶ Let  $f \in \mathcal{F}$ ,  $g \in L^2(d\mu)$  then

$$\left| \int fg d\mu - \sum_{i=1}^N w_i f(x_i) \right| \leq \|f\|_{\mathcal{F}} \left\| \mu_g - \sum_{i=1}^N w_i \kappa(x_i, \cdot) \right\|_{\mathcal{F}}, \quad (1)$$

where

$$\mu_g = \int g(x) \kappa(x, \cdot) d\mu(x)$$

is the **mean element of  $g$** .

- ▶ Once the nodes  $x_1, \dots, x_N$  are known, minimizing the RHS of (1) in  $w$  boils down to inverting an  $N \times N$  matrix.
- ▶  $\sum_{i=1}^N \hat{w}_i \kappa(x_i, \cdot)$  is the orthogonal projection of  $\mu_g$  onto

$$S = \text{Span}(\kappa(x_1, \cdot), \dots, \kappa(x_N, \cdot)).$$

- ▶ The squared volume in  $S$  of the parallelotope spanned by these functions is  $\det[\kappa(x_i, x_j)]$ .

- ▶ Let  $f \in \mathcal{F}$ ,  $g \in L^2(d\mu)$  then

$$\left| \int fg d\mu - \sum_{i=1}^N w_i f(x_i) \right| \leq \|f\|_{\mathcal{F}} \left\| \mu_g - \sum_{i=1}^N w_i \kappa(x_i, \cdot) \right\|_{\mathcal{F}}, \quad (1)$$

where

$$\mu_g = \int g(x) \kappa(x, \cdot) d\mu(x)$$

is the **mean element of  $g$** .


- ▶ Once the nodes  $x_1, \dots, x_N$  are known, minimizing the RHS of (1) in  $w$  boils down to inverting an  $N \times N$  matrix.
- ▶  $\sum_{i=1}^N \hat{w}_i \kappa(x_i, \cdot)$  is the orthogonal projection of  $\mu_g$  onto

$$S = \text{Span}(\kappa(x_1, \cdot), \dots, \kappa(x_N, \cdot)).$$

- ▶ The squared volume in  $S$  of the parallelotope spanned by these functions is  $\det[\kappa(x_i, x_j)]$ .

## Volume sampling

- ▶ Let  $x_1, \dots, x_N \sim Z^{-1} \det[\kappa(x_i, x_j)] d\mu(x_1) \dots d\mu(x_N)$
- ▶ Solve the linear program for the weights  $w_1, \dots, w_N$ .



### Volume sampling

- ▶ Let  $x_1, \dots, x_N \sim Z^{-1} \det[\kappa(x_i, x_j)] d\mu(x_1) \dots d\mu(x_N)$
- ▶ Solve the linear program for the weights  $w_1, \dots, w_N$ .

### Theorem (Belhadji, Bardenet, and Chainais, 2020)

Assume again  $\sum_{n=1}^N |\langle g, e_n \rangle|^2 \leq 1$ . Then

$$\mathbb{E} \left\| \mu_g - \sum_{i=1}^N w_i \kappa(x_i, \cdot) \right\|_{\mathcal{F}}^2 \leq \sigma_N (1 + \beta_N),$$

where  $\beta_N = \min_{M \in [2:N]} [(N - M + 1)\sigma_N]^{-1} \sum_{m \geq M} \sigma_m$ .

- ▶ It is known<sup>1</sup> that  $\inf_{\substack{Y \subset \mathcal{F} \\ \dim Y = N}} \sup_{\|g\|_{d\omega} \leq 1} \inf_{y \in Y} \|\mu_g - y\|_{\mathcal{F}}^2 = \sigma_{N+1}$ .

---

<sup>1</sup>Pinkus, 2012.



### Take-home message

- ▶ Without surprise, more smoothness means faster convergence.
- ▶ Random quadratures can be as fast as possible in RKHSs.

### Many open problems

- ▶ Practical relevance of RKHS hypothesis in applications?
- ▶ What should  $g \in L^2(\mu)$  be in  $\int fg d\mu$ ?
- ▶ Robustness to kernel choice.
- ▶ How do we efficiently sample from continuous volume sampling **without spectral knowledge**? See e.g. Rezaei and Gharan, 2019.
- ▶ Can faster-than-Monte Carlo error come with **limited** smoothness? Yes, see e.g. Bardenet and Hardy, 2020.

- Bardenet, R. and A. Hardy (2020). “Monte Carlo with Determinantal Point Processes”. In: *Annals of Applied Probability*.
- Belhadji, A., R. Bardenet, and P. Chainais (2020). “Kernel interpolation with continuous volume sampling”. In: *International Conference on Machine Learning (ICML)*.
- Dick, J. and F. Pillichshammer (2010). *Digital Nets and Sequences. Discrepancy Theory and Quasi-Monte Carlo Integration*. Cambridge University Press.
- Pinkus, A. (2012). *N-widths in Approximation Theory*. Vol. 7. Springer Science & Business Media.
- Rezaei, A. and S. O. Gharan (2019). “A Polynomial Time MCMC Method for Sampling from Continuous Determinantal Point Processes”. In: *International Conference on Machine Learning*, pp. 5438–5447.