AEGIS: DIFFERENTIABLE MARS CLIMATE MODEL WITH NEURAL CLOSURES

Sameera S Kashyap, Victor Cruz, Angel Yepez & Razvan Marinescu
Department of Computer Science
UC Santa Cruz
Santa Cruz, CA 95064, USA
{ssudars1, vcruz17, ayepezal, ramarine}@ucsc.edu

ABSTRACT

General circulation models (GCMs) are the primary tool for simulating planetary atmospheres. They play a vital role in understanding Mars's atmosphere, as forecasting its unique weather is mission-critical for operations such as entry, descent, and landing. Mars poses unusual challenges for these models, as observations are sparse compared to Earth. In addition, a thin CO_2 atmosphere alongside a radiatively active dust cycle creates a volatile atmosphere with large diurnal temperature swings and no true terrestrial analog for validation. Existing Mars GCMs, including the LMD PCM, the NASA Ames Mars GCM, and PlanetWRF, are mature and physically detailed but are implemented in legacy Fortran with finite-difference or finite-volume solvers, and they do not expose gradients for calibration or machine learning. Here we present AEGIS, a modular differentiable Mars climate model that couples Mars's unique atmospheric physics to the Dinosaur dynamical core, with interfaces for neural closures. We demonstrate coupled forward integration, physical-parameter derivatives, and neural training. Historical seasonal comparisons show terrain-correlated surface pressure but also warm temperature biases and substantial frost and wind errors. The atmosphere–frost inventory is constrained by a numerical projection; its small residual does not establish raw energy or momentum conservation. We separate historical results from a new fixed-update neural-radiation comparison and state the remaining long-run, calibration, and observational validation requirements.

1 INTRODUCTION

General circulation models (GCMs) connect the governing equations of atmospheric motion to the physical processes that shape planetary climate. Established Mars models include the LMD Mars GCM, the NASA Ames Mars GCM, and PlanetWRF (Forget et al., 1999; Haberle et al., 1993; Richardson et al., 2007). On Mars, radiative heating by dust and CO_2, exchange with a thermally active surface, and seasonal condensation onto polar caps jointly shape circulation, pressure, and frost. A useful model must represent these interactions while allowing individual assumptions to be tested (Kahre & Haberle, 2010).

Traditional GCMs contain uncertain parameterization coefficients whose coupled effects can complicate tuning and permit compensating errors (Forget et al., 1999). Parameter sweeps and ensemble calibration are often computationally expensive (McNeall et al., 2020). In addition, the forward-model interface does not generally expose gradients, and maintaining separate adjoint implementations incurs substantial development effort (Giering & Kaminski, 1998). Introducing a learned component can also require substantial changes to the coupling between numerical and physical routines. These challenges motivate models in which physical simulation and gradient optimization share the same implementation (Davenport et al., 2026).

Differentiable programming bridges this gap. When a simulation is implemented as differentiable numerical operations, automatic differentiation applies the chain rule through its computational graph to obtain derivatives of outputs with respect to inputs and parameters or learned weights (Baydin et al., 2018). This supports calibration, sensitivity analysis, inverse problems, and training over multiple

time steps. In particular, a neural parameterization can be optimized for its effect on a coupled trajectory, accounting for feedback between the learned tendency and the resolved dynamics (Rasp, 2020; Kashyap et al., 2024). NeuralGCM demonstrates the potential of this approach for Earth's weather and climate by coupling learned parameterizations to a differentiable atmospheric dynamical core (Kochkov et al., 2024). Its Dinosaur core solves the hydrostatic primitive equations using spectral horizontal coordinates and a sigma vertical coordinate. The JAX Circulation Model (JCM) uses the same core with an explicit suite of intermediate-complexity physical parameterizations, providing a setting for process-based studies and hybrid modeling for Earth's climate and atmosphere (Gelbrecht et al., 2023; Davenport et al., 2026). In a different physical domain, JAX MD illustrates how reusable numerical components and neural models can be composed within a common differentiable simulation environment (Schoenholz & Cubuk, 2020). AEGIS shares the Dinosaur backbone with JCM and NeuralGCM, but addresses a different physical regime and adds coupled state variables and operators needed for Mars.

In this work, we present AEGIS, a differentiable Mars Climate Model based on the Dinosaur Core. Unlike the Earth-focused configurations of its Dinosaur-based predecessors, AEGIS uses planetary radius, rotation, and thermodynamic scales at the dynamical interface. It adds Mars radiation and surface physics, evolves frost and regolith alongside the atmosphere, feeds CO_2 phase change into the dynamical core's prognostic log-pressure state, and projects the coupled atmosphere–frost inventory after each step. These additions make mass exchange and surface memory part of the differentiated trajectory.

We make three contributions:

1. We extend the Dinosaur core with planetary-unit rescaling, a coupled atmosphere–surface–frost–regolith state, CO_2 mass exchange into prognostic pressure, and an atmosphere–frost inventory projection.
2. We provide modular Mars radiation, surface, transport, and neural interfaces through which physical controls and learned components can enter the same coupled simulation.
3. We verify physical and neural gradients, evaluate Mars fields against reference products, and demonstrate calibration, year-long forward integration, and controlled neural experiments, and showcase them in Section 3.

2 METHODS

AEGIS couples a modular Mars physics suite to the Dinosaur spectral dynamical core, carrying the atmosphere, surface, CO_2 frost, and regolith in a single differentiable state (Section 2.1). Automatic differentiation through this coupled time step provides gradients for physical calibration and for training learned components that enter through a common neural interface (Section 2.2).

2.1 ATMOSPHERIC CONFIGURATIONS AND THE DYNAMICAL FRAMEWORK

AEGIS separates the description of a planetary atmosphere from the numerical machinery used to evolve it. A body configuration specifies radius a, gravity g, rotation rate Ω, specific gas constant R, heat capacity c_p, and reference temperature and pressure. We use Mars values for the planetary constants (NASA Goddard Space Flight Center, 2025). The dynamical equations consume these values through a common interface. Length and time are scaled by a and $(2\Omega)^{-1}$, respectively, so the nondimensional radius and rotation rate follow Dinosaur's conventions for each body, removing dependence on Earth-specific default scales while retaining the existing spectral operators.

The framework supports a hierarchy of atmospheric configurations. Dynamical configurations isolate resolved transport; physically forced configurations add radiation, surface exchange, mixing, and volatile reservoirs; hybrid configurations add a learned tendency or substitute a learned component at a selected physical interface. Complexity is therefore introduced through explicit components: a body definition establishes the mechanical and thermodynamic constants, while its radiation, condensables, and boundary conditions establish the atmospheric regime. The physically developed configuration evaluated here is Mars; extending to another atmosphere involves supplying the relevant process operators as well as its constants (Hu et al., 2025).

Dinosaur advances the hydrostatic primitive equations on a sphere (Kochkov et al., 2024; Davenport et al., 2026). The atmospheric state contains spectral vorticity and divergence, temperature relative to a reference profile, log surface pressure, optional transported tracers, and simulation time, on a vertical coordinate $\sigma = p/p_s$ that follows changes in atmospheric mass and the terrain-dependent lower boundary (Kochkov et al., 2024). Spectral transform models using sigma coordinates have a long history in Mars atmospheric modeling (Nayvelt et al., 1997). The main experiments use triangular truncation T21, a 64×32 longitude–latitude transform grid, and 12 equally spaced sigma layers.

The coupled state augments the atmospheric variables with surface temperature, pressure-equivalent CO_2 frost, and a vertical regolith temperature profile attached to their geographic columns. Writing the full state as $\boldsymbol{x}$, the coupled equations take the form

$$\frac{d\boldsymbol{x}}{dt} = \mathcal{D}_{\mathrm{imp}}(\boldsymbol{x}) + \mathcal{D}_{\mathrm{exp}}(\boldsymbol{x}) + \mathcal{P}_{\boldsymbol{\alpha}}(\boldsymbol{x}, t) + \mathcal{N}_{\boldsymbol{\theta}}(\boldsymbol{x}, t), \tag{1}$$

where $\mathcal{D}$ denotes resolved dynamics, $\mathcal{P}_{\boldsymbol{\alpha}}$ denotes physical parameterizations with parameters $\boldsymbol{\alpha}$, and $\mathcal{N}_{\boldsymbol{\theta}}$ denotes an optional neural tendency. A neural replacement for a physical component is composed inside $\mathcal{P}$ instead of being added a second time. Time integration uses Dinosaur's implicit–explicit SIL3 Runge–Kutta scheme (Whitaker & Kar, 2013), with linear gravity-wave terms treated implicitly and nonlinear dynamics and assembled physical tendencies on the explicit side. Planetary unit conversion, the augmented state, and the atmosphere–frost correction are implemented in AEGIS's surrounding operators.

The Mars operators in $\mathcal{P}_{\boldsymbol{\alpha}}$ span CO_2 and dust radiation, surface and regolith energy exchange, boundary-layer mixing and dry convection, and a CO_2 condensation cycle (Lacis & Oinas, 1991; Forget et al., 1999; Haberle et al., 2019; Petrosyan et al., 2011). The last is the distinctive Mars coupling: surface condensation and sublimation transfer mass between the atmosphere and frost, feeding the prognostic log surface pressure,

$$\left.\frac{d \log p_s}{dt}\right|_{\mathrm{phase}} = -\frac{1}{p_s}\frac{dI}{dt}, \tag{2}$$

Here, p_s is local surface pressure and $I = g m_{\mathrm{ice}}$ is the surface frost inventory in pressure units. Positive dI/dt denotes condensation onto the surface and lowers p_s; negative dI/dt denotes sublimation and raises p_s. The factor $1/p_s$ converts the pressure tendency to a log-pressure tendency. A per-step projection constrains the integrated atmosphere-plus-frost inventory to the specified mass budget. Its residual measures satisfaction of that constraint, not unprojected discretization error or full energy/momentum closure. Because CO_2 is both a radiative gas and a substantial mobile mass reservoir, carrying its phase change through the prognostic pressure state, alongside surface and subsurface temperatures (Kelly et al., 2006), makes this coupling part of the simulated system and its derivatives. The radiation, surface, boundary-layer, and CO_2-inventory operators are detailed in Appendix A.

2.2 Differentiation and neural components

Let $\Phi_{\boldsymbol{\alpha},\boldsymbol{\theta}}$ denote the operator that propagates a single time step, including the reservoir corrections. A trajectory is the repeated composition of $\boldsymbol{x}_{n+1} = \Phi_{\boldsymbol{\alpha},\boldsymbol{\theta}}(\boldsymbol{x}_n)$ while a loss function can be evaluated over an entire trajectory rollout as $\mathcal{L}(\boldsymbol{\alpha}, \boldsymbol{\theta}) = \ell\big(\mathcal{H}(\boldsymbol{x}_1), \boldsymbol{y}_1, \mathcal{H}(\boldsymbol{x}_2), \boldsymbol{y}_2, \ldots, \mathcal{H}(\boldsymbol{x}_N), \boldsymbol{y}_N\big)$, where $\mathcal{H}$ extracts evaluated fields and $\boldsymbol{y}_n$ denotes the target data at step n. JAX differentiates this discrete numerical trajectory using automatic differentiation. We use forward-mode differentiation for the three physical controls and reverse-mode differentiation for neural weights; forward mode efficiently evaluates derivatives with respect to a small number of controls, whereas reverse mode is appropriate for a scalar loss with many parameters (Baydin et al., 2018; JAX Developers, 2024a). Reverse-mode training through coupled rollouts, as in the heating experiment below, uses fixed-length scans with step rematerialization, which trades additional computation for reduced reverse-mode memory use by recomputing selected forward intermediates during the backward pass (JAX Developers, 2024b).

Learned components are exposed through two adapters that share one interface: a model is initialized from a random key into a JAX parameter tree, input normalization is fitted on training data only, and a `bind` call returns a pure JAX function that is passed to the coupled equations when the time step is




Figure 1: AEGIS architecture. Planetary constants and boundary data configure a shared GCM. Dinosaur supplies resolved dynamics; Mars physical operators supply column tendencies and surface evolution; neural components enter as bounded residuals or component replacements. The coupled time step advances atmosphere and surface reservoirs. Trajectory objectives connect the simulated fields to physical-parameter calibration and neural training through automatic differentiation. Dashed arrows denote this gradient-based update path.

built. The residual adapter adds a bounded heating tendency computed from current-state column features: the temperature profile, surface temperature and pressure, lowest-layer winds, geographic coordinates, and insolation, 21 features at T21/L12, as the equation

$$\dot{T}_{\mathrm{NN}} = h_{\max} \tanh f_{\boldsymbol{\theta}}(\boldsymbol{z}), \tag{3}$$

where $f_{\boldsymbol{\theta}}$ is the network, $\boldsymbol{z}$ the normalized column features, and $h_{\max}$ a fixed heating bound, converts it to the dynamical core's tendency units, and adds it to the physical tendencies while leaving the surface reservoirs under physical control; a zero-initialized output layer reproduces the physical model exactly at the start of training. The replacement adapter is a drop-in radiation component: it predicts interface fluxes from the column temperature and pressure profiles, dust, sunlight, albedo, and emissivity, bounds shortwave fluxes by the incoming sunlight, keeps longwave fluxes positive, imposes the top and surface boundary conditions exactly, and computes heating from flux convergence, so the column energy budget closes for any network weights. Both adapters accept a custom network through an `apply(params, features)` function and check the vertical grid and radiation configuration when bound. Because a bound component is part of the coupled step, its parameters can be trained on local targets or by reverse-mode differentiation through rollouts, and they are saved together with their normalization and metadata in validated checkpoints. Training protocols for the reported experiments are given in Appendix B.

3 Results

3.1 Verified gradients through the coupled Mars model

We verify derivatives through both a scalar atmospheric diagnostic and the objectives used for physical calibration and neural fitting. The physical derivative experiment differentiates final area- and layer-mean atmospheric temperature with respect to the three physical controls. It compares automatic differentiation (AD) with centered finite differences (FD) over six rollout horizons, from 1 to 148 steps, corresponding to 300–44,400 s at a 300 s timestep. For each control and horizon, FD estimates use perturbations 10^{-3}, 10^{-4}, and 10^{-5}, yielding 54 comparisons through the coupled dynamics and Mars physics given by Table 1. All 54 physical-sensitivity checks pass the recorded criterion. At the longest horizon, 44,400 s, the largest discrepancy is 3.56×10^{-7}. Agreement across perturbation sizes and integration horizons supports the use of the computed derivatives for physical-parameter sensitivity in these coupled simulations.

We extend this verification to multi-day rollouts and to several coupled diagnostics at once. Starting from a MY32, $L_s = 45^\circ$ ARCO-MACDA window, we compute the forward-mode Jacobian `jax.jacfwd` of global-mean surface temperature, surface pressure, and surface CO_2 frost with

Table 1: Derivative checks through coupled AEGIS trajectories. All comparisons satisfy $|g_{\mathrm{AD}} - g_{\mathrm{FD}}| \leq 10^{-8} + 10^{-4}|g_{\mathrm{FD}}|$.

Category	Objective	Horizon	Checks	Passed
Physical sensitivity	Final mean atmospheric temperature	1–148 steps	54	54/54
Physical calibration	Scalar fitting objective	task-dependent	9	9/9
Neural fitting	Multi-field trajectory objective	25 steps	3	3/3
Multi-diagnostic sensitivity	Mean surface T, p_s, and CO_2 frost	74–1480 steps	9	9/9

respect to the CO_2 longwave opacity multiplier β at its nominal value $\beta = 1$, through the complete coupled rollout at horizons of 0.25, 1, and 5 sols (74, 296, and 1480 steps). All nine derivatives agree with centered finite differences ($\epsilon = 10^{-3}$), with a maximum relative discrepancy of 3.99×10^{-6} (Fig. 2). The sensitivities are physically interpretable: increasing CO_2 longwave opacity warms the surface, with $\partial \bar{T}_s / \partial \beta$ growing from 0.488 to 0.785 and 0.945 K per unit β as the response accumulates, and the warmer surface suppresses condensation, so surface pressure rises while frost falls ($\partial \bar{p}_s / \partial \beta$ grows from 2.94×10^{-3} to 4.66×10^{-2} Pa). The pressure and frost derivatives are equal and opposite at every horizon, with their sum at most 9.3×10^{-11} Pa, so the gradient itself respects the atmosphere–frost mass inventory that the projection enforces on the state. A warm evaluation of the diagnostics and their Jacobian over the 5-sol rollout takes about 92 s on a single GPU, showing that exact sensitivities of the coupled mass exchange are available over multi-day horizons at modest cost.




Figure 2: Differentiable response of AEGIS to the CO_2 longwave opacity multiplier β at $\beta = 1$, from a MY32, $L_s = 45°$ window. Top: global-mean surface temperature, surface pressure, and surface CO_2 frost at rollout horizons of 0.25, 1, and 5 sols. Bottom: their derivatives with respect to β computed by forward-mode automatic differentiation through the full coupled rollout (lines) and by centered finite differences (crosses). The pressure and frost sensitivities are equal and opposite, reflecting conservation of the atmosphere–frost CO_2 inventory.

The historical twelve-window pilot reduces test temperature RMSE from 2.45505 to 2.39699 K, a 0.05805 K gain (Table 2). A paired-year percentile interval conditional on the two held-out Mars years is [0.03512, 0.08693] K; four windows and one fitted run do not establish population skill or optimizer uncertainty. Training loss was still decreasing over the last five of the 50 Adam updates, so convergence was not established. The validation-selected physical controls were (0.902924, 1.096341, 0.816336). These historical results require regeneration after the PBL and sidereal-clock fixes; the revised runner records bounded multi-start optimizer termination and projected gradients rather than equating a fixed update budget with convergence.

Table 2: Twelve-window reanalysis pilot: atmospheric temperature RMSE (K) at a single lead of 7397.937 s. Each split contains four seasonally distributed windows, combined as the square root of their mean squared errors. The physical baseline is shared by the two fitting experiments.

Configuration	Training	Validation	Test
Physical baseline	2.45	2.49	2.46
Calibrated physics	2.38	2.44	2.40
Coupled neural heating	2.42	2.49	2.46

3.2 Computational performance and long integrations

Published v1 reports an 86.6-minute one-year integration and an 11.4-hour ten-year integration at T21/L12, 450 s, float64. These are historical reports, not newly reproduced timings: the original GPU is identified only as `cuda:0`, and a complete public release of the headline fields, spinup, and timing inputs is pending. We withdraw the relative speed claim against NASA Ames because hardware, physics, grid, and output costs were not matched. The reproducibility guide specifies synchronized forward/gradient timings and separates compilation, warmup, and execution.

The available annual ablation archive instead uses 300 s, with a 150 s paired trajectory. Reanalysis of their 74 common output times gives 3.700 K equal-snapshot RMS temperature separation and 3.884 K under trapezoidal time weighting. These are transient sensitivity statistics, not climatological error estimates or a 450/300 s certification. Finite saved states do not establish physical realism or equilibration. Small post-projection inventory residuals must be reported separately from raw mass change and correction-induced energy and momentum changes. Differences between year-nine and year-ten global means must not be interchanged with spatial RMS differences; the headline metric definitions and raw fields remain to be reconciled.

3.3 Seasonal comparisons with MOLA topography

The MOLA-based lower boundary produces a strong geographic structure in surface pressure (Smith et al., 2001; NASA Planetary Data System, n.d.). Elevated regions contain less atmospheric column mass, while low-lying terrain supports higher surface pressure. Across our twelve season–reference comparisons, surface-pressure pattern correlations range from 0.975 to 0.989 (Table 3). The model reproduces the terrain-driven hydrostatic pressure structure set by the MOLA lower boundary across all four seasons and all three reference products. Figure 3 shows the terrain context and the seasonal fields at $L_s = 135^\circ$.

Surface-temperature correlations range from 0.732–0.821, but all twelve archived temperature biases are positive (+2.355 to +8.219 K). Frost RMSE ranges from 123.036 to 631.330 Pa-equivalent. Table 3 includes bias, RMSE, and correlation for all four fields, including frost and wind. High terrain-driven pressure correlation does not independently validate circulation or climatology.

The seasonal model uses opacity/exchange controls (0.25, 0.25, 1), selected after comparison with Ames; it is not an untuned or independent Ames validation. Calibration and neural baselines start at (1, 1, 1), while (0.6, 0.8, 1.4) is a synthetic teacher/gradient-test vector, not a fitted physical estimate. An independently calibrated, held-out seasonal evaluation remains required.

The seasonal comparison uses the documented sampling of each product: sparse model checkpoints within $\pm 5^\circ$ of the target season, five-sol Ames averages, daily ARCO-MACDA averages, and four-local-time MCD means. Wind levels and frost averaging periods also differ between products. We interpret these as model and reanalysis comparisons.

3.4 Physics ablations: process sensitivity

We experiment with an ablation suite that pairs the full Mars configuration with twelve process perturbations, such as removing boundary-layer mixing, dry convection, regolith conduction, or CO_2 exchange, CO_2 longwave opacity, and dust longwave opacity. Each is launched from the same initial state, so any divergence traces to the changed process, and we report the root-mean-square (RMS) separation from the paired full-physics trajectory. At a 44 400 s horizon, the perturbations separate




Figure 3: MOLA terrain on the comparison grid (top), and seasonal fields at $L_s = 135^\circ$ (bottom). The seasonal panels compare the physical model with NASA Ames, MCD 6.1, and ARCO-MACDA for surface temperature, pressure, near-surface wind speed, and surface CO_2 frost.

Table 3: Historical reference characterization, using the arithmetic mean of four seasonal statistics per row (not pooled spatial RMSE). Bias is model minus reference. Temporal sampling and wind heights are unmatched; opacity controls were Ames-informed.

Reference	Field	Bias	RMSE	Correlation
NASA Ames	Pressure (Pa)	-40.92	52.37	0.986
NASA Ames	Temperature (K)	+3.46	20.64	0.789
NASA Ames	Wind (m/s)	+9.75	11.91	0.442
NASA Ames	Frost (Pa-eq.)	+7.31	168.48	0.816
MCD 6.1	Pressure (Pa)	-41.56	57.18	0.977
MCD 6.1	Temperature (K)	+7.12	21.58	0.787
MCD 6.1	Wind (m/s)	+6.91	10.03	0.263
MCD 6.1	Frost (Pa-eq.)	-5.69	196.28	0.806
ARCO-MACDA	Pressure (Pa)	+3.43	40.72	0.988
ARCO-MACDA	Temperature (K)	+7.42	22.27	0.772
ARCO-MACDA	Wind (m/s)	+2.90	7.88	0.454
ARCO-MACDA	Frost (Pa-eq.)	-65.55	368.34	0.795

into a well-resolved sensitivity ranking (Fig. 4): regolith conduction and boundary-layer mixing dominate, followed by dry convection and the surface-exchange multipliers, while the calibrated radiative opacity multipliers produce smaller but clearly resolved changes. Halving the time step changes temperature by only 0.0042 K, a short-horizon numerical sensitivity check that does not certify long-horizon timestep convergence.

All thirteen archived configurations complete with finite sampled states. The full-physics year's maximum relative post-projection inventory residual is 4.28×10^{-9}, which verifies the inventory constraint only. The same archive contains a peak wind of 236.046 m s^{-1}, a minimum air temperature of 93.263 K, and a sampled maximum Mach number of 1.098. Air reaches 35.77 K below the model's layer-pressure CO_2 frost point. These diagnostics motivate timestep, drag, and saturation studies rather than establishing a single cause or physical validity.




Figure 4: Sensitivity of the T21/L12 Mars configuration. Bars show RMS differences from the paired full-physics trajectory over five saved times through 44 400 s; the right panel combines zonal and meridional wind RMSE by quadrature. All cases begin from the same state.

3.5 Neural components as coupled experiments

AEGIS exposes learned components through the same operator interface as its physical parameterizations. A neural module attaches at a chosen process either as a bounded additive residual or as a full component replacement. Because the module enters the coupled step, its effect propagates through the subsequent dynamics and physics, and it is trained by differentiating the resulting trajectory. We showcase two experiments using these interfaces exposed by the framework.

3.5.1 Learning radiative fluxes

We replace the compact physical radiation kernel with a structured learned flux operator that preserves the discrete radiative budget. Four-layer columns are first trained locally for 100 updates. Starting from that common fit, trajectory fine-tuning and local continuation each use 20 fixed updates (seed 0). This is an update-budget accuracy comparison, not equal computational cost. The v1 wall-time continuation happened to perform 234 updates, so its 0.416/0.979 K values are historical machine-dependent results and are not retained as the primary accuracy comparison. The new CPU float64 run and an independent repeat archive reports, histories, checkpoints, generated predictions, executed source hashes, and the actual update counts.

With fixed update counts, trajectory fine-tuning gives the lowest interpolation error (0.566 K), while continued local fitting gives the lowest higher-dust error (1.595 K). The local-fit baseline is 0.620/1.799 K, and continued local fitting is 0.592/1.595 K. Neither method wins on both splits. All variants complete the two-step T21 coupled execution check. The maximum held-out instantaneous radiation closure residual is 1.42×10^{-14} W m^{-2}; this component identity is not a full-GCM energy-conservation result. A repeated seed-0 run agrees within the recorded tolerance.

3.5.2 Recovering a physical parameter through a trajectory

The differentiable column system recovers a known 1.6 multiplier on a bulk sensible-heat conductance of $2\,\mathrm{W\,m^{-2}\,K^{-1}}$, using targets generated with fixed illumination, dust, and pressure per column. After 120 updates through 24-step trajectories, initial multipliers of 0.5, 1.0, and 3.0 converge to 1.59950, 1.59901, and 1.59696, all within 0.20% of the target (Fig. 6). Derivative checks at multipliers 0.5, 1.0, and 2.5 agree with finite differences to relative discrepancies below 3.41×10^{-10}. A derivative-free scalar reference reaches 1.60000007 in nine objective evaluations. This controlled recovery establishes parameter identifiability and a gradient-based inverse workflow through the shared differentiable column operators.

Table 4: Fixed-update, seed-0, 48-step column air-temperature RMSE (K). Both continuations start from the same 100-update local fit. Generated physical-kernel targets; not observational Mars skill.

Training	Interpolation	Higher dust
Initial local fit	0.620	1.799
Trajectory, 20 updates	0.566	1.653
Continued local, 20 updates	0.592	1.595




Figure 5: Fixed-update neural-radiation experiment at a 48-step column rollout (100 initial updates; 20 updates per continuation method; seed 0). Air-temperature RMSE is evaluated against the compact physical radiation kernel on held-out interpolation columns and higher-dust extrapolation columns (24 columns per split). The three bars compare the initial local fit, trajectory fine-tuning, and continued local fitting.




Figure 6: Physical inverse experiment from the neural-framework archive. Gradient-based optimization recovers a prescribed surface-exchange multiplier of 1.6 from three initial values over 120 updates. Each curve includes its initial value and the saved multiplier after every update. The parameter belongs to the controlled column model; the experiment tests the shared differentiable inference workflow.

4 Future Extensions

The present demonstrations span coarse-resolution physical simulations, coupled-heating seed, and structured neural component replacement, and they establish the differentiable physical and neural interfaces. The framework already supports the natural next steps: broader observational comparison, repeated neural experiments across seeds, and multi-year equilibration studies, each reusing the same coupled operators and trajectory objectives. The explicit planetary, radiation, and reservoir interfaces provide a route to experiments with changing Martian boundary conditions. Altered surface

albedo, imposed heating, or specified changes in volatile inventory could be introduced through these interfaces and evaluated with spatial climate diagnostics. Differentiable objectives could then connect intervention parameters to quantities such as regional temperature, frost persistence, or atmospheric pressure. For larger changes in atmospheric composition and mass, the body thermodynamics, opacity data, condensable processes, and applicable dynamical assumptions would be extended together.

5 Conclusion

AEGIS connects modular Mars operators to the Dinosaur dynamical core and differentiable physical and neural experiments. Archived gradient checks and controlled inverse experiments demonstrate useful discrete sensitivities. Historical reference comparisons expose temperature, wind, and frost errors and documented tuning; they do not establish independently validated climatology. The new fixed-update radiation comparison makes the learning budget reproducible and separates accuracy from machine-dependent timing. Converged reanalysis fitting, corrected-model seasonal and timestep studies, complete raw budget diagnostics, and a public release of the headline long-run fields remain necessary before stronger climate or performance claims.

Scope and reproducibility

The CO_2 cycle is surface-only: atmospheric condensation, latent heating aloft, and condensate sedimentation are absent. The frost relation uses the fixed-CO2/total-pressure approximation and is not a general gas-mixture calculation. Prescribed dust, equally spaced sigma levels, missing subgrid orographic drag, and numerical mass repairs limit long-run interpretation. A temperature clamp would not supply conservative missing physics. The revised implementation separates sidereal rotation (88642.663 s) from the solar sol (88775.244 s); historical nondimensional restarts must be reconstructed from dimensional fields before use with changed unit scales. Results from before and after these fixes must not be silently combined.

The repository archives the immutable arXiv v1 source, this revision, a claim-to-evidence map, the corrected radiation runs, and the recovered small-data calibration evidence under `paper/arxiv-2610.04081`. The simple command sequence in `docs/paper-reproduction.md` distinguishes executable small-data audits, fresh model runs, external inputs, and pending public fields. Historical figures retain their source revisions; only the fixed-update radiation figure is replaced by the new run. The revision does not assert that absent headline artifacts or new annual simulations have been reproduced.

References

Atilim Gunes Baydin, Barak A. Pearlmutter, Alexey Andreyevich Radul, and Jeffrey Mark Siskind. Automatic differentiation in machine learning: a survey. *Journal of Machine Learning Research*, 18(153):1–43, 2018. URL `http://jmlr.org/papers/v18/17-468.html`.

Ananyo Bhattacharya. ARCO-MACDA, 2026. URL `https://huggingface.co/datasets/ananyo01/ARCO-MACDA`. Dataset; accessed 23 September 2026; experiments pin archive revision 65a0bebd804b9c240752277e83f5737d58c6ee9c.

Ellen H. Davenport, J. Varan Madan, Rebecca Gjini, Jared Brzenski, Nick Ho, Tien-Yiao Hsu, Yueshan Liang, Zhixing Liu, Veeramakali Manivannan, Eric Pham, Rohith Vutukuru, Andrew I. L. Williams, Zhiqi Yang, Rose Yu, Nicholas J. Lutsko, Stephan Hoyer, and Duncan Watson-Parris. JCM v1.0: A Differentiable, Intermediate-Complexity Atmospheric Model. *EGUsphere*, 2026. doi: 10.5194/egusphere-2025-6266. URL `https://doi.org/10.5194/egusphere-2025-6266`. Preprint.

Francois Forget, Frederic Hourdin, Richard Fournier, Christophe Hourdin, Olivier Talagrand, Matthew Collins, Stephen R. Lewis, Peter L. Read, and Jean-Paul Huot. Improved general circulation models of the Martian atmosphere from the surface to above 80 km. *Journal of Geophysical Research: Planets*, 104(E10):24155–24175, 1999. doi: 10.1029/1999je001025. URL `https://doi.org/10.1029/1999je001025`.

Matthias Gelbrecht, Andrew White, Sebastian Bathiany, and Niklas Boers. Differentiable programming for earth system modeling. *Geoscientific Model Development*, 16(11):3123–3139, 2023. doi: 10.5194/gmd-16-3123-2023.

Ralf Giering and Thomas Kaminski. Recipes for adjoint code construction. *ACM Transactions on Mathematical Software*, 24(4):437–474, 1998. doi: 10.1145/293686.293695. URL https://doi.org/10.1145/293686.293695.

Robert M. Haberle, James B. Pollack, Jeffrey R. Barnes, Richard W. Zurek, Conway B. Leovy, James R. Murphy, Hilda Lee, and James Schaeffer. Mars atmospheric dynamics as simulated by the NASA Ames General Circulation Model: 1. The zonal-mean circulation. *Journal of Geophysical Research: Planets*, 98(E2):3093–3123, 1993. doi: 10.1029/92JE02946. URL https://doi.org/10.1029/92JE02946.

Robert M. Haberle, Melinda A. Kahre, Jeffery L. Hollingsworth, Franck Montmessin, R. John Wilson, Richard A. Urata, Amanda S. Brecht, Michael J. Wolff, Alexandre M. Kling, James R. Schaeffer, et al. Documentation of the NASA/Ames legacy Mars global climate model: Simulations of the present seasonal water cycle. *Icarus*, 333:130–164, 2019. doi: 10.1016/j.icarus.2019.03.026.

Zeyuan Hu, Akshay Subramaniam, Zhiming Kuang, Jerry Lin, Sungduk Yu, Walter M. Hannah, Noah D. Brenowitz, Josh Romero, and Michael S. Pritchard. Stable machine-learning parameterization of subgrid processes in a comprehensive atmospheric model learned from embedded convection-permitting simulations, 2025. URL https://arxiv.org/abs/2407.00124.

JAX Developers. Forward- and reverse-mode automatic differentiation in JAX. https://docs.jax.dev/en/latest/jacobian-vector-products.html, 2024a. Accessed: 2026-09-24.

JAX Developers. Gradient checkpointing with `jax.checkpoint` (`jax.remat`). https://docs.jax.dev/en/latest/gradient-checkpointing.html, 2024b. Accessed: 2026-09-24.

Melinda A. Kahre and Robert M. Haberle. Mars CO_2 cycle: Effects of airborne dust and polar cap ice emissivity. *Icarus*, 207(2):648–653, 2010. doi: 10.1016/j.icarus.2009.12.016.

Sameera S Kashyap, Raj Abhijit Dandekar, Rajat Dandekar, and Sreedath Panat. Modeling chaotic lorenz ode system using scientific machine learning, 2024. URL https://arxiv.org/abs/2410.06452.

N. J. Kelly, W. V. Boynton, K. Kerry, D. Hamara, D. Janes, R. C. Reedy, K. J. Kim, and R. M. Haberle. Seasonal polar carbon dioxide frost on mars: Co2 mass and columnar thickness distribution. *Journal of Geophysical Research: Planets*, 111(E3), 2006. doi: https://doi.org/10.1029/2006JE002678. URL https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2006JE002678.

Diederik P. Kingma and Jimmy Ba. Adam: A method for stochastic optimization. arXiv:1412.6980, 2014. URL https://arxiv.org/abs/1412.6980.

Dmitrii Kochkov, Janni Yuval, Ian Langmore, Peter Norgaard, Jamie Smith, Griffin Mooers, Milan Klower, James Lottes, Stephan Rasp, Peter Duben, Sam Hatfield, Peter Battaglia, Alvaro Sanchez-Gonzalez, Matthew Willson, Michael P. Brenner, and Stephan Hoyer. Neural general circulation models for weather and climate. *Nature*, 632(8027):1060–1066, 2024. doi: 10.1038/s41586-024-07744-y. URL https://doi.org/10.1038/s41586-024-07744-y.

Andrew A. Lacis and Valdar Oinas. A description of the correlated k distribution method for modeling nongray gaseous absorption, thermal emission, and multiple scattering in vertically inhomogeneous atmospheres. *Journal of Geophysical Research: Atmospheres*, 96(D5):9027–9063, 1991. doi: 10.1029/90jd01945. URL https://doi.org/10.1029/90jd01945.

Doug McNeall, Jonny Williams, Richard Betts, Ben Booth, Peter Challenor, Peter Good, and Andy Wiltshire. Correcting a bias in a climate model with an augmented emulator. *Geoscientific Model Development*, 13(5):2487–2509, 2020. doi: 10.5194/gmd-13-2487-2020.

NASA Goddard Space Flight Center. Mars fact sheet. https://nssdc.gsfc.nasa.gov/planetary/factsheet/marsfact.html, 2025. NASA National Space Science Data Center. Accessed 2026-09-23.

NASA Planetary Data System. Mars Global Surveyor Mars Orbiter Laser Altimeter: Mission Experiment Gridded Data Records. MOLA MEGDR archive, n.d. URL https://pds-geosciences.wustl.edu/mgs/mgs-m-mola-5-megdr-l3-v1/. Terrain asset used here: meg004/megt90n000cb.img; checksum recorded with the experiment.

Lev Nayvelt, Peter J. Gierasch, and Kerry H. Cook. Modeling and Observations of Martian Stationary Waves. *Journal of the Atmospheric Sciences*, 54(8):986–1013, 1997. doi: 10.1175/1520-0469(1997)054⟨0986:MAOOMS⟩2.0.CO;2.

Arakel Petrosyan, Boris Galperin, Søren E. Larsen, Stephen R. Lewis, A. Määttänen, et al. The martian atmospheric boundary layer. *Reviews of Geophysics*, 49(3), 2011. doi: 10.1029/2010RG000351.

Stephan Rasp. Coupled online learning as a way to tackle instabilities and biases in neural network parameterizations: general algorithms and Lorenz 96 case study (v1.0). *Geoscientific Model Development*, 13(5):2185–2196, 2020. doi: 10.5194/gmd-13-2185-2020. URL https://doi.org/10.5194/gmd-13-2185-2020.

Mark I. Richardson, Anthony D. Toigo, and Claire E. Newman. PlanetWRF: A general purpose, local to global numerical model for planetary atmospheric and climate dynamics. *Journal of Geophysical Research: Planets*, 112(E9), 2007. doi: 10.1029/2006JE002825. URL https://doi.org/10.1029/2006JE002825.

Samuel S. Schoenholz and Ekin D. Cubuk. JAX, M.D.: A framework for differentiable physics. In *Advances in Neural Information Processing Systems*, volume 33, 2020. URL https://arxiv.org/abs/1912.04232.

David E. Smith, Maria T. Zuber, Herbert V. Frey, James B. Garvin, James W. Head, Duane O. Muhleman, Gordon H. Pettengill, Roger J. Phillips, Sean C. Solomon, H. Jay Zwally, W. Bruce Banerdt, Thomas C. Duxbury, Matthew P. Golombek, Frank G. Lemoine, Gregory A. Neumann, David D. Rowlands, Oded Aharonson, Peter G. Ford, Anton B. Ivanov, Catherine L. Johnson, Patrick J. McGovern, James B. Abshire, Robert S. Afzal, and Xiaoli Sun. Mars Orbiter Laser Altimeter: Experiment summary after the first year of global mapping of Mars. *Journal of Geophysical Research: Planets*, 106(E10):23689–23722, 2001. doi: 10.1029/2000je001364. URL https://doi.org/10.1029/2000je001364.

Alex Valeanu, Kylash Rajendran, Luca Montabone, and Peter L. Read. Mars Analysis Correction Data Assimilation (MACDA): Atmospheric and surface fields produced from assimilated MGS/TES, ODY/THEMIS, and MRO/MCS observations, v2.0, 2026. URL https://catalogue.ceda.ac.uk/uuid/cd037a9ea387438fabf4d674dbe53088. Dataset; accessed 23 September 2026.

Jeffrey S. Whitaker and Sajal K. Kar. Implicit–Explicit Runge–Kutta Methods for Fast–Slow Wave Problems. *Monthly Weather Review*, 141(10):3426–3434, 2013. doi: 10.1175/mwr-d-13-00132.1. URL https://doi.org/10.1175/mwr-d-13-00132.1.

A MARS PHYSICS OPERATORS

MOLA elevation data provide the lower boundary (Smith et al., 2001; NASA Planetary Data System, n.d.). The raster is regridded onto the transform grid and converted into spectral orography. Prescribed dust fields can be supplied through seasonal or per-window adapters. These data remain separate from the numerical operators, making boundary-condition experiments straightforward.

Radiation combines CO_2 absorption and dust optical properties in a column flux calculation. The correlated-k approach represents nongray gaseous absorption using a reduced distribution of absorption coefficients (Lacis & Oinas, 1991). The main physical configuration uses bundled Ames-derived 12-band correlated-k tables, interpolated in temperature and log pressure, together with two-stream transfer. Solar and thermal fluxes are evaluated at layer interfaces, and atmospheric heating follows

from their convergence. For layer k, with pressure thickness $\Delta p_k = p_s \Delta \sigma_k$, the atmospheric heat capacity per unit area is

$$C_k = \frac{c_p \Delta p_k}{g}, \qquad \left.\frac{dT_k}{dt}\right|_{\mathrm{rad}} = \frac{Q_k}{C_k}, \tag{4}$$

where Q_k is radiative flux convergence. Computing atmospheric and surface heating from a common set of interface fluxes makes the column energy exchanges telescope to the top-of-atmosphere balance. This same construction is used for the neural radiation interface.

The surface temperature evolves from radiative, sensible, conductive, and latent heat exchange (Forget et al., 1999; Haberle et al., 2019). A bulk aerodynamic sensible heat flux couples the surface to the lowest atmospheric layer. The surface loss and atmospheric gain use the same flux, with atmospheric heating divided by the layer heat capacity. Surface stress transfers momentum, and Richardson-number-dependent vertical diffusion mixes momentum, heat, and tracers. The boundary-layer physics is motivated by Petrosyan et al. (2011); the implementation uses a batched tridiagonal linear solve. Dissipated resolved kinetic energy is returned as heat. Dry convective adjustment mixes unstable adjacent layers while conserving their sigma-weighted enthalpy. A 12-layer regolith conducts heat beneath the surface and supplies thermal memory across daily and seasonal forcing.

The CO_2 cycle couples temperature to atmospheric mass. Surface condensation releases latent heat and transfers mass from the atmosphere to frost; sublimation reverses the exchange. The local phase-change contributions satisfy

$$\left.\frac{dp_s}{dt}\right|_{\mathrm{phase}} = -\frac{dI}{dt}, \qquad \left.\frac{d\log p_s}{dt}\right|_{\mathrm{phase}} = -\frac{1}{p_s}\frac{dI}{dt}. \tag{5}$$

The frost temperature follows a pressure-dependent saturation relation, and an energy-limited option ties the exchange to the surface energy budget. After each complete time step, a projection repairs negative frost and transfers the corresponding deficit from the atmospheric column. A global pressure rescaling then matches the quadrature-integrated atmosphere-plus-frost inventory to its pre-step value, accounting for prescribed escape when enabled. This addresses the finite-step inventory error associated with advancing log pressure. Atmospheric pressure can still change seasonally as mass moves into and out of the surface reservoir.

B Calibration and neural-training protocol

Physical calibration and coupled heating are evaluated using twelve prespecified ARCO-MACDA reanalysis windows (Bhattacharya, 2026; Valeanu et al., 2026). Four windows from Mars years 24, 26, 29, and 31 are used for training; four from years 32–33 for validation; and four from years 34–35 for testing. Each split samples four seasonal quarters. Atmospheric initial conditions are constructed from the reference temperature, winds, and pressure. Surface and frost initialization use start-time fields, and the soil initialization uses preceding surface history. The native reference interval of 7397.937 s is integrated with 25 steps of 295.91748 s. Start-time dust and the orbital phase are matched for each window.

The training loss averages squared errors in temperature, zonal wind, and meridional wind, after scaling each field by its training-target standard deviation. It uses horizontal area weights, equal sigma-layer weights, and equal window weights. Feature normalization uses initial states from training data. Both fitting experiments perform 50 Adam updates (Kingma & Ba, 2014) with learning rate 10^{-3} and gradient-norm clipping at one. Validation selects the checkpoint, including the initial physical baseline among the candidates. Physical calibration adjusts CO_2 longwave opacity, dust longwave opacity, and surface exchange within bounds $[0.05, 2]$, $[0.05, 2]$, and $[0.25, 4]$. Neural fitting retains all three physical multipliers at one. Reported split RMSE is the square root of the mean per-window MSE, matching the figure-generation code.

C Additional seasonal reference comparisons

Figure 3 in the main text shows the seasonal fields at $L_s = 135°$. Figures 7 and 8 give the remaining aphelion-side and perihelion-side seasons, $L_s = 45°$ and $L_s = 315°$, under the same sampling conventions and the same set of reference products. The Ls225 transition-season panel is also included below; the table covers all four seasons and retains the historical sampling qualifications.




Figure 7: Seasonal fields at $L_s = 45°$. The panels compare the physical model with NASA Ames, MCD 6.1, and ARCO-MACDA for surface temperature, pressure, near-surface wind speed, and surface CO_2 frost, under the sampling conventions described in the main text.




Figure 8: Seasonal fields at $L_s = 315°$. The panels compare the physical model with NASA Ames, MCD 6.1, and ARCO-MACDA for surface temperature, pressure, near-surface wind speed, and surface CO_2 frost, under the sampling conventions described in the main text.




Figure 9: Seasonal diagnostic comparison at $L_s = 225°$, organized as a 4×4 grid. Each *row* is one surface field—surface temperature (K), surface pressure (Pa), near-surface wind speed ($\mathrm{m\,s^{-1}}$), and surface CO_2 frost (Pa-equivalent)—and each *column* is one source: the AEGIS physical model (leftmost) alongside the NASA Ames, MCD 6.1, and ARCO-MACDA reference products. All panels are longitude–latitude maps (0–360°E, 90°S–90°N) with the MOLA terrain contour overlaid for geographic reference, and the color scale is shared within each row so the model and the three references are directly comparable field by field. Reading across a row shows how closely the modeled field tracks each reference; reading down the leftmost column shows the modeled state across all four diagnostics. Table 3 summarizes the corresponding pattern correlations and RMSE.