

Dysfunctional Nucleus Tractus Solitarius: Its Crucial Role in Promoting Neuropathogenic Cascade of Alzheimer's Dementia—A Novel Hypothesis

Mak Adam Daulatzai

Received: 28 September 2011 / Revised: 16 November 2011 / Accepted: 15 December 2011 / Published online: 5 January 2012
© Springer Science+Business Media, LLC 2011

Abstract The pathophysiological mechanism(s) underlying Alzheimer's disease (AD) still remain unclear, and no disease-modifying or prophylactic therapies are currently available. Unraveling the fundamental neuropathogenesis of AD is an important challenge. Several studies on AD have suggested lesions in a number of CNS areas including the basal forebrain, hippocampus, entorhinal cortex, amygdala/insula, and the locus coeruleus. However, plausible unifying studies on the upstream factors that involve these heterogeneous regions and herald the onset of AD pathogenesis are not available. The current article presents a novel nucleus tractus solitarius (NTS) vector hypothesis that underpins several disparate biological mechanisms and neural circuits, and identifies relevant hallmarks of major presumptive causative factor(s) linked to the NTS, in older/aging individuals. Aging, obesity, infection, sleep apnea, smoking, neuropsychological states, and hypothermia—all activate inflammatory cytokines and oxidative stress. The synergistic impact of systemic proinflammatory mediators activates microglia and promotes neuroinflammation. Acutely, the innate immune response is protective defending against pathogens/toxins; however, when chronic, it causes neuroinflammation and neuronal dysfunction, particularly in brainstem and neocortex. The NTS in the brainstem is an essential multiple signaling hub, and an extremely important central integration site of baroreceptor, chemoreceptor, and a multitude of sensory afferents from gustatory, gastrointestinal, cardiac, pulmonary, and upper airway systems. Owing to persistent neuroinflammation, the dysfunctional

NTS exerts deleterious impact on nucleus ambiguus, dorsal motor nucleus of vagus, hypoglossal, parabrachial, locus coeruleus and many key nuclei in the brainstem, and the hippocampus, entorhinal cortex, prefrontal cortex, amygdala, insula, and basal forebrain in the neocortex. The neuronal and synaptic dysfunction emanating from the inflamed NTS may affect its interconnected pathways impacting almost the entire CNS—which is already primed by neuroinflammation, thus promoting cognitive and neuropsychiatric symptoms. The upstream factors discussed here may underpin the neuropathogenesis of AD. AD pathology is multifactorial; the current perspective underscores the value of attenuating disparate upstream factors—in conjunction with anticholinesterase, anti-inflammatory, immunosuppressive, and anti-oxidant pharmacotherapy. Amelioration of the NTS pathology may be of central importance in countering the neuropathological cascade of AD. The NTS, therefore, may be a potential target of novel therapeutic strategies.

Keywords Alzheimer's disease · Neuroinflammation · Nucleus tractus solitarius · Hypoglossal · Vagus nerve · Sleep apnea

Abbreviations

α 1-ACT	α 1-Antichymotrypsin
A β	Amyloid beta
Ach	Acetylcholine
BFB	Basal forebrain
BP	Blood pressure
AD	Alzheimer's disease
AHI	Apnea hypopnea index
AP	Area postrema
APR	Acute phase response
BFB	Basal forebrain

M. A. Daulatzai (✉)
Sleep Disorders Group, EEE Department, MSE,
University of Melbourne, ICT Building, 3rd Floor,
Room no. 344, 111 Barry Street, Parkville, VIC 3010, Australia
e-mail: makd@unimelb.edu.au

BMI	Basal metabolic index
BBB	Blood brain barrier
CBF	Cerebral blood flow
CCL2	Chemokine ligand 2
CD15	Immunological carbohydrate adhesion molecule
CIC	Circulating inflammatory cytokines,
c-fos	Proto-oncogene
CIM	Circulating inflammatory mediators
CNS	Central nervous system
COX-2	Cyclooxygenase-2
CRP	C-reactive protein
DMNV	Dorsal motor nucleus of the vagus nerve
DVC	Dorsal vagal complex
ERC	Entorhinal cortex
GMV	Gray matter volume
HIF-1	Hypoxia-inducible factor 1
ICAM 1	Intercellular adhesion molecule 1
IFN γ	Interferon gamma
IL	Interleukin
LPS	Lipopolysaccharide
MCI	Mild cognitive impairment
MIP-2	Macrophage inflammatory protein
NA	Nucleus ambiguus
NF- κ B	Nuclear factor of kappa light polypeptide gene enhancer in B-cells
NTS	Nucleus tractus solitarius
NO	Nitric oxide
NOS	Nitric oxide synthase
NK	Natural killer cell
OB	Olfactory bulb
OFC	Orbitofrontal cortex
ON	Olfactory nerve
OSA	Obstructive sleep apnea
PC	Piriform cortex
PFC	Prefrontal cortex
p-tau	Hyperphosphorylated tau
ROS	Reactive oxygen species
TGF- β	Transforming growth factor beta
THP-1	Human acute monocytic leukemia cell line
TNF	Tumor necrosis factor
VCAM-1	Vascular cell adhesion molecule-1
VNS	Vagus nerve stimulation

Introduction

Alzheimer's disease (AD) is an aging-associated disease affecting about 10% of those aged over 65, and over 45% of those aged 85. The two pathological hallmarks of AD are amyloid plaques composed of aggregated β -amyloid ($A\beta$) peptide, and neurofibrillary tangles composed of hyperphosphorylated tau protein [1]. There is a progressive

deterioration in cognitive and emotional functions in AD; unfortunately, as yet there is no definitive prophylaxis and/or curative treatment that stops and reverses the relentless neurodegenerative process of this disease. Progress in treatment strategies is hindered due to our poor knowledge of the fundamental upstream factors underlying the neuro-pathological cascade of this disease.

Aging is a biological process characterized by time-dependent, progressive, physiological declines and thus accompanied by increasing incidence of age-related diseases and attenuated CNS functions including those of sensory, motor, and cognitive. Aging has been suggested to be a state of chronic, low-grade molecular inflammation which links normal aging and the pathogenesis of age-related diseases [2–4]. Inflammation is considered pivotal in age-related physiological changes and the pathogenesis of many age-related diseases. Several studies implicate oxidative stress in the etiology of AD in man [2, 3, 5–10] and animals [8, 9]. Consequently, available data have established two facts: (a) aging-associated dysregulation of the immune system and (b) aging-associated alteration of redox status. Both processes inter-twine and exacerbate systemic inflammatory status, due to a wide variety of inflammatory mediators. Several studies have shown increased inflammation in old age [11–15]. Glial cells from old mice secrete more proinflammatory IL-6 and less of anti-inflammatory IL-10, compared to young adults [16]. An insidious close relationship exists between systemic infection \rightarrow neuroinflammation \rightarrow cognitive dysfunction in the aged [11, 17]. Stimulation of the peripheral innate immune system (e.g. with lipopolysaccharide, LPS) causes increased neuroinflammatory response in the brain of aged mice [18] and humans [16, 19]. Old mice [13] and old rats [20] infected with *Escherichia coli* possess increased hippocampal interleukin (IL)1 β and several other inflammatory cytokines, and undergo deficits in hippocampal-dependent memory, in comparison with similarly infected younger animals. This is because systemic circulating inflammatory cytokines (CIC) have been shown to impair synaptic function/plasticity [21, 22] and decrease gray matter volume [23, 24]. Consequently, an increase in neuroinflammatory response may be correlated with susceptibility to cognitive impairment [25]. Further, there is strong clinical evidence that AD is accompanied by an inflammatory response involving peripheral cytokines including IL-6, TNF- α , IL-1 β , IL-12 and IL-18 [14].

The nucleus of the solitary tract (NTS) is an important brainstem nucleus where “initial integration” occurs of a wide variety of heterogeneous afferent inputs including gustatory, gastrointestinal, cardiovascular, and respiratory, to name a few. The brainstem circuits control several cardinal functions encompassing all above mentioned afferents, sympathetic and parasympathetic systems, neurotransmitter,

and hormonal systems; all these involve the crucial NTS [26–31]. The neuroinflammatory response in the CNS impacts the neocortex and the brainstem, and primes its different regions. Although the importance of neuroinflammation in AD is well recognized [14, 16, 19, 32, 33], however, as yet there is no unifying hypothesis on the initial upstream factors that provoke and exacerbate the fundamental neuropathogenetic steps of AD. The current thesis is the first to draw attention and emphasize that in the wake of CNS neuroinflammation in aging—it is the degeneration and neuronal apoptosis particularly within the dysfunctional NTS—that disrupts crucial CNS pathways. Since the NTS is reciprocally interconnected with almost the entire CNS, the dysfunctional NTS exerts deleterious impact on various essential neurophysiological functions in the vulnerable aged. Thus, mediated by neuroinflammation, the NTS is posited to occupy a central position in promoting the early upstream factors of neuropathogenetic cascade—that may cause cognitive decline in AD.

Evidence for Systemic Proinflammatory Cytokines/Inflammation in the Elderly

Despite the normal immune surveillance and blood brain barrier (BBB) regulation, toxic substances, bacteria, and virus can enter the CNS, e.g. via transcytosis through endothelial cells-inducing cytokine and chemokine synthesis and causing neuronal dysfunction. The risk factors that increase proinflammatory cytokines and inflammation in the elderly include: (1) aging, (2) obesity, (3) infection, (4) cigarette smoking, (5) psychosocial Stress, (6) hypothermia, and (7) Obstructive sleep apnea, to name a few. The inflammatory insults can be acute, prolonged, repeated, and chronically synergistic. Further, the following evidence suggests that various stigmata including aging are associated with multiple cascades that arise from separate pathophysiological processes, each associated with dysfunctional cognitive sequelae.

Aging and Inflammation

During normal aging, dysregulated immune and inflammatory responses occur in both humans and animals. The NF- κ B transcription factor and NF- κ B-dependent gene transcription are activated in systemic inflammatory processes upregulating the transcription of the classical proinflammatory cytokines [2, 34–36], as well as other proinflammatory proteins such as adhesion molecules viz. VCAM-1 (P-, E-selectin) and ICAM-1 during aging [37]. The neurons of aged animals show greater vulnerability to CIC/neurotoxins than those of younger animals, thus reflecting a correlation between age-dependent

inflammatory responses and an increased risk of neurodegeneration [38, 39]. The middle-aged (compared with young) individuals show increased sensitivity to acute LPS administration, upregulated neuronal c-fos immunoreactivity, and a lower tolerance development; they mount enhanced immune responses even to a lower dose of endotoxin [38, 40]. There is copious evidence of occurrence of high cholesterol, atherosclerosis, and inflammation in both aging and AD [41]. Noradrenaline is a well known anti-inflammatory molecule. The findings of selective decrease in noradrenergic innervations of the hippocampal dentate gyrus (DG) and the frontal cortex (FC) [42] and neurodegeneration of LC neurons in the aging brains of amyloid precursor protein [APP(swe)] and presenilin-1 [PS1(DeltaE9)] mice mimic the neuropathology noted in the AD brains [43].

Obesity and Inflammation

There is substantial evidence that adipocytes are involved in inflammation [44] and obesity leads to reactive astrocytes, increased synthesis of adipokines [45], caspase-1 [46], proinflammatory cytokines such as TNF- α , IL-1 β , IL-6, IL-8, IL-18, E-selectin, and a decreased synthesis of antiinflammatory cytokine IL-10 [46–48]. Obesity is associated with chronic low-grade systemic inflammation during mid- and late-life and increases the risk for neurodegenerative diseases including AD [49–51]. Higher BMI was associated with decreased gray matter volume [52–54]. In the Framingham Heart study, the relationship between obesity and cognitive dysfunction has been shown (Wolf et al. [55]). Several studies have described an association between atherosclerosis [56], high cholesterol, diabetes mellitus, obesity in midlife [57–61] and the incidence of AD.

Infection and Inflammation

The nose is said to be the air conditioner of the respiratory system; however, invading allergens/viruses/bacteria impact the airways negatively, and promote proinflammatory cytokine synthesis both locally and systemically. The most common infection in humans is viral respiratory tract infection (common cold) caused mainly by rhinoviruses. In acute rhinitis and influenza, viruses enter the nose/eye, initiate infection and trigger the release of inflammatory mediators, evidenced by the presence of TNF- α , IL-8, and MIP-2 in nasal secretions [62, 63]. The epithelial damage [64] may lead to secondary bacterial infection, local inflammation, microvascular leakage, degradation of extracellular matrix components, and activation of the innate immune system (to combat the bacterial invasion) [64–66]. An important source of circulating LPS and

proinflammatory cytokines in the aged is excess of gut microbiota and gut inflammation [67–69]. In addition, about 40% of persons over 60 years of age are known to have *Helicobacter pylori*, which causes peptic ulcers by damaging the mucous coating of the stomach and duodenum. Increased cerebrospinal fluid *H. pylori* antibody has been documented in Alzheimer's disease [70]. Many elderly take anti-inflammatory medication which is known to erode the lining of the gut. Both infection and inflammation are sources of neuroinflammation including that found in the NTS; see the section below on “Invading Pathogens Cause NTS Dysfunction”. Even a single intraperitoneal injection of *E. coli* in the aged rats leads to specific deficits in long-term memory and long-lasting synaptic plasticity in hippocampus—this being strongly dependent on brain-derived neurotrophic factor (BDNF) [71].

Smoking and Inflammation

There is a positive association between tobacco smoking and increases in pro-inflammatory markers. A British study found higher CRP and fibrinogen levels in smokers relative to non-smokers [72]. Importantly, compared with never smokers, current cigarette smokers showed significantly higher levels of white cell count, hematocrit, blood and plasma viscosity, tissue plasminogen activator antigen, and fibrin D-dimer [72]. Another longitudinal study from Uppsala also found higher IL-6 levels in both current smokers and former smokers, compared to non-smokers [73]. A recent study has documented that chronic smoking is associated with inferior performance on measures of general intelligence, learning and memory [74].

Psychosocial Stress and Inflammation

Psychosocial stress alters innate immune function and aggravates inflammation. In mice, exposure to repeated social stress causes airway inflammation, altered corticosterone responsiveness [75], and augmented effects of LPS [76]. Further, in the absence of peripheral immune challenge, social disruption alone induced pulmonary inflammation and increased myeloperoxidase activity [77]. Remarkably, potentiation of CNS proinflammatory cytokine occurs in stressed animal [78]. Persons undergoing emotional stress of bereavement have an increase in IL-1 β and TNF receptor II in orbitofrontal cortex (OFC) [79]. Acute stress elicits increased natural killer (NK) cell counts in PFC, and increased systemic levels of IL-1 β and IL-6. Several data have reported that the stressors upregulate IL-6 gene expression in various brain regions, increase IL-1 β and IL-6 into the circulation, and cause proinflammatory activation in neurons [78–80]. Further, various psychiatric

stressors exert powerful impact on brainstem causing alteration in glutamate signaling genes SLC1A2, SLC1A3 and GLUL [81]. Interestingly, genome-wide transcriptional profiling conducted on healthy adults (25–40 year old) who grew up in low socioeconomic status early in life, showed activation of markers of systemic inflammation including IL-6, and immune activating transcription factor AP-1 activity [82]. In drug-free persons with major depression, gene set analysis showed up-regulation of a variety of proinflammatory cytokines, including interleukin IL-1 α , IL-2, IL-3, IL-5, IL-8, IL-9, IL-10, IL-12A, IL-13, IL-15, IL-18, interferon gamma (IFN γ), and lymphotoxin α . This study found that the PFC showed evidence of local inflammation, oxidative stress, and apoptosis [83]. The age-related increases in IL-1 β mRNA and protein are correlated with increased microglial stimulation, and this priming to an inflammatory phenotype may underpin enhanced neuroinflammation noted in aging [18], other age-related conditions, and psychosocial stress with potentially serious neuroinflammatory consequences. The above mentioned may also mediate the early neuropsychiatric and behavioral alterations noted in AD. Further, psychological stress has an inverse association with cognition [84].

Hypothermia and Inflammation

Hypothermia is a thermoregulatory response to systemic inflammation and metabolic deterioration in patients and experimental animals alike. Systemic infections/inflammation may alter the body temperature, and this may modulate the host response to infection. Hypothermic septic patients have a significantly worse outcome than the pyrexia or normothermic. Peri-operative hypothermia is common owing to anesthetic action, surgical procedures, and the drugs. The mechanisms regulating hypothermia are not fully understood, but NF-kappaB-dependent gene expression and cytokines such as TNF- α , IL-6, and IFN γ may be involved [85]. TNF- α functions as an endogenous cryogen inducing hypothermia and causes changes in cytokine generation comparable to those stimulated by zymosan, toxic shock syndrome toxin-1, or LPS. Further, hypothermia exerts stimulatory effect on TNF- α and IL-1 β generation in primary monocyte cultures and the human THP-1 monocyte cell line, and that hypothermia causes changes in TNF- α and IL-1 β mRNA accumulation. A threefold increase has been shown in reporter gene studies of the human TNF- α promoter at 32°C versus 37°C [85]. Flow cytometric analysis and simultaneous measurement of IL-1 β , IL-6, IL-8, IL-10, IL-12p70, and TNF- α in monocyte culture of healthy humans showed higher levels in supernatant under hypothermic conditions [86]. Furthermore, mild hypothermia affects the balance of pro- and anti-inflammatory cytokines leading to a pro-inflammatory

state [87]. There is an age-associated increase in cytokines in hypothermics; proinflammatory IL-1 β and IL-6, were higher in blood in aged endotoxemic mice (vs. the young) [88]. Hypothermia is a critical factor that causes deleterious impact on brain stem and neocortical functions. Additionally, hypothermia in the elderly promotes not only sensory and motor changes, but memory impairment as well [89].

Obstructive Sleep Apnea (OSA) and Inflammation

Snoring and OSA are caused by incomplete obstruction of the upper airway (UA) and up to 40% of the general population may suffer from these disordered breathing conditions. Snoring is associated with cytokine release from blood cells, and promotes inflammation [90]. Snoring involves vibration of the soft palate, pharyngeal walls, epiglottis and the tongue, and causes lesions of the UA mucosa, pharyngeal muscles, and their innervating nerves [91–93]. Therefore UA mucosa in snorers is edematous and inflamed. The levels of proinflammatory cytokines TNF- α and IL-6 are elevated in the uvula of nonapneic snorers, but are much higher in OSA [94, 95]. Mechanical vibration simulating snoring, triggered an inflammatory cascade in human bronchial epithelial cells, reflected by increased in IL-8 release [96]. Finally, higher BMI, alcohol consumption, and cigarette smoking exacerbate inflammation in snorers [97, 98]. Acute partial sleep deprivation as in OSA has been shown to result in weight gain/obesity [99] and increased risk of oxidative stress in OSA [100, 101].

In OSA, there is evidence of chronic systemic inflammation, evidenced by elevated levels of plasma CRP [102], soluble adhesion molecules [2], and leukocyte superoxide [103]. Further, there is an increase in levels of circulating ICAM-1, VCAM-1, and L-selectin in OSA [104]. Increased expression of adhesion molecules CD15 and CD11c from monocytes in OSA patients has been implicated in adverse effect on vascular proinflammatory/anti-inflammatory homeostasis [103]. The above has been confirmed by increased production of proinflammatory IL-4 cytokine, and a decreased production of IL-10 (anti-inflammatory cytokine), in patients with moderate to severe OSA (AHI of >10/h) [105]. An increase in lipid peroxidation and generation of reactive oxygen species (ROS) has been documented in OSA patients [106]. The ROS increase occurs in repetitive episodes of hypoxia in experimental animals also [107, 108]; the reoxygenation/reperfusion phase in OSA is said to be the culprit that promotes ROS production, oxidative stress, and neuronal apoptosis [108]. OSA has been shown to correlate with hypertension, which has been implicated in AD [109].

OSA is associated with unique cerebral alterations that may explain the behavioral and neurocognitive alterations

observed. The vascular pathologies associated with OSA lead to the impairment of optimal cerebral perfusion [110] which has been emphasized to be a critical factor in the development of cognitive dysfunction [111]. Several studies have documented that hypoxia due to OSA causes neuropathological changes and cognitive/memory decline [112, 113]. The latter may result due to decreased oxidative metabolism in the brain, and impairment of neurotransmission. Different techniques including transcranial Doppler, event-related potentials, MR spectroscopy, and structural and functional MRI have clearly demonstrated changes in blood flow, metabolism, morphology, and activation in neurocognition-related brain regions in the OSA patients [114–139]. Decreased cerebral activation during the working memory task in OSA patients reflects that these individuals possess impaired cerebral responses during executive function [140–142].

OSA: Endothelial Dysfunction/Inflammation

Normal endothelium plays an important role in regulating vasomotor tone and maintaining inflammatory and coagulation homeostasis. However, these functions are altered in OSA patients [143–145]. OSA is characterized by vascular inflammation noted by upregulation of cyclooxygenase-2 (COX-2) and inducible NOS (iNOS) in endothelial cells [144]. Upregulation of COX-2 in OSA may increase superoxide production resulting in oxidative stress, increased platelet activation, and endothelial dysfunction/vasoconstriction [146]. In OSA, aggregation and adhesion of circulating leukocytes to the vascular endothelium causes blood vessel inflammation, cause endothelial cell apoptosis with impaired endothelial repair [105–107, 326]. The reasons for endothelial dysfunction in OSA are repetitive hypoxia/reoxygenation during apneas and hypopneas and oxidant-related microcirculatory endothelial dysfunction [144, 145]. There is reduced nitric oxide (NO) availability in OSA patients [143]; importantly, the decreased eNOS activity and increased nitrotyrosine production (byproduct of NO degradation) in endothelial cells provide direct evidence that bioavailability of NO is reduced in OSA patients [144]. This reduced NO availability in OSA impacts endothelial function thus enhancing vulnerability for disorders [143–146]. Even in healthy subjects, sleep deprivation causes a 50% decline in vasodilation reflecting reduced endothelium-dependent NO availability [147], increase in Proinflammatory cytokines IL-6, TNF- α , CRP, platelet adhesion thus promoting coagulation cascade [148–150]. NO is swiftly scavenged by ROS, producing the toxic metabolite—peroxynitrate. Thus as expected, nitrotyrosine expression is greater (than controls) in endothelial cells depicting the presence of enhanced endothelial oxidative stress [144, 145]. The endothelial oxidative damage and

ROS production with a decrease in NO perpetuate a cyclical pattern of endothelial injury [151, 152]. Hence, in OSA patients (those having AHI of >10) repetitive apnea/non-apnea episodes promote inflammation, adversely impact endothelial function, upregulate cell death receptors and mitochondria-dependent apoptotic pathways, that culminate in endothelial apoptosis [153].

Nucleus of the Solitary Tract (NTS)

The NTS is a compact network of neurons; it has copious afferent and efferent pathways that affect central homeostatic control. An array of extensive connections indicates that the NTS is a key structure for sensory, autonomic and neuroendocrine functions. CNS areas that receive a direct projection from the NTS project back to this nucleus. This nucleus contains an enormous range of neuroactive substances; indeed, most of those identified within the CNS are also found in the NTS, as neurotransmitters and neuromodulators [154]. The NTS undoubtedly is a nucleus of tremendous importance subserving extensive key interaction with almost every system of the body. Copious and well documented evidence is available on this brainstem hub nucleus and its influence/impact throughout the CNS. The NTS, the area postrema (AP), and the dorsal motor nucleus of the vagus (DMNV) nerve are components of the dorsal vagal complex (DVC).

The NTS is located in the dorsal brainstem and is the primary site for termination and integration of sensory afferents, such as baroreceptor, chemoreceptor, nociceptors, and afferents from several key body systems, and from UA and tongue. The visceral afferents—i.e. abdominal sensory receptors project via the vagus (and glossopharyngeal) nerve to the DVC. The sensory afferents form synapses with one or more neurons in the NTS in a loose viscerotopic organization. Rostral NTS receives sensory inputs from the UA including the soft palate, epiglottis, pharynx, and the tongue. In intermediate NTS adjacent to area postrema and close to the midline, gastrointestinal inputs synapse, whereas respiratory tract and pulmonary sensory afferents synapse in lateral and ventrolateral regions of the intermediate NTS. Cardiovascular afferents synapse in the medial NTS [155]. In the most caudal NTS, however, cardiovascular, respiratory/pulmonary, and gastrointestinal afferents synapse mediolaterally as well [30, 156, 157]. Overlap of these heterogeneous sensory inputs increases in the caudal NTS where there is increased sensory convergence. Thus the NTS is the first CNS region for synaptic contact of the above afferents. The signal processing of sensory information from the heart, lungs, gut, UA, and the tongue at these synapses determines the output to all downstream NTS synapses in the reflex pathways.

The second-order NTS neurons, therefore, integrate the sensory information including the vagal afferent inputs, spatially and temporally, orchestrate an efferent output, and transmit it to various interconnected foci—including the hypoglossal nucleus and the parasympathetic preganglionic neurons of the DMNV [158].

Various circulating inflammatory mediators (CIM), may directly or indirectly gain access and influence the activity of NTS neurons. Adjacent to NTS is the AP; it lacks a BBB and its prominent axons project to the NTS [158]. The AP provides an anatomical pathway whereby CIM can affect the NTS neurons. The caudomedial NTS also lacks a competent BBB, it possesses complexes of fenestrated capillaries and perivascular spaces which provide the NTS neurons direct exposure to blood-borne CIM. The brain is an immunologically competent organ and expresses constitutively a number of chemokines and chemokine receptors in neurons, astrocytes, microglia, and oligodendrocytes [159, 160]. Peripheral LPS challenge causes a hyperactive microglial response in the aged brain associated with higher induction of inflammatory IL-1 β and a significant increase in IL-1 β mRNA expression in the brain of aged mice [36]. Microglia produce IFN- γ during acute infection; they produce IFN- γ mRNA and MHC class II mRNA expression following stimulation with IL-12 and/or IL-18 [161]. There is evidence that inflammatory mediators can influence the brainstem neuronal function directly and the NTS itself is a primary CNS detector of cytokines [162]. Indeed CIM can influence NTS neuronal function directly through local synthesis of inflammatory mediators [163]. Thus binding of CIM, e.g. IL-1 β to its receptors on the neuronal membrane, including those of the NTS, initiates signaling cascades upregulating transcription of genes including those of COX-2, TNF- α , and IL-6, and other mediators; these then recruit leukocytes and macrophages that release additional inflammatory cytokines [36, 159–164].

It is important to underscore that both afferent and efferent parasympathetic activity plays a crucial role in immunomodulation [165–167]. By having a “wandering” route through the body, the vagus (Xth cranial nerve) is uniquely equipped to provide an effective early warning system about the pathogenic presence, and feedback to the immune system when pathogens are cleared [165]. The majority of vagal fibers are sensory providing sensory information from a wide range of body systems; this includes relaying the detection/presence of pathogenic invaders [165–167]. It is in the NTS that the afferent and efferent components of the parasympathetic nervous system meet [168]. The NTS having neural-immune communication mounts a specific and effective response [165, 168]; through vagus it affects a broad array of physiological processes [169]. The NTS provides input to the DMNV, nucleus ambiguus (NA), hypoglossal and other

brainstem nuclei; these nuclei provide extensive efferent signals [170]. The efferent parasympathetic pathways constitute the “cholinergic anti-inflammatory pathway” [166, 171, 172]. Ascending from the NTS, the vagus reaches the parabrachial nucleus, thalamus, paraventricular nucleus, amygdala, hippocampus, insula, anterior cingulate cortex, and the PFC [173]. Chronic systemic/neuroinflammation may cause synaptic transmission perturbation and attenuate multitude of efferent signaling pathways from the dysfunctional NTS [170, 174–179]. A dysfunctional NTS would be deleterious to a multitude of CNS foci and body systems that have reciprocal projection to this nucleus. Taken together, the abovementioned studies suggest that the dysfunctional NTS may be responsible for promoting neuropsychiatric and cognitive disturbances noted in the elderly and in AD.

Disparate Afferent(s) and the NTS-Mediated Responses

Maintaining cardiorespiratory fitness (CRF) mitigates structural and functional brain changes related with aging, and preserves cognitive function [180]. Thus, higher CRF preserves brain volume thereby minimizing brain atrophy. In transgenic mice, exercise decreased neuropathology in both cortical and hippocampal regions decreasing amyloid load [181]. The relationship between lower CRF and brain atrophy is present in early AD [182]. This can be explained as follows: (1) CRF has the propensity to attenuate AD pathology and brain atrophy; (2) AD pathology inherently modifies/downregulates CRF; and (3) there is a possible underlying factor such as the dysfunctional NTS that downregulates CRF and enhances neuropathologic burden in AD. It is plausible that based on its functional status, the NTS may influence the CRF and related phenomena [183–186] (see below).

Owing to its immense importance, baroreceptor example is used here as a prototype. Delineation of multitude of afferent receptors vis-à-vis NTS is beyond the scope of this paper. However, some afferents and the involvement of the NTS have been described elsewhere in the above sections.

A number of epidemiological studies have shown a strong association between AD and cardiovascular risk factors, particularly hypertension. Hypertension increases with advancing age. Hypertension is a cardiovascular risk factor [187] which is associated with AD [59, 60, 188]. The densities of SP and NFT in hypertensive subjects are elevated compared to controls [189, 190]. The pathophysiological relationship between hypertension and cognition is important, in that, hypoperfusion and neurodegeneration are expected to be possible underlying mechanisms [191, 192]. Hypertension is also associated with disruptions in neurovascular coupling, decreased perfusion, vascular reserve capacity, microvascular disease, stroke, cognitive

decline, and dementia. The baroreceptor reflex is a negative feedback system; baroreceptor stimulation decreases hypertension [193–200]. Conversely, NTS lesions are known to interrupt central baroreceptor pathways [196]. Artificial stimulation of baroreceptor afferents increases parasympathetic outflow but decreases sympathetic outflow to the cardiovascular system [193–200]. Indeed, age-associated changes in baroreflex regulation may involve NTS catecholaminergic mechanisms [201, 202] suggesting that any dysfunctional NTS physiology may contribute to altered regulation of parasympathetic discharge [203].

Cardiovascular system is affected by oxidative stress (ROS) and expression of subclinical proinflammatory TNF- α , IL-6, and CRP [204]. Increased sympathetic activation is a risk factor that potentiates cardiac hypertrophy, fibrosis, and the renin-angiotensin system [205–207]. However, enhancing vagal tone, via vagal nerve stimulation (VNS), exerts anti-sympathetic action through well characterized mechanisms [208–210]. Interestingly, in addition to anti-sympathetic effects, the vagal activation (a cholinergic effect) also mediates anti-inflammatory actions. In LPS-stimulated human macrophage culture, proinflammatory cytokines such as TNF- α , IL-1 β , IL-6, and IL-18 were suppressed by ACh [211]. In LPS-induced endotoxemic shock, bilateral VNS ameliorated hypertension and prevented surging of plasma TNF- α [211]. Equally important, VNS mediated ACh release activates the endothelial dependent NO synthesis—a protective mechanism against hypertension and atherosclerosis [212]. Furthermore, a 3-month VNS stimulation improves Left Ventricular function and hemodynamics, together with normalization of TNF- α , IL-6 [213], and NO (mRNA and protein expression) [214]. The above mentioned suggests the possible in vivo importance of integrity of the NTS.

The NTS is extensively interconnected; this network may play an important role in several regulations/functions, in addition to cardiovascular system [195, 215]. Another example of the ubiquitous nature of the NTS is given below. The inflammatory signal induced by LPS reaches the pre-optic anterior hypothalamic area (POA). Subdiaphragmatic vagotomy or abdominal perivagal lidocaine administration, or lidocaine injection into the NTS prevented the LPS action. These data suggest that LPS function is mediated by the vagus nerve which conveys the signal to the NTS that in turn stimulates norepinephrine release within the POA [216]. VNS stimulates both vagal afferents and efferents [217], and neural activation of NTS enhancing the brain-derived anti-inflammatory response [216, 218, 219]. The vagal nerve, therefore, is an important modulator of the immune system [172, 220]; it controls inflammation and modulates the immune response through a ‘nicotinic anti-inflammatory pathway’ dependent on the α 7-nicotinic ACh receptor [221]. Vagus nerve commences from the NTS, and

supportive evidence of cognitive effects of VNS has been well documented [221–225].

Finally, the combination of advanced age in conjunction with other stigmata described above, and the dysfunctional NTS may interact in a complex pattern and provide pathophysiological impetus that leads to cognitive decline and dementia. The latter is promoted due to interactions between cortico-cortical and subcortical disconnection (synaptic dysfunction), neuroinflammation, altered cholinergic/NTS transmission, autonomic disturbance, dysfunctional vascular endothelium, altered regulation of blood flow, deposition of A β and NFT, and neuronal degeneration/death in several key brain areas. These attenuate decision-making, cognition and memory, thus promoting AD.

The NTS Vis-à-Vis Deficits in Known Key Brain Regions in AD

AD is multifactorial and heterogeneous and may involve more than one etiopathogenetic mechanisms. One of the major hallmarks of AD is the degeneration of BFB cholinergic neurons indicating an important role of the cholinergic system in learning and memory. Studies have emphasized the critical role of pro-inflammatory cytokines as common mediators of cholinergic neuronal damage [226]. In contrast to late AD, patients with mild cognitive impairment (MCI), apparently show no cholinergic neurodegeneration. There is abundant evidence that A β may trigger cholinergic dysfunction through action on $\alpha 7$ nicotinic acetylcholine receptors, affecting NGF signaling in cholinergic neurons [227, 228]. Mesulam concluded that “cholinergic loss is neither a primary pathogenetic factor of AD nor the principal correlate of its clinical manifestations” [229]. However, as the cholinergic dysfunction progresses, it does influence cognitive deficits. These observations are consistent with the clinical evidence that cholinergic therapies (use of cholinesterase inhibitors) alone are unable to treat AD.

A major feature of AD is the degeneration of noradrenergic locus coeruleus (LC) neurons [230–232]. LC is the sole source of noradrenaline (NA) to neocortex and the hippocampus [233]. Attention, memory, arousal, vigilance and mood are affected by NA signaling [233]. LC damage has been shown to upregulate gliosis and inflammation [234–237], excitotoxicity [238], oxidative stress [238, 239], and A β deposition in the hippocampus and neocortex [238]. LC degeneration has been shown to reduce NGF and/or BDNF in LC projection sites in AD [231]. Although NA agonists may improve cognition, their side effects would include tachycardia and other arrhythmias. Interestingly, the effect of AD on LC neurons was greater than that on the BFB cholinergic neurons [240]. When A β was

injected in rat retrosplenial cortex, it caused a higher neuronal loss in the LC than in the BFB (nucleus basalis) [241]. Importantly, degeneration of NA neurons does not occur as a result of normal aging [242, 243]. Hypoxia has deleterious effect on neurons including those of LC and hippocampus [244–248]. They become persistently impaired after exposure to hypoxia/reoxygenation, modeling sleep apnea in adult mice [249]. The LC degeneration, therefore, is an event being closely associated with neuroinflammation and OSA. Vagal activation (or inhibition) has a correlated effect on the LC neurons, thus reflecting the influence of the NTS on the LC function [250].

The entorhinal cortex (ERC), located inside the rhinal sulcus in the olfactory area, is spatially associated with the amygdaloid complex rostrally and hippocampal formation caudally. The olfactory structures are the only primary sensory system components that have direct projections to the ERC; in addition, the cingulate cortex and thalamic afferents also innervates ERC [251]. Electrophysiological studies have confirmed the selective olfactory projection to the ERC [252–255]. The ERC forwards sensory information to the hippocampus via the perforant pathway and receives back the hippocampal information [256]. A substantial body of literature has documented that memory and learning deficits associated with AD are attributable to neuronal degeneration in the ERC. ERC ablations alone may produce memory deficits [257]. AD pathology begins in the ERC, and in AD the atrophy rate in the ERC is higher than in the hippocampus [258]. In mild cases of AD, 60% of the neurons are lost in the ERC, whereas 90% loss occurs in advanced AD [259]. The ERC atrophy precedes hippocampal atrophy and the ERC volume loss is dominant over the hippocampal volume loss in MCI [260]. Such preferential damage to the ERC contributes to the etiology of AD. ERC is particularly vulnerable to tau pathology in aging and following olfactory dysfunction. The upstream pathological factors in the latter (including infection, inflammation and environmental toxins) would have an adverse impact on ERC function. Further, there are projections from the NTS that may impact olfactory bulb function in AD [261–266]. The ERC lesion, therefore, would impair working memory and induce anterograde memory deficits [258–260, 267].

Metabolic and functional homeostasis in CNS is maintained due to the complex output of endocrine, autonomic, and behavioral control circuits. The functional relationship between the NTS neurons and a host of key foci and modulation of synaptic activity [170] identifies a pivotal role of the NTS in memory and cognition. Sensory signals from the cardiovascular, respiratory, gastrointestinal, and other systems (see the section on the NTS) are delivered to the NST; subsequently efferent signals arising within the NTS are relayed to various brain regions. The axonal

projections from the NTS to the hypothalamus and limbic regions is said to modulate behavioral arousal, and coordinate endocrine [268–273] and behavioral responses to stress (including infection and inflammation) [272, 273]. The potential physiological significance of these connections and mechanisms is immense in terms of central control of a vast array of functions. Finally, since stimulation of vagus nerve (arises from the NTS as mentioned above) has been reported to enhance memory in animals and humans, the underlying neuroanatomical-physiological substrate primarily responsible for modulating memory is the NTS. The dysfunctional NTS conceivably would lead to dysfunctional synaptic activity and may negatively impact memory in aging and AD [71, 201–203, 250, 264–266]. Accordingly, since the NTS may have an etiological role in AD, it represents an appropriate target for treatment of attention, arousal, cognition and memory dysfunction.

Discussion

Inflammatory Pathology in Aging and AD

Aging is a low-grade chronic subclinical state of inflammation that may promote age-associated diseases including AD. Ongoing inflammatory cascades may enhance pathologies including vascular endothelial cell damage, ischemia, hypoxia/hypoxemia, and cardiovascular conditions [2–4, 11–13, 18, 36, 164]. These may further exacerbate the expression of proinflammatory cytokines in the elderly [106–108]. Systemic inflammation has an impact on the microglia and switches them to a primed phenotype. Thus activated microglia become hyperresponsive to inflammatory stimuli [3, 18, 36, 164] in the aged CNS; they release proinflammatory cytokines that may exceed the anti-inflammatory mediators [22, 38, 39] and promote neurodegeneration [109, 274–277]. In both humans and transgenic mouse models of AD, the increased inflammatory modulators are correlated with the onset of AD neuropathology [109, 276, 277]. The former include the proinflammatory cytokines such as IL-1 β , IL-6, TNF- α and TGF- β , chemokines, the complement pathway, coagulation factors; α -2-macroglobulin and α 1-ACT (acute-phase reactive proteins) [25]. The Rotterdam Study (6,713 subjects) found a relationship between elevated levels of α 1antichymotrypsin, CRP, IL-6, soluble ICAM and VCAM, and an increased risk for AD [278].

Aging Decreases NTS Volume

Applying strict exclusion criteria, 152 “healthy older adults without any disease”, (mean age = 66), except for “undiagnosed sleep-related breathing disorders (SRBD)”, and an

AHI of 21 (\pm 15) for males, and 16.7 (\pm 12.9) for females, were studied [279]. The absence of subjective daytime sleepiness and a lack of involvement of any cortical or subcortical structures in these elderly underscored their overall health. Furthermore, their whole brain Gray matter volume (GMV) scans obtained using voxel-based MRI morphometry did not differ significantly between subjects with and without SRBD. However, bilaterally, a significant inverse relationship was found between AHI and GMV in the pontomedullary zone—involving (1) NTS, (2) NA, and (3) the DMNV [279]. The symmetrical atrophy of these key brainstem nuclei was interpreted as a function of episodic hypoxic and sleep-disordered conditions. The GMV loss in specific brainstem nuclei in asymptomatic elderly reflects an ongoing silent subclinical anatomical and pathophysiological change. Apart from their subclinical breathing abnormalities during sleep, the only other abnormalities these elderly possibly suffered from were subclinical ongoing decreases in olfactory, gustatory, and somatosensory modalities of senescence [89, 280, 281], and aging-related inflammation rendering the brainstem nuclei dysfunctional, including the NTS. Conceivably, during aging decreases occur in sensory modalities (thalamocortical system) and this may affect the NTS \rightarrow Hypoglossal \rightarrow genioglossus activity, attenuating the pharyngeal patency; this would result in apnea/hypopnea and hence hypoxia/hypoxemia during sleep. The recurring intermittent hypoxic episodes during sleep may potentiate neuropathology in key brainstem regions, and neocortex including the parietal, temporal, and frontal lobes, and the BFB. Conceivably, these neuropathological alterations may underpin cognitive dysfunction.

Hypoxic Injury and NTS Pathology

The NTS is a pivotal region for regulating the set-point of arterial pressure; CIH has been linked to elevations of sympathetic activity and mean arterial pressure. A recent study reports that the NTS hypoxia increases arterial pressure [282]. Hypoxia stimulates the expression of inflammatory cytokines (TNF- α , IL-1 β), chemokines (IL-8, MCP-1/CCL2), and adhesion molecules (ICAM-1) in the brain and in cultured human and mouse astrocytes, and brain endothelial cells [283, 284]. Hypoxia-induced upregulation of inflammatory genes is regulated by different transcription factors including activator protein-1, NF κ B, and HIF-1. HIF-1 is an essential molecule that regulates oxygen homeostasis and mediates hypoxia-induced expression of proinflammatory mediators including IL-1 β [284–286]. Astroglial cells are the most abundant cells in the brain and play an important role in the initiation and progression of hypoxia-induced neuroinflammation. OSA subjects suffer from microarousals and enhanced sympathetic activity; however, it is important to highlight

that age-related oxidative stress itself is sufficient to promote vascular inflammation even in the absence of other common risk factors [286]. Recent experimental data suggest that ROS and multiple proinflammatory pathways may converge on NF- κ B to enhance its transcriptional activity, and induce endothelial activation in aged vasculature [286]. Furthermore, in the NTS neurons, hypoxia decreases the amplitude of excitatory postsynaptic currents [287].

Significant cognitive impairment occurs in patients with moderate to severe OSA, and this is correlated with tissue damage in regions involved in cognition. Various data have shown focal loss of grey matter volume (GMV) and hence neuronal loss in cognitively relevant brain regions in OSA/hypoxia [288]. In animal model of OSA, the cortical neuronal cell death due to CIH has been documented causing neurocognitive dysfunction [108, 289]. Neurocognitive dysfunction associated with sleep apnea has been documented in elderly patients with MCI [290]. Another mechanism analogous to CIH is the ischemic hypoxia/reperfusion-related reoxygenation, where enhanced ROS generation may cause damage. Decreased brain perfusion has been documented after apneic episodes in OSA patients [291]; further, hypoxia has been shown to cause severe pathology in the NTS neurons [292].

Neuronal Apoptosis in the NTS

OSA is characterised by repetitive obstruction of the UA during sleep due to failure of the principal dilator muscle genioglossus to oppose the pharyngeal collapse. The net result is a condition characterized by intermittent apnea and hypoxia. However, despite its great importance little is known about the pathways proximal/prior to genioglossus dysfunction and UA collapse. The important role of snoring/OSA in causing degeneration of UA afferent nerve fibers that impacts the sensory limb of the sensory-motor reflex has been emphasized [280, 281]. It was also pointed out that the NTS that integrates UA sensory inputs may be rendered dysfunctional owing to paucity of UA sensory stimulation. Since the NTS projects to the hypoglossal neurons [28, 29] it impacts the genioglossus function that causes UA obstruction/collapse \rightarrow snoring to overcome the UA negative pressure \rightarrow apneic hypoxia. The question still remains as to what causes the NTS dysfunction in the first place that evokes a cascade culminating in UA obstruction and snoring; the pathology of sensory nerve fibers of UA being downstream to snoring and pharyngeal suction collapse. It is posited here that it is the ongoing systemic inflammatory episodes (from disparate sources detailed above) and the resultant neuroinflammation which are the harbinger of neuronal dysfunction/degeneration in CNS in general, and the NTS neurons in particular.

Consequently, neuroinflammation, ROS production, oxidative stress gene activation, protein oxidation, lipid and nucleic acid oxidation, alter the neuronal homeostasis in the NTS and other brainstem nuclei. This conclusion is supported by experimental studies employing ROS/Neuronal degeneration approach [293]. Consequently, neuroinflammation, intermittent/episodic apnea/hypopnea, and ROS/oxidative stress would synergize to augment the NTS pathology [19, 108, 292, 294, 295]. There is compelling evidence, therefore, to support the above explanation for neuronal dysfunction—and the epicentre of this pathology is the multifunctional highly interconnected NTS.

Hypoxia and other risk factors described above upregulate innate inflammation and neuroinflammation causing release of proinflammatory cytokines from several brain foci. OSA was found to be independently associated with TNF- α level [293]; this cytokine has been shown to alter synaptic transmission and plasticity [296], and it is a crucial mediator of neuronal cell death. The NTS is highly vulnerable to ischemia and/or hypoxia owing to its intense metabolic activity. Important data are available on the anatomic distribution of apoptosis of neurons and glial cells in the medullary nuclei of adult subjects who died of hypoxic injury [294]. The apoptotic index was: NTS 11.2, DMNV 6.8, and hypoglossal nucleus 6.6. In another study of human autopsied brainstems (15 adults, 25–58 year old), the percentage of TUNEL-positive neurons in the NTS was 8.5% and DMNV was 5.4%, and the apoptosis was ascribed to hypoxic-ischemic injury [295]. In a guinea pig model of intermittent apnea (mimicking closely to that occurs in humans) recent elegant studies have clearly documented that recurrent periods of apnea induce apoptotic neurodegeneration in nuclei throughout the brainstem [291]. Although apoptotic neurons were widespread, the highest density of apoptotic neurons was present in the facial, hypoglossal, ambiguus, DMNV, and the NTS [297]. In addition to the nuclei involved in the regulation of respiratory, cardiovascular, sleep/wake, and autonomic functions, a great number of apoptotic neurons were present in other areas of the pons and medulla as well. Since the NTS is a major coordinating center, apoptotic degeneration of neurons in this particular nucleus and those linked to this nucleus may contribute significant deficits in a number of functions throughout the CNS.

Invading Pathogens Cause NTS Dysfunction

In addition to inflammation, hypertension, diabetes mellitus, hypoxia/hypoxemia, and attenuated cerebral blood flow (CBF), the Intra-cerebral presence of microorganism is also an important risk factor in promoting cognitive dysfunction and AD. An invading pathogen may generate an inflammatory acute response and commence a cascade

of signalling mechanisms, including activation of cytokines, complement system, and recruitment of the immune cells to the site of inflammation. An example of recurring acute viral insult is herpes simplex virus type 1 (HSV1) that remains in a latent state in the peripheral nervous system. The life-long dormancy is punctuated by repeated activation (due to different stresses), reaching different brain areas such as frontal and temporal sites (known to be involved in AD)—and causing damage. The virus is ubiquitous infecting about 90% of adult population, and PCR has confirmed the viral presence in brains of the elderly with and without AD [298]. Other viri including influenza PR8 virus may gain entry via nasal and/or oral portals, promote CNS neuroinflammation, and cause widespread neuronal dysfunction, including those of the NTS. Similar to HSV1, hepatitis, HIV, West Nile virus, *Chlamydia pneumoniae* and *Borrelia burgdorferi* (to name just a few), are capable of causing CNS pathology [299], as are other bacteria [70]. Injection of pseudorabies virus in the rat genioglossus produced viral labeling in neurons of the hypoglossal and the NTS nuclei [300]. When present in the brain cells, microorganism(s) enhance microglial activation, cause release of neurotoxic agents, and lead to neuronal injury and degeneration. Thus stigmata such as cerebral infection, neuroinflammation, hypoxia/hypoxemia, and perturbed CBF are assumed to occur in the NTS, and may impact its neuronal function adversely. In order to gain entry into the NTS, any infection need not alter the integrity/permeability of the BBB. Microorganisms can cross the BBB and enter the brain through the so-called “Trojan horse mechanism” [301]. *Chlamydia pneumoniae* e.g. is known to reside in the peripheral blood monocyte and leucocytes, and could exploit the Trojan horse mechanism to enter the CNS over an intact BBB [302]. NTS is adjacent to the AP [158] and lacks a BBB; the AP has prominent axonal projections to the NTS, and provides an anatomical pathway whereby various circulating inflammation-causing mediators can gain access and affect the NTS activity.

Olfactory Nerve Transmits Infection to the NTS

The olfactory nerve (ON) is an important pathway for transmission of viruses into the brain owing to its direct contact with the environment. When influenza virus PR8 is intranasally introduced in mice, signals from the ON triggers the acute phase response (APR); however, the ON transection prior to influenza viral infection reduced the number of viral antigen-TNF- α - and IL1 β -immunoreactive cells in the olfactory bulb (OB) and other interconnected brain regions [302]. As early as 15 h post-infection, the number of cells expressing TNF- α and IL1- β mRNAs and proteins increase in the OB, as well as in neurons within

different regions of the brain including the amygdala, hypothalamus, and central autonomic system regions [302]. All above mentioned areas project reciprocally to the NTS (see section on NTS above).

Vagus Nerve Transmits Infection to the NTS

By having a “wandering” route through the body, the vagus is uniquely equipped to provide (a) an effective early warning system about the pathogenic presence, and (b) a feedback to the immune system when pathogens are cleared. Several studies have demonstrated that neurotropic viruses travels to the CNS mainly via the vagus nerve. In mice infected intranasally with influenza (either virus strain 24a5b or avian influenza A-H5N3), the virus travel to the NTS via the vagus nerve [303, 304], moreover, the nucleus primarily affected was the NTS [303, 304]. Following administration of the neurotropic avian influenza virus H5N1 in C57BL/6J mice, the virus is transmitted into the CNS via the vagus. Other infections including the fatal transmissible spongiform encephalopathies or prion disease spread from the periphery to the brain, through an oral route. The first target area in the brain of orally infected scrapie is the DMNV, followed rapidly by the NTS—again via the vagus nerve [305]. The above data clearly show that viri from the oral, nasal or lower respiratory mucosa are transmitted to the CNS via the vagus; they spread trans-synaptically in the CNS inducing lesions in the brainstem nuclei, including the NTS. Thus, via vagus, microbes in the periphery are relayed to the NTS for mounting a specific and effective response; it is here in the NTS where the afferent and efferent aspects of the parasympathetic nervous system meet [168]. The efferent parasympathetic pathway forms “the cholinergic anti-inflammatory pathway” [166, 171, 172]. On the efferent side, since the NTS has direct and indirect connections to a wide range of neural structures in the CNS, it has the capacity to influence their physiological processes [165].

An Inflamed NTS Is a Dysfunctional NTS

Brain mRNA is replete with mRNA of systemic cytokines; systemic IL1- β e.g. increases brain IL1- β mRNA, and vagotomy (subdiaphragmatic) blocks this [306]. Interestingly, relative to control (saline), systemic administration of proinflammatory cytokines TNF- α or IL1- β also increased IL1- β , TNF- α and IL-6 mRNAs in the NTS, but did not increase the anti-inflammatory IL-10 cytokine [307]. Recent studies have implicated the microvasculature inflammation in the brainstem, specifically in the NTS, in the pathogenesis of hypertension [308, 309]. It has been shown that vessels within brainstem regions of hypertensive animals are inflamed and release ROS and cytokines;

these pathological messengers then alter neuronal activity in the NTS [310–314]. The inflammatory signaling molecules released from the brainstem microvasculature exert deleterious impact on neuronal excitability in the NTS; this then alters both the central set point of arterial pressure and its reflex control. Furthermore, the brainstem vessel inflammation conceivably elevates the resistance to blood flow thus causing inadequate perfusion. This then presents a scenario of sympathetically mediated hypertension in response to inadequate blood perfusion. Conceivably, the above pathology synergized with hypoxia and other risk factors negatively impacts the NTS activity further; an inflamed and dysfunctional NTS consequently may cause widespread disruption of many key physiological functions in both brainstem and neocortex.

Recent studies have shown high levels of proinflammatory molecules expressed in the NTS of the spontaneously hypertensive rat (SHR), an animal model of human essential hypertension. In the NTS of SHR, the gp39 precursor was up-regulated [310, 311]. The gp39 precursor is homologous to chitinase 3-like protein 1, also known as human cartilage-gp39 or YKL40. Although functional roles for gp39 precursor are not fully known, high levels of this molecule are present in many inflammatory conditions including rheumatoid arthritis, glioblastoma, inflammatory bowel disease, atherosclerosis, asthma, and indeed AD. Furthermore, gp39 precursor promotes chemotaxis, and an up-regulation of gp39 precursor in the NTS of the hypertensive rats reflects a pervasive inflammatory state that may attenuate neuronal activity in both inflamed NTS, as well as its interconnected foci [310–314]. In AD, the dysfunctional NTS may differentially affect functional integrity of its projected pathways/neurons, and compromise linked key brainstem nuclei, and neocortical regions including the hippocampus, BFB, hypothalamus, insula, PFC, amygdala, and other cortical regions. There are numerous examples of neural network degradation in aging and AD. Finally, it is untenable that the NTS remains structurally and functionally insulated from the pathological effects of local and distant neuroinflammation.

The Current Hypothesis

Several CNS areas including the BFB, hippocampus, ERC, amygdala/insula, and the LC are known to be lesioned in AD. To date, a plausible unifying hypothesis that links well characterized upstream factors and AD pathology encompassing heterogeneous dysfunctional regions in AD brain is not available. Based on well documented evidence, it is posited here that the NTS is a brainstem hub nucleus that has an enormous influence/impact throughout the CNS. The current article presents a novel NTS vector hypothesis

that underpins several disparate biological mechanisms and neural circuits, and identifies relevant hallmarks of major presumptive causative factor(s) that link the NTS to AD neuropathogenesis.

The GMV decrease occurs in the CNS due to aging, inflammation, obesity, and hypoxia, as mentioned above. It is well documented that inflammatory and oxidative stress processes trigger cascade of neurodegenerative events including synaptic loss, neurotransmitter deficits, and neuronal dysfunction [315]. Input from the inflamed and dysfunctional NTS is deemed to negatively impact several foci, including the hypoglossal motor neurons. The dysfunctional hypoglossal activity causes genioglossal dysfunction, which provokes UA functional perturbations. To overcome the latter, snoring is employed as an effort to ameliorate UA obstruction that promotes periodic apnea, i.e. OSA. The resulting periodic UA obstruction/collapse and hypopnea/apnea cause intermittent hypoxia and hypoxemia. Chronic intermittent hypoxia/reoxygenation is toxic to neurons, which are already subjected to neuroinflammation and GMV. A synergistic effect of the above factors may exert deleterious impact and compromise the NTS function. Perturbations in the NTS activity cause dysfunctional processing of sensory afferents from the olfactory, gustatory, UA somatosensory, cardiovascular baroreceptor and chemoreceptor, as well as afferents from the gastrointestinal, respiratory, and other thoraco-abdominal viscera. The vagus arises from the NTS—a dysfunctional NTS may compromise and attenuate the efferent activity of the vagus; since vagus has an anti-inflammatory parasympathetic function—this will be attenuated as well. The projections from a pathological NTS (and the several nuclei it projects to, e.g. the LC, NA, DMNV, and parabrachial), to the PFC, amygdala, insula, BFB, ERC, and the hippocampus would conceivably compromise activity in the above CNS areas thus contributing to neuropsychiatric and cognitive pathology of AD (Fig. 1). An ongoing attenuation/compromise (over a period of decades) of function in many CNS pathways may result in widespread synaptic and neuronal perturbations, their disruption, degeneration, and eventual death.

Neuroinflammation—Amyloid Deposition and Neurofibrillary Tangles

Although A β deposition is considered a hall mark of AD, it occurs in unimpaired normal elderly as well [316]. This indicates that the time window for its slow rate of accumulation in clinically silent period may be quite large in the aged. An indolent sub-clinical inflammatory state persisting in the elderly (see inflammatory risk factors discussed above) causes neuroinflammation and cerebrovascular pathology in




Fig. 1 Schematic representation of the hypothesis. Several risk factors in aging enhance proinflammatory cytokines inducing systemic inflammation. This promotes neuroinflammation; activated microglia further augment central inflammation affecting neuronal, glial and vascular endothelial cell functions. The box with the asterisk denotes that dysfunctional hypoglossal motoneurons cause genioglossus dysfunction resulting in upper airway obstruction/collapse. This leads to OSA, periodic hypoxia, and hypoxemia. Inflammation, intermittent hypoxia, and oxidative stress provide a rich milieu for the NTS inflammation and dysfunction. Being a central hub for disparate afferent processing, and owing to its widespread projections in the CNS, the dysfunctional NTS would negatively impact several key physiological functions, causing impairment in cognition and memory. The ongoing synaptic dysfunction and neuronal degeneration over a prolonged period (aging), promote neuropsychiatric and cognitive deterioration, and provoke Alzheimer's dementia

the vulnerable. This enhances priming of the microglia and astrocytes, infiltration of perivascular macrophages and peripheral immunocytes that may promote neuronal dysfunction/degeneration. These processes may cause reactive A β generation. In upregulated neuroinflammation, there is a mismatch between the level of A β synthesis and its removal by microglia; this may lead to further inflammation and excess of A β deposition. Thus, in neuroinflammation, A β formation/deposition is a downstream phenomenon. Furthermore, A β has been shown to function as an antimicrobial peptide (AMP) being active against common and clinically relevant microorganisms [317, 318]. Since several infectious agents have been reported in CNS of the elderly

and AD patients [70] (see the section on invading pathogens cause NTS dysfunction), A β may be synthesized as part of a response triggered by the immune system.

AD is characterized by neurofibrillar degeneration; the latter occurs as intraneuronal NFT in dementia. Hyperphosphorylated tau (p-tau) is considered to occupy a central position in AD neuropathogenesis. In Tg4510 mice, p-tau is associated with age-related microglial activation, and in normal aging up to 9 mo CD45+ microglia increase in parallel with tau pathology [319]. 1–3 month old young mice are capable of clearing soluble p-tau; however, in 5.5 month-old mice insoluble tau aggregates are present [319, 320]. In LPS injected APP mice, activated microglia cleared A β [321]. However, LPS-induced microglial activation exacerbated p-tau pathology in Tg4510 mice [319]. Interestingly, peripheral administration of LPS showed no changes of APP processing, while p-tau was increased in both hippocampus and the ERC [322]—important regions in AD pathogenesis. Moreover, proinflammatory cytokine IL-1 β caused tau phosphorylation in neurons [323]. In 3xTgAD mice also, chronic TNF signalling enhanced p-tau causing neuronal death [324]. The above studies support the potential role of inflammatory mediators in enhancing p-tau pathology and underscore that inflammatory stimuli can and do facilitate tau phosphorylation potentiating tauopathy. Similar results however have been obtained in rats undergoing chronic neuroinflammation following LPS administration [325]. Such a scenario may occur in the CNS and the NTS in particular, in the elderly and AD patients (who possess a chronic inflammatory state as delineated above). In sum, the dysfunctional NTS may affect the functional integrity of a large number of key pathways/synapses/neurons, and provoke cognitive dysfunction/dementia in the elderly.

Conclusions

The immune system and inflammation are implicated in aging and a wide variety of neurodegenerative conditions. Relatively common sources of systemic inflammation may be significant risk factors to potentiate neuroinflammation in the elderly. This would cause microvascular changes, switching of microglial phenotype and activity, and perturbation of physiological functions in the CNS including the key brainstem nuclei, viz. the NTS and the hypoglossal. Their dysfunctional activity results in genioglossal dysfunction that may lead to UA obstruction and intermittent hypoxia/hypoxemia. The latter has widespread impact on CNS structure and function by enhancing further inflammation and exacerbating GMV decrease in aging. Neuroinflammation plus intermittent hypoxia synergize to promote escalation of CNS pathology. The NTS is the

central integration hub for UA somatosensory, gastrointestinal, gustatory, baroreceptor, chemoreceptor, and several other afferents e.g. from the amygdala, PFC, and hypothalamus, as well as sympathetic and parasympathetic systems. On the efferent side, the NTS projects to most, if not the entire, CNS. A pragmatic hypothesis is posited on the neuropathogenesis of AD emphasizing the important role of upstream factors. The current hypothesis implicates the inflamed and dysfunctional NTS as an upstream central player in the neuropathogenesis of cognitive and neuropsychiatric impairment. In defining etiopathology of AD, the current hypothesis is the first to connect: Aging → proinflammatory mediators, neuroinflammation and oxidative stress → dysfunctional NTS → impaired function of hypoglossal nucleus → depressed genioglossal function and hence snoring/OSA → hypoxia and hypoxemia → further exacerbation of neuroinflammation/oxidative stress → widespread neuronal degeneration → global deleterious impact on key physiological functions in CNS → neuropsychiatric, memory and cognitive dysfunction → AD. Vagus arises from the NTS; the vagal function is anti-inflammatory and hence important in ameliorating inflammation and upregulating functions towards homeostasis. However, a dysfunctional NTS would compromise and attenuate vagal activity, resulting in increased CNS susceptibility to the effects of disparate pathological factors, and facilitating neuropathogenesis in aging and AD. The implications of the current perspective are considerable and future research is warranted.

Conflict of interest I have no conflict of interest.

References

- Giannakopoulos P, Hof PR, Michel JP, Guimon J, Bouras C (1997) Cerebral cortex pathology in aging and Alzheimer's disease: a quantitative survey of large hospital-based geriatric and psychiatric cohorts. *Brain Res Brain Res Rev* 25:217–245
- Chung HY, Cesari M, Anton S, Marzetti E, Giovannini S, Seo AY, Carter C, Yu BP, Leeuwenburgh C (2009) Molecular inflammation: underpinnings of aging and age-related diseases. *Ageing Res Rev* 8:18–30
- Deng XH, Bertini G, Xu YZ, Yan Z, Bentivoglio M (2006) Cytokine-induced activation of glial cells in the mouse brain is enhanced at an advanced age. *Neuroscience* 141:645–661
- Choi DY, Zhang J, Bing G (2010) Aging enhances the neuroinflammatory response and alpha-synuclein nitration in rats. *Neurobiol Aging* 31:1649–1653
- Mattson MP (2002) Oxidative stress, perturbed calcium homeostasis, and immune dysfunction in Alzheimer's disease. *J Neurovirol* 8:539–550
- Butterfield DA (2002) Amyloid beta-peptide (1–42)-induced oxidative stress and neurotoxicity: implications for neurodegeneration in Alzheimer's disease brain. A review. *Free Radic Res* 36:1307–1313
- Praticò D (2008) Evidence of oxidative stress in Alzheimer's disease brain and antioxidant therapy: lights and shadows. *Ann N Y Acad Sci* 1147:70–78
- Clark TA, Lee HP, Rolston RK, Zhu X, Marlatt MW, Castellani RJ, Nunomura A, Casadesus G, Smith MA, Lee HG, Perry G (2010) Oxidative stress and its implications for future treatments and management of Alzheimer disease. *Int J Biomed Sci* 6: 225–227
- Head E, Liu J, Hagen TM, Muggenburg BA, Milgram NW, Ames BN, Cotman CW (2002) Oxidative damage increases with age in a canine model of human brain aging. *J Neurochem* 82: 375–381
- Montiel T, Quiroz-Baez R, Massieu L, Arias C (2006) Role of oxidative stress on beta-amyloid neurotoxicity elicited during impairment of energy metabolism in the hippocampus: protection by antioxidants. *Exp Neurol* 200:496–508
- Holmes C, El-Okli M, Williams AL, Cunningham C, Wilcockson D, Perry VH (2003) Systemic infection, interleukin 1beta, and cognitive decline in Alzheimer's disease. *J Neurol Neurosurg Psychiatry* 74:788–789
- Richwine AF, Sparkman NL, Dilger RN, Buchanan JB, Johnson RW (2009) Cognitive deficits in interleukin-10-deficient mice after peripheral injection of lipopolysaccharide. *Brain Behav Immun* 23:794–802
- Chen J, Buchanan JB, Sparkman NL, Godbout JP, Freund GG, Johnson RW (2008) Neuroinflammation and disruption in working memory in aged mice after acute stimulation of the peripheral innate immune system. *Brain Behav Immun* 22:301–311
- Swardfager W, Lanctôt K, Rothenburg L, Wong A, Cappell J, Herrmann N (2010) A meta-analysis of cytokines in Alzheimer's disease. *Biol Psychiatry* 68:930–941
- Ye SM, Johnson RW (2001) An age-related decline in interleukin-10 may contribute to the increased expression of interleukin-6 in brain of aged mice. *Neuroimmunomodulation* 9:183–192
- Tan ZS, Seshadri S (2010) Inflammation in the Alzheimer's disease cascade: culprit or innocent bystander? *Alzheimers Res Ther* 2:6
- Wofford JL, Loehr LR, Schwartz E (1996) Acute cognitive impairment in elderly ED patients: etiologies and outcomes. *Am J Emerg Med* 14:649–653
- Godbout JP, Chen J, Abraham J, Richwine AF, Berg BM, Kelley KW, Johnson RW (2005) Exaggerated neuroinflammation and sickness behavior in aged mice following activation of the peripheral innate immune system. *FASEB J* 19:1329–1331
- Zotova E, Nicoll JA, Kalara R, Holmes C, Boche D (2010) Inflammation in Alzheimer's disease: relevance to pathogenesis and therapy. *Alzheimers Res Ther* 2:1
- Barrientos RM, Higgins EA, Biedenkapp JC et al (2006) Peripheral infection and aging interact to impair hippocampal memory consolidation. *Neurobiol Aging* 27:723–732
- Tanaka S, Ide M, Shibutani T et al (2006) Lipopolysaccharide-induced microglial activation induces learning and memory deficits without neuronal cell death in rats. *J Neurosci Res* 83: 557–566
- Perry VH, Cunningham C, Holmes C (2007) Systemic infections and inflammation affect chronic neurodegeneration. *Nat Rev Immunol* 7:161–167
- Torelli F, Moscufo N, Garreffa G, Placidi F, Romigi A, Zannino S, Bozzali M, Fasano F, Giulietti G, Djonlagic I, Malhotra A, Marciani MG, Guttmann CR (2011) Cognitive profile and brain morphological changes in obstructive sleep apnea. *Neuroimage* 54:787–793
- Canessa N, Castronovo V, Cappa SF, Aloia MS, Marelli S, Falini A, Alemanno F, Ferini-Strambi L (2011) Obstructive sleep apnea: brain structural changes and neurocognitive

- function before and after treatment. *Am J Respir Crit Care Med* 183:1419–1426
25. Wyss-Coray T (2006) Inflammation in Alzheimer disease: driving force, bystander or beneficial response? *Nat Med* 12:1005–1015
 26. Loewy AD (1990) Central autonomic pathways. In: Loewy AD, Spyer KM (eds) *Central regulation of autonomic functions*. Oxford University Press, Oxford, pp 88–103
 27. Loewy AD (1998) Viruses as transneuronal tracers for defining neural circuits. *Neurosci Biobehav Rev* 22:679–684
 28. Beckman ME, Whitehead MC (1991) Intramedullary connections of the rostral nucleus of the solitary tract in the hamster. *Brain Res* 557:265–279
 29. Borke RC, Nau ME, Ringler RL Jr (1983) Brain stem afferents of hypoglossal neurons in the rat. *Brain Res* 269:47–55
 30. Onai T, Saji M, Miura M (1987) Functional subdivisions of the nucleus tractus solitarius of the rat as determined by circulatory and respiratory responses to electrical stimulation of the nucleus. *J Auton Nerv Syst* 21:195–202
 31. Streefland C, Jansen K (1999) Intramedullary projections of the rostral nucleus of the solitary tract in the rat: gustatory influences on autonomic output. *Chem Senses* 24:655–664
 32. Akiyama H, Barger S, Barnum S, Bradt B, Bauer J, Cole GM (2000) Inflammation and Alzheimer's disease. *Neurobiol Aging* 21:383–421
 33. Eikelenboom P, Veerhuis R, Scheper W, Rozemuller AJ, van Gool WA, Hoozemans JJ (2006) The significance of neuroinflammation in understanding Alzheimer's disease. *J Neural Transm* 113:1685–1695
 34. Böhner H, Qiu F, Zimmermann T, Zhang Y, Jilmer T, Männel D, Böttiger BW, Stern DM, Waldherr R, Saeger HD, Ziegler R, Bierhaus A, Martin E, Nawroth PP (1997) Role of NF-kappaB in the mortality of sepsis. *J Clin Invest* 100:972–985
 35. Makarov SS (2000) NF-kappaB as a therapeutic target in chronic inflammation: recent advances. *Mol Med Today* 6:441–448
 36. Henry CJ, Huang Y, Wynne AM, Godbout JP (2009) Peripheral lipopolysaccharide (LPS) challenge promotes microglial hyperactivity in aged mice that is associated with exaggerated induction of both pro-inflammatory IL-1beta and anti-inflammatory IL-10 cytokines. *Brain Behav Immun* 23:309–317
 37. Zou Y, Jung KJ, Kim JW, Yu BP, Chung HY (2004) Alteration of soluble adhesion molecules during aging and their modulation by calorie restriction. *FASEB J* 18:320–322
 38. Kohman RA, Crowell B, Kusnecov AW (2010) Differential sensitivity to endotoxin exposure in young and middle-age mice. *Brain Behav Immun* 24:486–492
 39. Sawada H, Hishida R, Hirata Y, Ono KK, Suzuki H, Muramatsu S, Nakano I, Nagatsu T, Sawada M (2007) Activated microglia affect the nigro-striatal dopamine neurons differently in neonatal and aged mice treated with 1-methyl-4-phenyl-1, 2, 3, 6-tetrahydropyridine. *J Neurosci Res* 85:1752–1761
 40. Gaykema RP, Balachandran MK, Godbout JP, Johnson RW, Goehler LE (2007) Enhanced neuronal activation in central autonomic network nuclei in aged mice following acute peripheral immune challenge. *Auton Neurosci* 131:137–142
 41. Casserly I, Topol E (2004) Convergence of atherosclerosis and Alzheimer's disease: inflammation, cholesterol, and misfolded proteins. *Lancet* 363:1139–1146
 42. Ishida Y, Shirokawa T, Miyaishi O, Komatsu Y, Isobe K (2000) Age-dependent changes in projections from locus coeruleus to hippocampus dentate gyrus and frontal cortex. *Eur J Neurosci* 12:1263–1270
 43. O'Neil JN, Mouton PR, Tizabi Y, Ottinger MA, Lei DL, Ingram DK, Manaye KF (2007) Catecholaminergic neuronal loss in locus coeruleus of aged female dtg APP/PS1 mice. *J Chem Neuroanat* 34:102–107
 44. Kennedy DJ, Kuchibhotla S, Westfall KM, Silverstein RL, Morton RE, Febbraio MA (2011) CD36-dependent pathway enhances macrophage and adipose tissue inflammation and impairs insulin signalling. *Cardiovasc Res* 89:604–613
 45. Annone F, Lapadula G (2010) Obesity and inflammation-targets for OA therapy. *Curr Drug Targets* 11:586–598
 46. Vandanmagsar B, Youm YH, Ravussin A, Galgani JE, Stadler K, Mynatt RL, Ravussin E, Stephens JM, Dixit VD (2011) The NLRP3 inflammasome instigates obesity-induced inflammation and insulin resistance. *Nat Med* 17:179–188
 47. Look C, Morano I, Ehrhart-Bornstein M, Bornstein SR, Lamounier-Zepter V (2011) Adipocyte-derived factors suppress heart contraction. *Int J Obes* 35:84–90
 48. Zanni MV, Stanley TL, Makimura H, Chen CY, Grinspoon SK (2010) Effects of TNF-alpha antagonism on E-selectin in obese subjects with metabolic dysregulation. *Clin Endocrinol (Oxf)* 73:48–54
 49. Kivipelto M, Ngandu T, Fratiglioni L, Viitanen M, Kåreholt I, Winblad B, Helkala EL, Tuomilehto J, Soininen H, Nissinen A (2005) Obesity and vascular risk factors at midlife and the risk of dementia and Alzheimer disease. *Arch Neurol* 62:1556–1560
 50. Rosengren A, Skoog I, Gustafson D, Wilhelmsen L (2005) Body mass index, other cardiovascular risk factors, and hospitalization for dementia. *Arch Intern Med* 165:321–326
 51. Whitmer RA, Gunderson EP, Barrett-Connor E (2005) Obesity in middle age and future risk of dementia: a 27 year longitudinal population based study. *Br Med J* 330:1360
 52. Gustafson D, Lissner L, Bengtsson C, Björkelund C, Skoog I (2004) A 24-year follow-up of body mass index and cerebral atrophy. *Neurology* 63:1876–1881
 53. Walther K, Birdsill AC, Glisky EL, Ryan L (2010) Structural brain differences and cognitive functioning related to body mass index in older females. *Hum Brain Mapp* 31:1052–1064
 54. Raji CA, Ho AJ, Parikshak NN, Becker JT, Lopez OL, Kuller LH, Hua X, Leow AD, Toga AW, Thompson PM (2010) Brain structure and obesity. *Hum Brain Mapp* 31:353–364
 55. Wolf PA, Beiser A, Elias MF, Au R, Vasan RS, Seshadri S (2007) Relation of obesity to cognitive function: importance of central obesity and synergistic influence of concomitant hypertension. The Framingham Heart Study. *Curr Alzheimer Res* 4:111–116
 56. Hofman A, Ott A, Breteler MM, Bots ML, Slieter AJ, van Harskamp F, van Duijn CN, Van Broeckhoven C, Grobbee DE (1997) Atherosclerosis, apolipoprotein E, and prevalence of dementia and Alzheimer's disease in the Rotterdam Study. *Lancet* 349:151–154
 57. Chandra V, Pandav R (1998) Gene-environment interaction in Alzheimer's disease: a potential role for cholesterol. *Neuroepidemiology* 17:225–232
 58. Romas SN, Tang MX, Berglund L, Mayeux R (1999) APOE genotype, plasma lipids, lipoproteins, and AD in community elderly. *Neurology* 53:517–521
 59. Kivipelto M, Helkala EL, Laakso MP, Hanninen T, Hallikainen M, Alhainen K et al (2002) Apolipoprotein E epsilon4 allele, elevated midlife total cholesterol level, and high midlife systolic blood pressure are independent risk factors for late-life Alzheimer disease. *Ann Intern Med* 137:149–155
 60. Kivipelto M, Helkala EL, Hanninen T, Laakso MP, Hallikainen M, Alhainen K et al (2001) Midlife vascular risk factors and late-life mild cognitive impairment: a population-based study. *Neurology* 56:1683–1689
 61. Peila R, Rodriguez BL, Launer LJ (2002) Type 2 diabetes, APOE gene, and the risk for dementia and related pathologies: the Honolulu-Asia Aging Study. *Diabetes* 51:1256–1262
 62. Winther B, Gwaltney JM Jr, Mygind N, Hendley JO (1988) Viral-induced rhinitis. *Am J Rhinol* 12:17–20

63. Gwaltney JM (2002) Clinical significance and pathogenesis of viral respiratory infections. *Am J Med* 112(Suppl 6A):13S–18S
64. Bruewer M, Luegering A, Kucharzik T, Parkos CA, Madara JL, Hopkins AM, Nusrat A (2003) Proinflammatory cytokines disrupt epithelial barrier function by apoptosis independent mechanisms. *J Immunol* 171:6164–6172
65. Vieira SM, Lemos HP, Grespan R, Napimoga MH, Dal-Secco D, Freitas A, Cunha TM, Verri WA Jr, Souza-Junior DA, Jamur MC, Fernandes KS, Oliver C, Silva JS, Teixeira MM, Cunha FQ (2009) A crucial role for TNF- α in mediating neutrophil influx induced by endogenously generated or exogenous chemokines, KC/CXCL1 and LIX/CXCL5. *Br J Pharmacol* 158:779–789
66. Wang S, Quang Le T, Chida J, Cisse Y, Yano M, Kido H (2010) Mechanisms of matrix metalloproteinase-9 upregulation and tissue destruction in various organs in influenza A virus infection. *J Med Invest* 57:26–34
67. de La Serre CB, Ellis CL, Lee J, Hartman AL, Rutledge JC, Raybould HE (2010) Propensity to high-fat diet-induced obesity in rats is associated with changes in the gut microbiota and gut inflammation. *Am J Physiol Gastrointest Liver Physiol* 299:G440–G448
68. Cani PD, Bibiloni R, Knauf C, Waget A, Neyrinck AM, Delzenne NM, Burcelin R (2010) Changes in gut microbiota control metabolic endotoxemia-induced inflammation in high-fat diet-induced obesity and diabetes in mice. *Diabetes* 57:1470–1481
69. Lupp C, Robertson ML, Wickham ME et al (2007) Host-mediated inflammation disrupts the intestinal microbiota and promotes the overgrowth of *Enterobacteriaceae*. *Cell Host Microbe* 2:119–129
70. Kountouras J, Boziki M, Gavalas E, Zavos C, Deretzi G, Grigoriadis N, Tsolaki M, Chatzopoulos D, Katsinelos P, Tzilves D, Zabouri A, Michailidou I (2009) Increased cerebrospinal fluid *Helicobacter pylori* antibody in Alzheimer's disease. *Int J Neurosci* 119:765–777
71. Chapman TR, Barrientos RM, Ahrendsen JT, Hoover JM, Maier SF, Patterson SL (2011) Aging and infection reduce expression of specific brain-derived neurotrophic factor mRNAs in hippocampus. *Neurobiol Aging* [Epub ahead of print]
72. Wannamethee SG, Lowe GD, Shaper AG, Rumley A, Lennon L, Whincup PH (2005) Associations between cigarette smoking, pipe/cigar smoking, and smoking cessation, and haemostatic and inflammatory markers for cardiovascular disease. *Eur Heart J* 26:1765–1773
73. Helmersson J, Larsson A, Vessby B, Basu S (2005) Active smoking and a history of smoking are associated with enhanced prostaglandin F(2 α), interleukin-6 and F2-isoprostane formation in elderly men. *Atherosclerosis* 181:201–207
74. Durazzo TC, Meyerhoff DJ, Nixon SJ (2011) A comprehensive assessment of neurocognition in middle-aged chronic cigarette smokers. *Drug Alcohol Depend* [Epub ahead of print]
75. Bailey MT, Kinsey SG, Padgett DA, Sheridan JF, Leblebicioglu B (2009) Social stress enhances IL-1 β and TNF- α production by *Porphyromonas gingivalis* lipopolysaccharide-stimulated CD11b+ cells. *Physiol Behav* 98:351–358
76. Gibb J, Hayley S, Gandhi R, Poulter MO, Anisman H (2008) Synergistic and additive actions of a psychosocial stressor and endotoxin challenge: circulating and brain cytokines, plasma corticosterone and behavioral changes in mice. *Brain Behav Immun* 22:573–589
77. Curry JM, Hanke ML, Piper MG, Bailey MT, Bringardner BD, Sheridan JF, Marsh CB (2010) Social disruption induces lung inflammation. *Brain Behav Immun* 24:394–402
78. Frank MG, Baratta MV, Sprunger DB, Watkins LR, Maier SF (2007) Microglia serve as a neuroimmune substrate for stress-induced potentiation of CNS pro-inflammatory cytokine responses. *Brain Behav Immun* 21:47–59
79. O'Connor MF, Irwin MR, Wellisch DK (2009) When grief heats up: pro-inflammatory cytokines predict regional brain activation. *Neuroimage* 47:891–896
80. Jankord R, Zhang R, Flak JN, Solomon MB, Albertz J, Herman JP (2010) Stress activation of IL-6 neurons in the hypothalamus. *Am J Physiol Regul Integr Comp Physiol* 299:R343–R351
81. Bernard R, Kerman IA, Thompson RC, Jones EG, Bunney WE, Barchas JD AF, Schatzberg AF, Myers RM, Akil H, Watson SJ (2011) Altered expression of glutamate signaling, growth factor, and glia genes in the locus coeruleus of patients with major depression. *Mol Psychiatry* 16:634–646
82. Chen E, Miller GE, Kobor MS, Cole SW (2011) Maternal warmth buffers the effects of low early-life socioeconomic status on pro-inflammatory signaling in adulthood. *Mol Psychiatry* 16:729–737
83. Shelton RC, Claiborne J, Sidoryk-Wegrzynowicz M, Reddy R, Aschner M, Lewis DA, Mirmics K (2011) Altered expression of genes involved in inflammation and apoptosis in frontal cortex in major depression. *Mol Psychiatry* 16:751–762
84. Peters JL, Weisskopf MG, Spiro A 3rd, Schwartz J, Sparrow D, Nie H et al (2010) Interaction of stress, lead burden, and age on cognition in older men: the VA Normative Aging Study. *Environ Health Perspect* 118:505–510
85. Fairchild KD, Singh IS, Patel S, Drysdale BE, Viscardi RM, Hester L, Lazusky HM, Hasday JD (2004) Hypothermia prolongs activation of NF- κ B and augments generation of inflammatory cytokines. *Am J Physiol Cell Physiol* 287:C422–C431
86. Matsui T, Ishikawa T, Takeuchi H, Okabayashi K, Maekawa T (2006) Mild hypothermia promotes pro-inflammatory cytokine production in monocytes. *J Neurosurg Anesthesiol* 18:189–193
87. Matsui T, Kakeda T (2008) IL-10 production is reduced by hypothermia but augmented by hyperthermia in rat microglia. *J Neurotrauma* 25:709–715
88. Saito H, Sherwood ER, Varma TK, Evers BM (2003) Effects of aging on mortality, hypothermia, and cytokine induction in mice with endotoxemia or sepsis. *Mech Ageing Dev* 124:1047–1058
89. Daulatzai M (2010) Conversion of elderly to Alzheimer's dementia: role of confluence of hypothermia and senescent stigmata—the plausible pathway. *J Alzheimers Dis* 21:1039–1063
90. Guasti L, Marino F, Cosentino M, Maroni L, Maresca AM, Colombo F, Maio RC, Castiglioni L, Saporiti F, Loraschi A, Gaudio G, Bernasconi A, Laurita E, Grandi AM, Venco A (2011) Cytokine production from peripheral blood mononuclear cells and polymorphonuclear leukocytes in patients studied for suspected obstructive sleep apnea. *Sleep Breath* 15:3–11
91. Woodson BT, Garancis JC, Toohill RJ (1991) Histopathologic changes in snoring and obstructive sleep apnea syndrome. *Laryngoscope* 101:1318–1322
92. Bassiouny A, Nasr S, Mashaly M, Ayad E, Qotb M, Atef A (2009) Electron microscopy study of peripheral nerves in the uvulae of snorers and obstructive sleep apnoea patients. *J Laryngol Otol* 123:203–207
93. Stål PS, Lindman R, Johansson B (2009) Capillary supply of the soft palate muscles is reduced in long-term habitual snorers. *Respiration* 77:303–310
94. Loubaki L, Jacques E, Semlali A, Biardel S, Chakir J, Sériès F (2008) Tumor necrosis factor- α expression in uvular tissues differs between snorers and apneic patients. *Chest* 134:911–918
95. Almendros I, Acerbi I, Puig F, Montserrat JM, Navajas D, Farré R (2007) Upper-airway inflammation triggered by vibration in a rat model of snoring. *Sleep* 30:225–227

96. Puig F, Rico F, Almendros I, Montserrat JM, Navajas D, Farre R (2005) Vibration enhances interleukin-8 release in a cell model of snoring-induced airway inflammation. *Sleep* 28:1312–1316
97. Bloom JW, Kaltenborn WT, Quan SF (1988) Risk factors in a general population for snoring. Importance of cigarette smoking and obesity. *Chest* 93:678–683
98. Nagayoshi M, Yamagishi K, Tanigawa T, Sakurai S, Kitamura A, Kiyama M, Imano H, Ohira T, Sato S, Sankai T, Iso H (2011) Risk factors for snoring among Japanese men and women: a community-based cross-sectional study. *Sleep Breath* 15:63–69
99. Van de Berg JF, Neven AK, Tulen JH, Hofman A, Witteman JC, Miedema HM, Tiemeier H (2008) Actigraphic sleep duration and fragmentation are related to obesity in the elderly: the Rotterdam Study. *Int J Obes (Lond)* 32:1083–1090
100. Lavie L (2003) Obstructive sleep apnea syndrome—an oxidative stress disorder. *Sleep Med Rev* 7:35–51
101. Yamauchi M, Nakano H, Maekawa J, Okamoto Y, Ohnishi Y, Suzuki T, Kimura H (2005) Oxidative stress in obstructive sleep apnea. *Chest* 127:1674–1679
102. Shamsuzzaman AS, Winnicki M, Lanfranchi P, Wolk R, Kara T, Accurso V, Somers VK (2002) Elevated C-reactive protein in patients with obstructive sleep apnea. *Circulation* 105:2462–2464
103. Dyugovskaya L, Lavie P, Lavie L (2002) Increased adhesion molecules expression and production of reactive oxygen species in leukocytes of sleep apnea patients. *Am J Respir Crit Care Med* 165:859–860
104. Ohga E, Nagase T, Tomita T, Teramoto S, Matsuse T, Katayama H, Ouchi Y (1999) Increased levels of circulating ICAM-1, VCAM-1, and L-selectin in obstructive sleep apnea syndrome. *J Appl Physiol* 87:10–14
105. Dyugovskaya L, Lavie P, Lavie L (2005) Lymphocyte activation as a possible measure of atherosclerotic risk in patients with sleep apnea. *Ann N Y Acad Sci* 1051:340–350
106. Lavie L, Vishnevsky A, Lavie P (2004) Evidence for lipid peroxidation in obstructive sleep apnea. *Sleep* 27:123–128
107. Chen L, Einbinder E, Zhang Q, Hasday J, Balke CW, Scharf SM (2005) Oxidative stress and left ventricular function with chronic intermittent hypoxia in rats. *Am J Respir Crit Care Med* 172:915–920
108. Xu W, Chi L, Row BW, Xu R, Ke Y, Xu B, Luo C, Kheirandish L, Gozal D, Liu R (2004) Increased oxidative stress is associated with chronic intermittent hypoxia-mediated brain cortical neuronal cell apoptosis in a mouse model of sleep apnea. *Neuroscience* 126:313–323
109. Carnevale D, Lembo G (2011) ‘Alzheimer-like’ pathology in a murine model of arterial hypertension. *Biochem Soc Trans* 39:939–944
110. de la Torre JC (2002) Alzheimer disease as a vascular disorder: nosological evidence. *Stroke* 33:1152–1162
111. Johnson KA, Albert MS (2000) Perfusion abnormalities in prodromal AD. *Neurobiol Aging* 21:289–292
112. Kiratli PO, Demir AU, Volkan-Salanci B, Demir B, Sahin A (2010) Cerebral blood flow and cognitive function in obstructive sleep apnea syndrome. *Hell J Nucl Med* 13:138–143
113. Peled N, Abinader EG, Pillar G, Sharif D, Lavie P (1999) Nocturnal ischemic events in patients with obstructive sleep apnea syndrome and ischemic heart disease: effects of continuous positive air pressure treatment. *J Am Coll Cardiol* 34:1744–1749
114. Corfield DR, Meadows GE (2006) Control of cerebral blood flow during sleep and the effects of hypoxia. *Adv Exp Med Biol* 588:65–73
115. Ayalon L, Peterson S (2007) Functional central nervous system imaging in the investigation of obstructive sleep apnea. *Curr Opin Pulmon Med* 13:479–483
116. Macey KE, Macey PM, Woo MA, Henderson LA, Frysinger RC, Harper RK, Alger JR, Yan-Go F, Harper RM (2006) Inspiratory loading elicits aberrant fMRI signal changes in obstructive sleep apnea. *Resp Physiol Neurobiol* 151:44–60
117. Ismail Z, Herrmann N, Francis PL et al (2009) A SPECT study of sleep disturbance and Alzheimer’s disease. *Dementia Geriatr Cogn Disord* 27:254–259
118. Kakeda S, Korogi Y (2010) The efficacy of a voxel-based morphometry on the analysis of imaging in schizophrenia, temporal lobe epilepsy, and Alzheimer’s disease/mild cognitive impairment: a review. *Neuroradiology* 52:711–721
119. Hayakawa T, Terashima M, Kayukawa Y, Ohta T, Okada T (1996) Changes in cerebral oxygenation and hemodynamics during obstructive sleep apneas. *Chest* 109:916–921
120. Engleman HM, Kingshott RN, Martin SE et al (2000) Cognitive function in the sleep apnea/hypopnea syndrome (SAHS). *Sleep* 23(Suppl 4):S102–S108
121. Cohen-Zion M, Stepnowsky C, Marler Shochat T, Kripke DF, Ancoli-Israel S (2001) Changes in cognitive function associated with sleep disordered breathing in older people. *J Am Geriatr Soc* 49:1622–1627
122. Shamsuzzaman AS, Gersh BJ, Somers VK (2003) Obstructive sleep apnea: implications for cardiac and vascular disease. *JAMA* 290:1906–1914
123. Altay T, Gonzales ER, Park TS, Gidday JM (2004) Cerebrovascular inflammation after brief episodic hypoxia: modulation by neuronal and endothelial nitric oxide synthase. *J Appl Physiol* 96:1223–1230
124. Cohen-Zion M, Stepnowsky C, Johnson S, Marler M, Dimsdale JE, Ancoli-Israel S (2004) Cognitive changes and sleep disordered breathing in elderly: differences in race. *J Psychosom Res* 56:549–553
125. Butt M, Dwivedi G, Khair O, Lip GY (2010) Obstructive sleep apnea and cardiovascular disease. *Int J Cardiol* 139:7–16
126. Mou L, Gates A, Mosser VA, Tobin A, Jackson DA (2006) Transient hypoxia induces sequestration of M1 and M2 muscarinic acetylcholine receptors. *J Neurochem* 96:510–519
127. Zimmerman ME, Aloia MS (2006) A review of neuroimaging in obstructive sleep apnea. *J Clin Sleep Med* 2:461–471
128. Quan SF, Wright R (2006) Obstructive sleep apnea-hypopnea and neurocognitive functioning in the Sleep Heart Health Study. *Sleep Med* 7:498–507
129. Gu XQ, Haddad GG (2001) Decreased neuronal excitability in hippocampal neurons of mice exposed to cyclic hypoxia. *J Appl Physiol* 91:1245–1250
130. Alchanatis M, Zias N, Deligiorgis N et al (2008) Comparison of cognitive performance among different age groups in patients with obstructive sleep apnea. *Sleep Breath* 12:17–24
131. Gale SD, Hopkins RO (2004) Effects of hypoxia on the brain: neuroimaging and neuropsychological findings following carbon monoxide poisoning and obstructive sleep apnea. *J Int Neuropsychol Soc* 10:60–71
132. Spira AP, Blackwell T, Stone KL, Redline S, Cauley JA, Ancoli-Israel S, Yaffe K (2008) Sleep-disordered breathing and cognition in older women. *J Am Geriatr Soc* 56:45–50
133. Carlson BW, Neelon VJ, Carlson JR, Hartman M, Bliwise DL (2011) Cerebral oxygenation in wake and during sleep and its relationship to cognitive function in community-dwelling older adults without sleep disordered breathing. *J Gerontol A Biol Sci Med Sci* 66:150–156
134. Walker MP (2009) The role of sleep in cognition and emotion. *Ann N Y Acad Sci* 1156:168–197
135. Bartlett DJ, Rae C, Thompson CH, Byth K, Joffe DA, Enright T, Grunstein RR (2004) Hippocampal area metabolites relate to severity and cognitive function in obstructive sleep apnea. *Sleep Med* 5:593–596
136. Bliwise DL (1996) Is sleep apnea a cause of reversible dementia in old age? *J Am Geriatr Soc* 44:1407–1409

137. Allahdadi KJ, Duling LC, Walker BR, Kanagy NL (2008) Eucapnic intermittent hypoxia augments endothelin-1 vasoconstriction in rats: role of PKC δ . *Am J Physiol Heart Circ Physiol* 294:H920–H927
138. Farré R, Montserrat JM, Navajas D (2008) Morbidity due to obstructive sleep apnea: insights from animal models. *Curr Opin Pulm Med* 14:530–536
139. Bullitt E, Zeng D, Mortamet B et al (2010) The effects of healthy aging on intracerebral blood vessels visualized by magnetic resonance angiography. *Neurobiol Aging* 31:290–300
140. Beebe DW, Groesz L, Wells C, Nichols A, McGee K (2003) The neuropsychological effects of obstructive sleep apnea: a meta-analysis of norm-referenced and case-controlled data. *Sleep* 26:298–307
141. Thomas RJ, Rosen BR, Stern CE, Weiss JW, Kwong KK (2005) Functional imaging of working memory in obstructive sleep-disordered breathing. *J Appl Physiol* 98:2226–2234
142. Ayalon L, Ancoli-Israel S, Aka AA, McKenna BS, Drummond SP (2009) Relationship between obstructive sleep apnea severity and brain activation during a sustained attention task. *Sleep* 32:373–381
143. Kato M, Roberts-Thomson P, Phillips BG et al (2000) Impairment of endothelium-dependent vasodilation of resistance vessels in patients with obstructive sleep apnea. *Circulation* 102:2607–2610
144. Jelic S, Padeletti M, Kawut SM et al (2008) Inflammation, oxidative stress and repair capacity of the vascular endothelium in obstructive sleep apnea. *Circulation* 117:2270–2278
145. Patt BT, Jarjoura D, Haddad DN, Sen CK, Roy S, Flavahan NA, Khayat RN (2010) Endothelial dysfunction in the microcirculation of patients with obstructive sleep apnea. *Am J Respir Crit Care Med* 182:1540–1545
146. Li RC, Row BW, Gozal E, Kheirandish L, Fan Q, Brittan KR, Guo SZ, Sachleben LR Jr, Gozal D (2003) Cyclooxygenase 2 and intermittent hypoxia-induced spatial deficits in the rat. *Am J Respir Crit Care Med* 168:469–475
147. Takase B, Akima T, Uehata A, Ohsuzu F, Kurita A (2004) Effect of chronic stress and sleep deprivation on both flow-mediated dilation in the brachial artery and the intracellular magnesium level in humans. *Clin Cardiol* 27:223–227
148. Shearer WT, Reuben JM, Mullington JM, Price NJ, Lee BN, Smith EO, Szuba MP, Van Dongen HP, Dinges DF (2001) Soluble TNF- α receptor 1 and IL-6 plasma levels in humans subjected to the sleep deprivation model of spaceflight. *J Allergy Clin Immunol* 107:165–170
149. Vgontzas AN, Zoumakis E, Bixler EO, Lin HM, Follett H, Kales A, Chrousos GP (2004) Adverse effects of modest sleep restriction on sleepiness, performance, and inflammatory cytokines. *J Clin Endocrinol Metab* 89:2119–2126
150. von Känel R, Dimsdale JE, Ancoli-Israel S, Mills PJ, Patterson TL, McKibbin CL, Archuleta C, Grant I (2006) Poor sleep is associated with higher plasma proinflammatory cytokine interleukin-6 and procoagulant marker fibrin D-dimer in older caregivers of people with Alzheimer's disease. *J Am Geriatr Soc* 54:431–437
151. Vasquez-Vivar J, Kalyanaraman B, Martasek P, Hogg N, Masters BS, Karoui H, Tordo P, Pritchard KA Jr (1998) Superoxide generation by endothelial nitric oxide synthase: the influence of cofactors. *Proc Natl Acad Sci USA* 95:9220–9225
152. Xia Y, Tsai AL, Berka V, Zweier JL (1998) Superoxide generation from endothelial nitric-oxide synthase. A Ca²⁺/calmodulin-dependent and tetrahydrobiopterin regulatory process. *J Biol Chem* 273:25804–25808
153. Dhar-Masareno M, Cárcamo JM, Golde DW (2005) Hypoxia-reoxygenation-induced mitochondrial damage and apoptosis in human endothelial cells are inhibited by vitamin C. *Free Radic Biol Med* 38:1311–1322
154. Jean A (1991) The nucleus tractus solitarius: neuroanatomic, neurochemical and functional aspects. *Arch Int Physiol Biochim Biophys* 99:A3–A52
155. Andresen MC, Kunze DL (1994) Nucleus tractus solitarius—gateway to neural circulatory control. *Annu Rev Physiol* 56:93–116
156. Seller H, Illert M (1969) The localization of the first synapse in the carotid sinus baroreceptor reflex pathway and its alteration of the afferent input. *Pflugers Arch* 306:1–19
157. Andresen MC, Mendelowitz D (1996) Sensory afferent neurotransmission in caudal nucleus tractus solitarius—common denominators. *Chem Senses* 21:387–395
158. Aylwin ML, Horowitz JM, Bonham AC (1998) Non-NMDA and NMDA receptors in the synaptic pathway between area postrema and nucleus tractus solitarius. *Am J Physiol* 275:H1236–H1246
159. Bajetto A, Bonavia R, Barbero S, Schettini G (2002) Characterization of chemokines and their receptors in the central nervous system: physiopathological implications. *J Neurochem* 82:1311–1329
160. Emch GS, Hermann GE, Rogers RC (2001) TNF- α -induced c-Fos generation in the nucleus of the solitary tract is blocked by NBQX and MK-801. *Am J Physiol Regul Integr Comp Physiol* 281:R1394–R1400
161. Kawanokuchi J, Mizuno T, Takeuchi H, Kato H, Wang J, Mitsuma N, Suzumura A (2006) Production of interferon- γ by microglia. *Mult Scler* 12:558–564
162. Hermann GE, Emch GS, Tovar CA, Rogers RC (2001) c-Fos generation in the dorsal vagal complex after systemic endotoxin is not dependent on the vagus nerve. *Am J Physiol Regul Integr Comp Physiol* 280:R289–R299
163. Dantzer R (2004) Cytokine-induced sickness behaviour: a neuroimmune response to activation of innate immunity. *Eur J Pharmacol* 500:399–411
164. Rozovsky I, Finch CE, Morgan TE (1998) Age-related activation of microglia and astrocytes: in vitro studies show persistent phenotypes of aging, increased proliferation, and resistance to down-regulation. *Neurobiol Aging* 19:97–103
165. Sternberg EM (2006) Neural regulation of innate immunity: a coordinated nonspecific response to pathogens. *Nat Rev Immunol* 6:318–328
166. Tracey KJ (2007) Physiology and immunology of the cholinergic anti-inflammatory pathway. *J Clin Invest* 117:289–296
167. Van der Zanden EP, Boeckxstaens GE, De Jonge WJ (2009) The vagus nerve as a modulator of intestinal inflammation. *Neurogastroenterol Motil* 21:6–17
168. Maier SF, Goehler LE, Fleshner M, Watkins LR (1998) The role of the vagus nerve in cytokine-to-brain communication. *Ann N Y Acad Sci* 840:289–300
169. Groves DA, Brown VJ (2005) Vagal nerve stimulation: a review of its applications and potential mechanisms that mediate its clinical effects. *Neurosci Biobehav Rev* 29:493–500
170. Castle M, Comoli E, Loewy AD (2005) Autonomic brainstem nuclei are linked to the hippocampus. *Neuroscience* 134:657–669
171. Rosas-Ballina M, Tracey KJ (2009) The neurology of the immune system: neural reflexes regulate immunity. *Neuron* 15(64):28–32
172. Rosas-Ballina M, Tracey KJ (2009) Cholinergic control of inflammation. *J Intern Med* 265:663–679
173. Ter Horst GJ, Postema F (1997) Forebrain parasympathetic control of heart activity: retrograde transneuronal viral labeling in rats. *Am J Physiol* 273:H2926–H2930

174. Weber CS, Thayer JF, Rudat M, Wirtz PH, Zimmermann-Viehoff F, Thomas A, Perschel FH, Arck PC, Deter HC (2010) Low vagal tone is associated with impaired post stress recovery of cardiovascular, endocrine, and immune markers. *Eur J Appl Physiol* 109:201–211
175. Espinosa-Oliva AM, de Pablos RM, Villarán RF, Argüelles S, Venero JL, Machado A, Cano J (2011) Stress is critical for LPS-induced activation of microglia and damage in the rat hippocampus. *Neurobiol Aging* 32:85–102
176. Tagliari B, Tagliari AP, Schmitz F, da Cunha AA, Dalmaz C, Wyse AT (2011) Chronic variable stress alters inflammatory and cholinergic parameters in hippocampus of rats. *Neurochem Res* 36:487–493
177. Ricardo JA, Koh TE (1978) Anatomical evidence of direct projection from the nucleus of the solitary tract to the hypothalamus, amygdala, and forebrain structures in the rat. *Brain Res* 153:1–26
178. Schwaber JS, Kapp BS, Higgins GA, Rapp PR (1982) Amygdaloid and basal forebrain direct connections with the nucleus of the solitary tract and the dorsal motor nucleus. *J Neurosci* 2:1424–1438
179. Clayton EC, Williams CL (2000) Adrenergic activation of the nucleus tractus solitarius potentiates amygdala norepinephrine release and enhances retention performance in emotionally arousing and spatial memory tasks. *Behav Brain Res* 112:151–158
180. Barnes DE, Yaffe K, Satariano WA, Tager IB (2003) A longitudinal study of cardiorespiratory fitness and cognitive function in healthy older adults. *J Am Geriatr Soc* 51:459–465
181. Adlard PA, Perreau VM, Pop V, Cotman CW (2005) Voluntary exercise decreases amyloid load in a transgenic model of Alzheimer's disease. *J Neurosci* 25:4217–4221
182. Heyn P, Abreu BC, Ottenbacher KJ (2004) The effects of exercise training on elderly persons with cognitive impairment and dementia: a meta-analysis. *Arch Phys Med Rehabil* 85:1694–1704
183. Black JE, Isaacs KR, Anderson BJ, Alcantara AA, Greenough WT (1990) Learning causes synaptogenesis, whereas motor activity causes angiogenesis, in cerebellar cortex of adult rats. *Proc Natl Acad Sci USA* 87:5568–5572
184. Cotman CW, Berchtold NC (2002) Exercise: a behavioral intervention to enhance brain health and plasticity. *Trends Neurosci* 25:295–301
185. Neeper SA, Gómez-Pinilla F, Choi J, Cotman C (1995) Exercise and brain neurotrophins. *Nature* 373:109
186. van Praag H, Shubert T, Zhao C, Gage FH (2005) Exercise enhances learning and hippocampal neurogenesis in aged mice. *J Neurosci* 25:8680–8685
187. Drager LF, Diegues-Silva L, Diniz PM, Bortolotto LA, Pedrosa RP, Couto RB, Marcondes B, Giorgi DM, Lorenzi-Filho G, Krieger EM (2010) Obstructive sleep apnea, masked hypertension, and arterial stiffness in men. *Am J Hypertens* 23:249–254
188. Miralbell J, Soriano JJ, Spulber G, López-Cancio E, Arenillas JF, Bargallo N, Galán A et al (2011) Structural brain changes and cognition in relation to markers of vascular dysfunction. *Neurobiol Aging* [Epub ahead of print]
189. Sparks DL, Scheff SW, Liu H, Landers TM, Coyne CM, Hunsaker JC 3rd (1995) Increased incidence of neurofibrillary tangles (NFT) in non-demented individuals with hypertension. *J Neurol Sci* 131:162–169
190. Makris TK, Thomopoulos C, Papadopoulos DP, Bratsas A, Papazachou O, Massias S, Michalopoulou E, Tsioufis C, Stefanadis C (2009) Association of passive smoking with masked hypertension in clinically normotensive nonsmokers. *Am J Hypertens* 22:853–859
191. Qiu C, Winblad B, Fratiglioni L (2005) The age-dependent relation of blood pressure to cognitive function and dementia. *Lancet Neurol* 4:487–499
192. Girouard H, Iadecola C (2006) Neurovascular coupling in the normal brain and in hypertension, stroke, and Alzheimer disease. *J Appl Physiol* 100:328–335
193. Mohaupt MG, Schmidli J, Luft FC (2007) Management of uncontrollable hypertension with a carotid sinus stimulation device. *Hypertension* 50:825–828
194. Sloan JA, Illig KA, Bisognano JD (2007) Improved control of resistant hypertension with device-mediated electrical carotid sinus baroreflex stimulation. *J Clin Hypertens* 9:716–719
195. Loewy AD, McKellar S (1980) The neuroanatomical basis of central cardiovascular control. *Fed Proc* 39:2495–2503
196. Ferrario CM, Barnes KL, Bohonek S (1981) Neurogenic hypertension produced by lesions of the nucleus tractus solitarius alone or with sinoaortic denervation in the dog. *Hypertension* 3:112–118
197. Machado BH, Mauad H, Chianca Júnior DA, Haibara AS, Colombari E (1997) Autonomic processing of the cardiovascular reflexes in the nucleus tractus solitarius. *Braz J Med Biol Res* 30:533–543
198. Minson JB, Llewellyn-Smith IJ, Arnold LF, Pilowsky PM, Chalmers JP (1997) C-fos expression in central neurons mediating the arterial baroreceptor reflex. *Clin Exp Hypertens* 19:631–643
199. Paton JF, Deuchars J, Ahmad Z, Wong LF, Murphy D, Kasparov S (2001) Adenoviral vector demonstrates that angiotensin II-induced depression of the cardiac baroreflex is mediated by endothelial nitric oxide synthase in the nucleus tractus solitarius of the rat. *J Physiol* 531:445–458
200. Waki H, Kasparov S, Wong LF, Murphy D, Shimizu T, Paton JF (2003) Chronic inhibition of endothelial nitric oxide synthase activity in nucleus tractus solitarius enhances baroreceptor reflex in conscious rats. *J Physiol* 546:233–242
201. Itoh H, Buñag RD (1992) Catecholaminergic nucleus tractus solitarius lesions in anesthetized rats alter baroreflexes differently with age. *Mech Ageing Dev* 64:69–84
202. Itoh H, Buñag RD (1993) Age-related reduction of reflex bradycardia in conscious rats by catecholaminergic nucleus tractus solitarius lesions. *Mech Ageing Dev* 67:47–63
203. Ebert TJ, Morgan BJ, Barney JA, Denahan T, Smith JJ (1992) Effects of aging on baroreflex regulation of sympathetic activity in humans. *Am J Physiol* 263:H798–H803
204. Hamaad A, Sosin M, Blann AD, Patel J, Lip GY, MacFadyen RJ (2005) Markers of inflammation in acute coronary syndromes: association with increased heart rate and reductions in heart rate variability. *Clin Cardiol* 28:570–576
205. Francis GS (1989) The relationship of the sympathetic nervous system and the renin–angiotensin system in congestive heart failure. *Am Heart J* 118:642
206. Zhang W, Huang BS, Leenen FH (1999) Brain renin–angiotensin system and sympathetic hyperactivity in rats after myocardial infarction. *Am J Physiol* 276:H1608–H1615
207. Teisman AC, Westerink BH, van Veldhuisen DJ, Scholtens E, de Zeeuw D, van Gilst WH (1999) Direct interaction between the sympathetic and renin–angiotensin system in myocardial tissue: a microdialysis study in anaesthetised rats. *J Auton Nerv Syst* 78:117–121
208. Kalsner S, Quillan M (1988) Presynaptic interactions between acetylcholine and adrenergic antagonists on norepinephrine release. *J Pharmacol Exp Ther* 244:879–891
209. Giessler C, Dhein S, Ponicke K, Brodde OE (1999) Muscarinic receptors in the failing human heart. *Eur J Pharmacol* 375:197–202
210. Renaudon B, Bois P, Bescond J, Lenfant J (1997) Acetylcholine modulates I(f) and IK(ACh) via different pathways in rabbit sino-atrial node cells. *J Mol Cell Cardiol* 29:969–975

211. Borovikova LV, Ivanova S, Zhang M, Yang H, Botchkina GI, Watkins LR, Wang H, Abumrad N, Eaton JW, Tracey KJ (2000) Vagus nerve stimulation attenuates the systemic inflammatory response to endotoxin. *Nature* 405:458–462
212. Augustyniak RA, Thomas GD, Victor RG, Zhang W (2005) Nitric oxide pathway as new drug targets for refractory hypertension. *Curr Pharm Des* 11:3307–3315
213. Gupta R, Imai M, Jiang A, Wang M, Sabba H (2006) Chronic therapy with selective electric vagus nerve stimulation normalizes plasma concentration of tissue necrosis factor? Interleukin-6 and B-type natriuretic peptide in dogs with heart failure. *J Am Coll Cardiol* 47:77A
214. Gupta R, Mishra S, Rastogi S, Imai M, Zaca V, Sabba H (2006) Chronic therapy with electric vagus nerve stimulation normalizes mRNA and protein expression of nitric oxide synthase in myocardium of dogs with heart failure. *Eur Heart J* 27:477A
215. Dean JB, Putnam RW (2010) The caudal solitary complex is a site of central CO(2) chemoreception and integration of multiple systems that regulate expired CO(2). *Respir Physiol Neurobiol* 173:274–287
216. Yilmaz MS, Goktalay G, Millington WR, Myer BS, Cutrera RA, Feleder C (2008) Lipopolysaccharide-induced hypotension is mediated by a neural pathway involving the vagus nerve, the nucleus tractus solitarius and alpha-adrenergic receptors in the preoptic anterior hypothalamic area. *J Neuroimmunol* 203:39–49
217. Handforth A, DeGiorgio CM, Schachter SC, Uthman BM, Naritoku DK, Tecoma ES, Henry TR, Collins SD, Vaughn BV et al (1998) Vagus nerve stimulation therapy for partial-onset seizures: a randomized active-control trial. *Neurology* 51:48–55
218. Kralh SE, Clark KB, Smith DC, Browning RA (1998) Locus coeruleus lesions suppress the seizure attenuating effects of vagus nerve stimulation. *Epilepsia* 39:709–714
219. Naritoku DK, Terry WJ, Helfert RH (1995) Regional induction of Fos immunoreactivity in the brain by anticonvulsant stimulation of the vagus nerve. *Epilepsy Res* 22:53–62
220. Kessler W, Traeger T, Westerholt A, Neher F, Mikulcak M, Müller A, Maier S, Heidecke CD (2006) The vagal nerve as a link between the nervous and immune system in the instance of polymicrobial sepsis. *Langenbecks Arch Surg* 391:83–87
221. Ulloa L (2005) The vagus nerve and the nicotinic anti-inflammatory pathway. *Nat Rev Drug Discov* 4:673–684
222. Boon P, Moors I, De Herdt V, Vonck K (2006) Vagus nerve stimulation and cognition. *Seizure* 15:259–263
223. Helmstaedter C, Hoppe C, Elger CE (2001) Memory alterations during acute high-intensity vagus nerve stimulation. *Epilepsy Res* 47:37–42
224. Ghacibeh GA, Shenker JJ, Shenal B, Uthman BM, Heilman KM (2006) The influence of vagus nerve stimulation on memory. *Cogn Behav Neurol* 19:119–122
225. Ghacibeh GA, Shenker JJ, Shenal B, Uthman BM, Heilman KM (2006) Effect of vagus nerve stimulation on creativity and cognitive flexibility. *Epilepsy Behav* 8:720–725
226. Hartig W, Bauer A, Brauer K, Grosche J, Hortobagyi T, Penke B, Schliebs R, Harkany T (2002) Functional recovery of cholinergic basal forebrain neurons under disease conditions: old problems, new solutions? *Rev Neurosci* 13:95–165
227. Schliebs R, Arendt T (2011) The cholinergic system in aging and neuronal degeneration. *Behav Brain Res* 221:555–563
228. Nyakas C, Granic I, Halmy LG, Banerjee P, Luiten PG (2011) The basal forebrain cholinergic system in aging and dementia. Rescuing cholinergic neurons from neurotoxic amyloid- β 42 with memantine. *Behav Brain Res* 221:594–603
229. Mesulam M (2004) The cholinergic system of Alzheimer's Disease: pivotal factor or side show? *Learning & Memory* 11:43–49
230. Grudzien A, Shaw P, Weintraub S, Bigio E, Mash DC, Mesulam MM (2007) Locus coeruleus neurofibrillary degeneration in aging, mild cognitive impairment and early Alzheimer's disease. *Neurobiol Aging* 28:327–335
231. Counts SE, Mufson EJ (2010) Noradrenaline activation of neurotrophic pathways protects against neuronal amyloid toxicity. *J Neurochem* 113:649–660
232. Insua D, Suárez ML, Santamarina G, Sarasa M, Pesini P (2010) Dogs with canine counterpart of Alzheimer's disease lose noradrenergic neurons. *Neurobiol Aging* 31:625–635
233. Foote SL, Bloom FE, Aston-Jones G (1983) Nucleus locus ceruleus: new evidence of anatomical and physiological specificity. *Physiol Rev* 63:844–914
234. Heneka MT, Galea E, Gavrilyuk V, Dumitrescu-Ozimek L, Daeschner J, O'Banion MK, Weinberg G, Klockgether T, Feinstein DL (2002) Noradrenergic depletion potentiates beta-amyloid-induced cortical inflammation: implications for Alzheimer's disease. *J Neurosci* 22:2434–2442
235. Heneka MT, Ramanathan M, Jacobs AH, Dumitrescu-Ozimek L, Bilkei-Gorzo A, Debeir T, Sastre M et al (2006) Locus ceruleus degeneration promotes Alzheimer pathogenesis in amyloid precursor protein 23 transgenic mice. *J Neurosci* 26:1343–1354
236. Heneka MT, Nadrigny F, Regen T, Martinez-Hernandez A, Dumitrescu-Ozimek L, Terwel D (2010) Locus ceruleus controls Alzheimer's disease pathology by modulating microglial functions through norepinephrine. *Proc Natl Acad Sci USA* 107:6058–6063
237. Kalinin S, Gavrilyuk V, Polak PE, Vasser R, Zhao J, Heneka MT, Feinstein DL (2007) Noradrenaline deficiency in brain increases beta-amyloid plaque burden in an animal model of Alzheimer's disease. *Neurobiol Aging* 28:1206–1214
238. Madrigal JL, Kalinin S, Richardson JC, Feinstein DL (2007) Neuroprotective actions of noradrenaline: effects on glutathione synthesis and activation of peroxisome proliferator activated receptor delta. *J Neurochem* 103:2092–2101
239. Traver S, Salthun-Lassalle B, Marien M, Hirsch EC, Colpaert F, Michel PP (2005) The neurotransmitter noradrenaline rescues septal cholinergic neurons in culture from degeneration caused by low-level oxidative stress. *Mol Pharmacol* 67:1882–1891
240. Zarow C, Lyness SA, Mortimer JA, Chui HC (2003) Neuronal loss is greater in the locus coeruleus than nucleus basalis and substantia nigra in Alzheimer and Parkinson diseases. *Arch Neurol* 60:337–341
241. Gonzalo-Ruiz A, González I, Sanz-Anquela JM (2003) Effects of beta-amyloid protein on serotonergic, noradrenergic, and cholinergic markers in neurons of the pontomesencephalic tegmentum in the rat. *J Chem Neuroanat* 26:153–169
242. Ohm TG, Busch C, Bohl J (1997) Unbiased estimation of neuronal numbers in the human nucleus coeruleus during aging. *Neurobiol Aging* 18:393–399
243. Goldman G, Coleman PD (1981) Neuron numbers in locus coeruleus do not change with age in Fisher 344 rat. *Neurobiol Aging* 2:33–36
244. Fujiwara N, Higashi H, Shimoji K, Yoshimura M (1987) Effects of hypoxia on rat hippocampal neurones in vitro. *J Physiol* 384:131–151
245. Shimoji K, Higashi H, Fujiwara N, Fukuda S, Yoshimura M (1989) Hypoxia in brain slices. *Biomed Biochim Acta* 48:S149–S154
246. Nieber K, Sevcik J, Illes P (1995) Hypoxic changes in rat locus coeruleus neurons in vitro. *J Physiol* 486:33–46
247. Yang JJ, Chou YC, Lin MT, Chiu TH (1997) Hypoxia-induced differential electrophysiological changes in rat locus coeruleus neurons. *Life Sci* 61:1763–1773
248. Ma S, Mifflin SW, Cunningham JT, Morilak DA (2008) Chronic intermittent hypoxia sensitizes acute hypothalamic-pituitary-adrenal stress reactivity and Fos induction in the rat locus coeruleus in response to subsequent immobilization stress. *Neuroscience* 154:1639–1647

249. Zhu Y, Fenik P, Zhan G, Mazza E, Kelz M, Aston-Jones G, Veasey SC (2007) Selective loss of catecholaminergic wake active neurons in a murine sleep apnea model. *J Neurosci* 27: 10060–10071
250. Manta S, Dong J, Debonnel G, Blier P (2009) Enhancement of the function of rat serotonin and norepinephrine neurons by sustained vagus nerve stimulation. *J Psychiatry Neurosci* 34: 272–280
251. Insausti R, Amaral DG, Cowan WM (1987) The entorhinal cortex of the monkey: II. Cortical afferents. *J Comp Neurol* 264: 356–395
252. Liu P, Bilkey DK (1997) Parallel involvement of perirhinal and lateral entorhinal cortex in the polysynaptic activation of hippocampus by olfactory inputs. *Hippocampus* 7:296–306
253. Mouly A-M, Litaudon P, Chabaud P, Ravel N, Gervais R (1998) Spatiotemporal distribution of a late synchronized activity in olfactory pathways following stimulation of the olfactory bulb in rats. *Eur J Neurosci* 10:1128–1135
254. Chapman CA, Racine RJ (1997) Piriform cortex efferents to the entorhinal cortex in vivo: kindling-induced potentiation and the enhancement of long-term potentiation by low-frequency piriform cortex or medial septal stimulation. *Hippocampus* 7:257–270
255. Th VanGroen, Lopes da Silva FH, Wadman WJ (1987) Synaptic organization of olfactory inputs and local circuits in the entorhinal cortex, a current source density analysis in the cat. *Exp Brain Res* 67:615–622
256. Witter MP, Amaral DG (1991) Entorhinal cortex of the monkey: V. Projections to the dentate gyrus, hippocampus, and subicular complex. *J Comp Neurol* 307:437–459
257. Zola-Morgan S, Squire LR, Amaral DG, Suzuki WA (1989) Lesions of perirhinal and parahippocampal cortex that spare the amygdala and hippocampal formation produce severe memory impairment. *J Neurosci* 9:4355–4370
258. Du AT, Schuff N, Amend D, Laakso MP, Hsu YY et al (2001) Magnetic resonance imaging of the entorhinal cortex and hippocampus in mild cognitive impairment and Alzheimer's disease. *J Neurol Neurosurg Psychiatry* 71:441–447
259. Gómez-Isla T, Price JL, McKeel DW Jr, Morris JC, Growdon JH, Hyman BT (1996) Profound loss of layer II entorhinal cortex neurons occurs in very mild Alzheimer's disease. *J Neurosci* 16:4491–4500
260. Pennanen C, Kivipelto M, Tuomainen S, Hartikainen P, Hanninen T, Laakso MP, Hallikainen M et al (2004) Hippocampus and entorhinal cortex in mild cognitive impairment and early AD. *Neurobiol Aging* 25:303–310
261. Christen-Zaech S, Kraftsik R, Pillevuit O, Kiraly M, Martins R, Khalili K, Miklossy J (2003) Early olfactory involvement in Alzheimer's disease. *Can J Neurol Sci* 30:20–25
262. Wilson RS, Arnold SE, Schneider JA, Tang Y, Bennett DA (2007) The relationship between cerebral Alzheimer's disease pathology and odour identification in old age. *J Neurol Neurosurg Psychiatry* 78:30–35
263. Velayudhan L, Lovestone S (2009) Smell identification test as a treatment response marker in patients with Alzheimer disease receiving donepezil. *J Clin Psychopharmacol* 29:387–390
264. Ruggiero DA, Anwar S, Kim J, Glickstein SB (1998) Visceral afferent pathways to the thalamus and olfactory tubercle: behavioral implications. *Brain Res* 799:159–171
265. Guevara-Guzman R, Garcia-Diaz DE, Solano-Flores LP, Wayner MJ, Armstrong DL (1991) Role of the paraventricular nucleus in the projection from the nucleus of the solitary tract to the olfactory bulb. *Brain Res Bull* 27:447–450
266. García-Díaz DE, Aguilar-Baturoni HU, Guevara-Aguilar R, Wayner MJ (1984) Vagus nerve stimulation modifies the electrical activity of the olfactory bulb. *Brain Res Bull* 12:529–537
267. de Toledo-Morrell L, Goncharova I, Dickerson B, Wilson RS, Bennett DA (2000) From healthy aging to early Alzheimer's disease: in vivo detection of entorhinal cortex atrophy. *Ann N Y Acad Sci* 911:240–253
268. Kerfoot EC, Chattillion EA, Williams CL (2008) Functional interactions between the nucleus tractus solitarius (NTS) and nucleus accumbens shell in modulating memory for arousing experiences. *Neurobiol Learn Mem* 89:47–60
269. Danielsen EH, Magnuson DJ, Gray TS (1989) The central amygdaloid nucleus innervation of the dorsal vagal complex in rat: a *Phaseolus vulgaris* leucoagglutinin lectin anterograde tracing study. *Brain Res Bull* 22:705–715
270. Saha S, Batten TF, Henderson Z (2000) A GABAergic projection from the central nucleus of the amygdala to the nucleus of the solitary tract: a combined anterograde tracing and electron microscopic immunohistochemical study. *Neuroscience* 99:613–626
271. Saha S, Henderson Z, Batten TF (2002) Somatostatin immunoreactivity in axon terminals in rat nucleus tractus solitarius arising from central nucleus of amygdala: coexistence with GABA and postsynaptic expression of sst2A receptor. *J Chem Neuroanat* 24:1–13
272. Zhang X, Cui J, Tan Z, Jiang C, Fogel R (2003) The central nucleus of the amygdala modulates gut-related neurons in the dorsal vagal complex in rats. *J Physiol* 553:1005–1018
273. Rinaman L (2010) Ascending projections from the caudal visceral nucleus of the solitary tract to brain regions involved in food intake and energy expenditure. *Brain Res* 1350:18–34
274. McGeer PL, Itagaki S, Boyes BE, McGeer EG (1988) Reactive microglia are positive for HLA-DR in the substantia nigra of Parkinson's and Alzheimer's disease brains. *Neurology* 38:1285–1291
275. Tansey MG, McCoy MK, Frank-Cannon TC (2007) Neuroinflammatory mechanisms in Parkinson's disease: potential environmental triggers, pathways, and targets for early therapeutic intervention. *Exp Neurol* 208:1–25
276. Rogers J, Mastroeni D, Leonard B, Joyce J, Grover A (2007) Neuroinflammation in Alzheimer's disease and Parkinson's disease: are microglia pathogenic in either disorder? *Int Rev Neurobiol* 82:235–246
277. Tuppo EE, Arias HR (2005) The role of inflammation in Alzheimer's disease. *Int J Biochem Cell Biol* 37:289–305
278. Engelhart MJ, Geerlings MI, Meijer J, Kiliaan A, Ruitenberg A, van Swieten JC, Stijnen T, Hofman A, Witteman JC (2004) Inflammatory proteins in plasma and the risk of dementia: the Rotterdam Study. *Arch Neurol* 61:668–672
279. Celler S, Peyron R, Faillenot I, Pichot V, Alabdullah M, Gaspoz JM, Laurent B, Barthélémy JC, Roche F (2009) Undiagnosed sleep-related breathing disorders are associated with focal brainstem atrophy in the elderly. *Hum Brain Mapp* 30:2090–2097
280. Daulatzai M (2010) Early stages of pathogenesis in memory impairment during normal senescence and Alzheimer's disease. *J Alzheimers Dis* 20:355–367
281. Daulatzai M (2011) Role of sensory stimulation in amelioration of obstructive sleep apnea. *Sleep Disorders* 2011:596879
282. Waki H, Bhuiyan ME, Gouraud SS, Takagishi M, Hatada A, Kohsaka A, Paton JF, Maeda M (2011) Acute reductions in blood flow restricted to the dorsomedial medulla induce a pressor response in rats. *J Hypertens* 29:1536–1545
283. Stanimirovic D, Shapiro A, Wong J, Hutchison J, Durkin JP (1997) The induction of ICAM-1 in human cerebrovascular endothelial cells (HCEC) by ischemia-like conditions promotes enhanced neutrophil/HCEC adhesion. *J Neuroimmunol* 76:193–205
284. Mojsilovic-Petrovic J, Callaghan D, Cui H, Dean C, Stanimirovic DB, Zhang W (2007) Hypoxia-inducible factor-1 (HIF-1) is involved in the regulation of hypoxia-stimulated expression of

- monocyte chemoattractant protein-1 (MCP-1/CCL2) and MCP-5 (Ccl12) in astrocytes. *J Neuroinflammation* 4:12
285. Zhang W, Mojsilovic-Petrovic J, Callaghan D, Jones A, Cui H, Howlett C, Stanimirovic D (2006) Evidence that hypoxia-inducible factor-1 (HIF-1) mediates transcriptional activation of Interleukin-1 β (IL-1 β) in astrocyte cultures. *J Neuroimmunol* 174:63–73
 286. Csiszar A, Wang M, Lakatta EG, Ungvari Z (2008) Inflammation and endothelial dysfunction during aging: role of NF-kappaB. *J Appl Physiol* 105:1333–1341
 287. Ohi Y, Ishii Y, Sasahara M, Haji A (2010) Involvement of platelet-derived growth factor-BB and its receptor-beta in hypoxia-induced depression of excitatory synaptic transmission in the nucleus tractus solitarius of mice. *J Pharmacol Sci* 112:477–481
 288. Macey PM, Macey KE, Henderson LA, Alger JR, Frysinger RC, Woo MA, Yan-Go F, Harper RM (2003) Functional magnetic resonance imaging responses to expiratory loading in obstructive sleep apnea. *Respir Physiol Neurobiol* 138:275–290
 289. Nair D, Dayyat EA, Zhang SX, Wang Y, Gozal D (2011) Intermittent hypoxia-induced cognitive deficits are mediated by NADPH oxidase activity in a murine model of sleep apnea. *PLoS One* 6:e19847
 290. Kim SJ, Lee JH, Lee DY, Jhoo JH, Woo JI (2011) Neurocognitive dysfunction associated with sleep quality and sleep apnea in patients with mild cognitive impairment. *Am J Geriatr Psychiatry* 19:374–381
 291. Zhu Y, Fenik P, Zhan G, Sanfillipo-Cohn B, Naidoo N, Veasey SC (2008) Eif-2a protects brainstem motoneurons in a murine model of sleep apnea. *J Neurosci* 28:2168–2178
 292. Kaur C, Viswanathan S, Ling EA (2011) Hypoxia-induced cellular and vascular changes in the nucleus tractus solitarius and ventrolateral medulla. *J Neuropathol Exp Neurol* 70:201–217
 293. Steiropoulos P, Papanas N, Nena E, Antoniadou M, Serasli E, Papoti S, Hatzizisi O, Kyriazis G, Tzouveleakis A, Maltezos E, Tsara V, Bouros D (2010) Inflammatory markers in middle-aged obese subjects: does obstructive sleep apnea syndrome play a role? *Mediat Inflamm* 2010:675320
 294. Porzionato A, Macchi V, Guidolin D, Sarasin G, Parenti A, De Caro R (2008) Anatomic distribution of apoptosis in medulla oblongata of infants and adults. *J Anat* 212:106–113
 295. Stecco C, Porzionato A, Macchi V, Sarasin G, Calcagno A, Parenti A, De Caro R (2005) Detection of apoptosis in human brainstem by TUNEL assay. *Ital J Anat Embryol* 110:255–260
 296. Batti L, O'Connor JJ (2010) Tumor necrosis factor-alpha impairs the recovery of synaptic transmission from hypoxia in rat hippocampal slices. *J Neuroimmunol* 218:21–27
 297. Zhang JH, Fung SJ, Xi M, Sampogna S, Chase MH (2010) Apnea produces neuronal degeneration in the pons and medulla of guinea pigs. *Neurobiol Dis* 40:251–264
 298. Jamieson GA, Maitland NJ, Itzhaki RF (1992) Herpes simplex virus type 1 DNA sequences are present in aged normal and Alzheimer's disease brain but absent in lymphocytes. *Arch Gerontol Geriatr* 15:197–201
 299. Wozniak MA, Itzhaki RF (2010) Antiviral agents in Alzheimer's disease: hope for the future? *Ther Adv Neurol Disord* 3:141–152
 300. Fay RA, Norgren R (1997) Identification of rat brainstem multisynaptic connections to the oral motor nuclei using pseudorabies virus. III. Lingual muscle motor systems. *Brain Res Rev* 25:291–311
 301. Williams AE, Blakemore WF (1990) Monocyte-mediated entry of pathogens into the central nervous system. *Neuropathol Appl Neurobiol* 16:377–392
 302. Leyva-Grado VH, Churchill L, Wu M, Williams TJ, Taishi P, Majde JA, Krueger JM (2009) Influenza virus- and cytokine-immunoreactive cells in the murine olfactory and central autonomic nervous systems before and after illness onset. *J Neuroimmunol* 211:73–83
 303. Shinya K, Shimada A, Ito T, Otsuki K, Morita T, Tanaka H, Takada A, Kida H, Umemura T (2000) Avian influenza virus intranasally inoculated infects the central nervous system of mice through the general visceral afferent nerve. *Arch Virol* 145:187–195
 304. Matsuda K, Park CH, Sunden Y, Kimura T, Ochiai K, Kida H, Umemura T (2004) The vagus nerve is one route of transneuronal invasion for intranasally inoculated influenza a virus in mice. *Vet Pathol* 41:101–107
 305. McBride PA, Schulz-Schaeffer WJ, Donaldson M, Bruce M, Diring H, Kretschmar HA, Beekes M (2001) Early spread of scrapie from the gastrointestinal tract to the central nervous system involves autonomic fibers of the splanchnic and vagus nerves. *J Virol* 75:9320–9327
 306. Hansen MK, Taishi P, Chen Z, Krueger JM (1998) Vagotomy blocks the induction of interleukin-1beta (IL-1beta) mRNA in the brain of rats in response to systemic IL-1beta. *J Neurosci* 18:2247–2253
 307. Churchill L, Taishi P, Wang M, Brandt J, Cearley C, Rehman A, Krueger JM (2006) Brain distribution of cytokine mRNA induced by systemic administration of interleukin-1beta or tumor necrosis factor alpha. *Brain Res* 1120:64–73
 308. Doba N, Reis DJ (1973) Acute fulminating neurogenic hypertension produced by brainstem lesions in the rat. *Circ Res* 32:584–593
 309. Talman WT, Perrone MH, Reis DJ (1981) Acute hypertension after the local injection of kainic acid into the nucleus tractus solitarii of rats. *Circ Res* 48:292–298
 310. Waki H, Gouraud SS, Maeda M, Paton JF (2008) Specific inflammatory condition in nucleus tractus solitarii of the SHR: novel insight for neurogenic hypertension? *Auton Neurosci* 142:25–31
 311. Waki H, Gouraud SS, Maeda M, Paton JF (2008) Gene expression profiles of major cytokines in the nucleus tractus solitarii of the spontaneously hypertensive rat. *Auton Neurosci* 142:40–44
 312. Waki H, Gouraud SS, Maeda M, Paton JF (2010) Evidence of specific inflammatory condition in nucleus tractus solitarii of spontaneously hypertensive rats. *Exp Physiol* 95:595–600
 313. Waki H, Gouraud SS, Maeda M, Raizada MK, Paton JF (2011) Contributions of vascular inflammation in the brainstem for neurogenic hypertension. *Respir Physiol Neurobiol* 178:422–428
 314. Paton JF, Waki H (2009) Is neurogenic hypertension related to vascular inflammation of the brainstem? *Neurosci Biobehav Rev* 33:89–94
 315. Johnston H, Boutin H, Allan SM (2011) Assessing the contribution of inflammation in models of Alzheimer's disease. *Biochem Soc Trans* 39:886–890
 316. Mintun MA, Larossa GN, Sheline YI, Dence CS, Lee SY, Mach RH, Klunk WE, Mathis CA, DeKosky ST, Morris JC (2006) [11C]PIB in a nondemented population: potential antecedent marker of Alzheimer disease. *Neurology* 67:446–452
 317. Soscia SJ, Kirby JE, Washicosky KJ et al (2010) The Alzheimer's disease-associated amyloid β -protein is an antimicrobial peptide. *PLoS One* 5:e9505
 318. Kobayashi N, Masuda J, Kudoh J, Shimizu N, Yoshida T (2008) Binding sites on tau proteins as components for antimicrobial peptides. *Biocontrol Sci* 13:49–56
 319. Lee DC, Rizer J, Selenica ML, Reid P, Kraft C, Johnson A, Blair L, Gordon MN, Dickey CA, Morgan D (2010) LPS- induced inflammation exacerbates phospho-tau pathology in rTg4510 mice. *J Neuroinflamm* 7:56

320. Dickey C, Kraft C, Jinwal U, Koren J, Johnson A, Anderson L, Lebson L, Lee D, Dickson D, de Silva R, Binder LI, Morgan D, Lewis J (2009) Aging analysis reveals slowed tau turnover and enhanced stress response in a mouse model of tauopathy. *Am J Pathol* 174:228–238
321. Quinn J, Montine T, Morrow J, Woodward WR, Kulhanek D, Eckenstein F (2003) Inflammation and cerebral amyloidosis are disconnected in an animal model of Alzheimer's disease. *J Neuroimmunol* 137:32–41
322. Kitazawa M, Oddo S, Yamasaki TR, Green KN, LaFerla FM (2005) Lipopolysaccharide-induced inflammation exacerbates tau pathology by a cyclin-dependent kinase 5-mediated pathway in a transgenic model of Alzheimer's disease. *J Neurosci* 25: 8843–8853
323. Li Y, Liu L, Barger SW, Griffin WS (2003) Interleukin-1 mediates pathological effects of microglia on tau phosphorylation and on synaptophysin synthesis in cortical neurons through a p38-MAPK pathway. *J Neurosci* 23:1605–1611
324. Janelins MC, Mastrangelo MA, Park KM, Sudol KL, Narrow WC, Oddo S, LaFerla FM, Callahan LM, Federoff HJ, Bowers WJ (2008) Chronic neuron-specific tumor necrosis factor- α expression enhances the local inflammatory environment ultimately leading to neuronal death in 3xTg-AD mice. *Am J Pathol* 173:1768–1782
325. Rosi S, Ramirez-Amaya V, Vazdarjanova A, Worley PF, Barnes CA, Wenk GL (2005) Neuroinflammation alters the hippocampal pattern of behaviorally induced Arc expression. *J Neurosci* 25:723–731
326. El Solh AA, Akinnusi ME, Baddoura FH, Mankowski CR (2007) Endothelial cell apoptosis in obstructive sleep apnea. A link to endothelial dysfunction. *Am J Respir Crit Care Med* 175: 1186–1191