


Experimental Data types include:


Sequences - Typically <u>Next-Generation DNA Sequencing (NGS)</u>.


3D Protein Structures - X-ray crystallography or

Nuclear magnetic resonance spectroscopy (NMR)

Gene Expression Data - Microarrays









First, find a family of **Homologous** sequences.

 APFELVISWKLIVESPAINCDWRTENGLANDSGMLVNOWPAI
 APYELVISQWKLIVESNPAINKDWRTYENGLANDSGMLVNOWAI
 APFELVISWKLIVESNPAINCDWRTENGLANDSGMLVNOWAI
 APFELVISQWKLIVESNPAINCDWRTENGLANDSGMLVNOWAI
 APYELVISWKLIVESNPINCDWRTENGLANDRSGMLINOWAI
 APFELVISQWKLIVESNPAINCDWRTENGLANDSGMLVNOWLI
 APFELVISQWKLIVESNPAINDWRTENGLANDSGMLVNOWAI
 APYELVISWKLIVESNPAINCDWRTENGLANDSGMLLNOWMI




Then, align by inserting "-"s representing <u>InDels</u>, in each sequence.

 APFELVIS-WKLIVES-PAINCDWRT-ENGLANDSGMLV-NOWPAI	
 APYELVISQWKLIVESNPAINKDWRTYENGLANDSGMLV-NOW-AI	
 APFELVIS-WRLIVESNPAINCDWRT-ENGLANDSGMLV-NOW-AI	
 APFELVISQWKLIVESNPAINCDWRT-ENGLANDSGMLV-NOW-AI	
 APYELVIS-WKLIVESNP-INCDWRT-ENGLANDRSGMLINOW-AI	
 APFELVISQWKLIVESNPAINCDWRT-ENGLANDSGMLV-NOW-LI	
 APFELVISQWKLIVESNPAIN-DWRT-ENGLANDSGMLV-NOW-AI	
 APYELVIS-WKLIVESNPAINCDWRT-ENGLANDSGMLL-NOW-MI	




Next, identify the columns where Substitutions and/or InDels have been predicted.

 APFELVIS-WKLIVES-PAINCDWRT-ENGLANDSGMLV-NOWPAI	
 APYELVISQWKLIVESNPAINKDWRTYENGLANDSGMLV-NOW-AI	
 APFELVIS-WRLIVESNPAINCDWRT-ENGLANDSGMLV-NOW-AI	
 APFELVISQWKLIVESNPAINCDWRT-ENGLANDSGMLV-NOW-AI	
 APYELVIS-WKLIVESNP-INCDWRT-ENGLANDRSGMLINOW-AI	
 APFELVISQWKLIVESNPAINCDWRT-ENGLANDSGMLV-NOW-LI	
 APFELVISQWKLIVESNPAIN-DWRT-ENGLANDSGMLV-NOW-AI	
 APYELVIS-WKLIVESNPAINCDWRT-ENGLANDSGMLL-NOW-MI	




Then, identify the columns where full Conservation has been predicted.

```
... APFELVIS-WKLIVES-PAINCDWRT-ENGLANDSGMLV-NOWPAI ...
APYELVISQWKLIVESNPAINKDWRTYENGLANDSGMLV-NOW-AI ...
APFELVIS-WRLIVESNPAINCDWRT-ENGLANDSGMLV-NOW-AI ...
APFELVISQWKLIVESNPAINCDWRT-ENGLANDSGMLV-NOW-AI ...
APYELVIS-WKLIVESNP-INCDWRT-ENGLANDRSGMLINOW-AI ...
APFELVISQWKLIVESNPAINCDWRT-ENGLANDSGMLV-NOW-LI ...
APFELVISQWKLIVESNPAINCDWRT-ENGLANDSGMLV-NOW-AI ...
APFELVISQWKLIVESNPAIN-DWRT-ENGLANDSGMLV-NOW-AI ...
APYELVIS-WKLIVESNPAINCDWRT-ENGLANDSGMLV-NOW-MI ...
```



Finally ... Identify the **Glorious Message**!!!!.

```
-PAINCDWRT-
                                     SGMLV-
                                              PAI
                 NPAINKDWRTY
                                     SGMLV-
                                              -AI
      ISOWKI
       IS-WRL
                 NPAINCDWRT-
                                     SGMLV-
                                              -AI
        OWKI
                 NPA NCDWRT-
                                     SGMLV-
                                               -AI
APF
                                     RSGMLI
                 NP-INCDWRT-
                                               -AI
                 NPAINCDWRT-
                                     SGMLV-
      ISOWKLI
                                              LI
APF
                 NPAIN-DWRT-
                                     SGMLV-
      /ISOWKL
                                               HAI
APF
APYELVIS-WKL
                 NPAINCDWRT-
                                     SGMLL-
                                              MT
```




Database searching is the most common Bioinformatics process by far.

Database searching is pairwise comparison repeated many times.

Non-optimal comparison methods are essential for practical reasons.

A list of matches, ordered by the improbability of occurring just by chance is generated.




Searching for Homologous Sequences in a Sequence Database.

Database searching seeks "Similarity". Users seek "Homology".

Query	KLYF	RPQTPE	RPQTPEPPPPPPPPPPP					LPAAPPQP	
Similarity	+L P		+P	P	P	PP	P	PP	PP+P
Database Entry	RLTPPQPLMMPPRPTPPTPLPPATLTVPPRP								

Homology?


Or 2 proteins including a lot of Prolines??



Searching for Homologous Sequences in a Sequence Database.

Database searching seeks "Similarity". Users seek "Homology".






Searching for Homologous Sequences in a Sequence Database.

Database searching seeks "Similarity". Users seek "Homology".



Homology?


A very unconvincing alignment!!



Database searching seeks "Similarity". Users seek "Homology".

TTAGCAAGATCAGCCCTAACTCGGCATCTT											
Query	L	A	R	S	С	L	T	R	H	L	Homology?
Similarity	L	A	R	S	C	L	T	R	H	L	
Database Entry	L	A	R	S	С	L	T	R	Н	L	Probable a perfect
CTTGCGCGCTCTGTCTTGACGAGACACTTA									protein match??		

In all circumstances - always align at the protein level wherever possible.



Largely a matter of finding short sequences within longer ones.

Computationally trivial.

A concrete example is required:

Restriction Mapping

Detecting Restriction Enzyme Recognition Sites is complicated by their redundancy.

Few Recognition Sites can be simply defined using only the codes A, C, G and T.

The solution is to use the <u>Nucleotide</u> <u>Ambiguity Codes</u> defined by <u>IUPAC</u>.



Unambiguous site (EcoRI):

G/AATC

Ambiguous site (PpuMI):

RG/GWCCY—

Cut here


And here

TTAGCAAGATCAGGACCTACTCGGCATCTTCCTGGGTCCC

RGGWCCY

IUPAC DNA Alphabet

Code	<u> </u>	Meaning				
A		A				
С		С				
G		G				
T/U		T/U				
M `a	a M ino`	A C				
R `]	pu R ine`	A G				
7 W	₩eak`	$\mathbf{A} \mid \mathbf{T}$				
s `:	Strong`	C∣G				
Y `;	p Y rimidine`	C T				
K `1	Keto`	G T				
v `1	not T`	A C G				
H `1	not G`	$\mathbf{A} \mid \mathbf{C} \mid \mathbf{T}$				
D `1	not C`	$\mathbf{A} \mid \mathbf{G} \mid \mathbf{T}$				
B `1	not A`	C G T				
N `a	a N y`	A C G T				



Patterns can be derived manually to represent conserved regions of MSAs

Simple where conservation is 100%

CQVLNPYYHWGQCGGIGWSGPTVCASGTT ...

. CQYSNDYYHWGQCGGIGWSGCKTCTSGTT ...

. CHVLNPYYQWGQCGGIGWTPSTTCASPYT ...

. CSTLNPYYVWGQCGGIGWSGPTNCAPGSA ..

.. CVYSNDYYVWGQCGGIGWSGPTCCASGST ...

WGQCGGIGW

Pattern

Data Systems Programming Statistics Data Generation Data Analysis Searching for simple sequence patterns Sequences in Proteins.

Not so easy where conservation is less than perfect

An Amino Acid Alphabet including all ambiguities is not practical!

The solution is a <u>simple syntax for</u> <u>ambiguous amino acid sequences</u>.

... CQVLNPYYHWKQCGGLGWSGPTVCASGTT ...

... CQYSNDYYHWGQCPGIGWSGCKTCTSGTT ...


... CHVLNPYYQWAQCFGVGWTPSTTCASPYT ...

... CSTLNPYYVWLQCYGIGWSGPTNCAPGSA ...


... CVYSNDYYVWAQCGGVGWSGPTCCASGST ...

W{P}QCxG[LIV]GW Pattern

NOT a P Anything Lorlor V












Only highly conserved regions can be described usefully.

Patterns cannot weight possibilities by frequency.










Patterns do not reflect commonly accepted substitutions.




Again, start with an MSA of instances of the feature to be modelled.

Create a "suitable" representation of the relevant portion of MSA

Compare the model along other protein sequences was illustrated for simple patterns.

Where matches are detected, the corresponding protein property is likely to occur.




A variety of simple models have been developed (e.g. <u>Position Weight Matrices</u>) for a number of purposes, including:

- Gene discovery in bacteria genomes (DNA)
- Early versions of 2D protein Structure Prediction
- Transmembrane Alpha Helix prediction

- TATA box Detection (DNA)
- Helix-Turn-Helix (HTH) Prediction
- Prediction of Coiled Coils

The most powerful and prolific current profiles are Hidden Markov Models (HMMs)




Broadly, the estimation of evolutionary history from available evidence.

"Evidence" does not <u>have</u> to be a carefully crafted MSA of Orthologous sequences from a range of organisms.


However, in the context of Bioinformatics, it invariably is.

Typically, conclusions of Phylogenetic analysis are represented as **Evolutionary Trees**.

Which are very Beautiful!!



My personal preference is for trees that place <u>ME</u> as far away from a <u>MOUSE</u> as possible!!!!




Phylogeny is another example of an analysis based on MSAs.

One very effective Phylogenetic strategy is to seek an answer to the question:

"What is the most probable Evolutionary Tree, given I believe this MSA to be perfect?"


Reinforcing how central is the role of Statistics in Bioinformatics.



Essentially predicting the locations of Alpha Helices, Beta Sheets and Turns.

Modern methods employ Machine Learning to generate Artificial Neural Networks.

That is profiles computed by "learning" from observation of examples.




Better predictions are obtained from MSA data than from individual protein sequences.

General principle being, the more information offered, the more reliable the prediction.

Some systems will automatically generate an MSA if offered a solitary protein sequence.

Prediction will be based on the MSA, computed by iterative database searching.



Predicting Tertiary Structure directly from Primary Structure is not currently practical.

De novo protein structure prediction requires better algorithms and more computing power.

Homology modelling requires a reliable Tertiary Structure for a homologous protein.

Tertiary Structure for a protein is predicted by comparison with the homologous structure.

Homology modelling is hampered by low volumes and uneven spread of available structures.

And now ... Once again ... Your turn! Some issue for consideration, discussion and reaction

The Bioinformatics topics mentioned here do not constitute a comprehensive list. What would suggest is missing ... in order of importance?

The term <u>algorithm</u> was mentioned once or twice. There are slightly differing definitions. Pick the one you like best and justify your selection.

Define the three terms <u>Homologue</u>, <u>Paralogue</u> and <u>Orthologue</u>, being ever assiduous to ignore offensive American misspellings!

The is but one basic strategy for computing Pairwise Alignments that is considered optimal. However, this strategy can be implemented to compute either Global Alignments or Local Alignments.

Just informally, how do these two possibilities differ?

Generally speaking, would you compute MSAs using a Global or a Local approach? Briefly justify your choice.

Generally speaking, would you conduct Database Similarity searches using a Global or a Local approach? Briefly justify your choice.

"Sequence alignment only makes sense for sequences representing Homologous entities"

A profound observation made by the ever sagacious David Philip Judge whilst sipping an eventide cup of <u>Tesco</u>'s very cheapest tea in the penthouse suite of his Ivory Tower (personal communication, 2016.06.10).

Consider and comment upon this fundamental truth.

"A Multiple Alignment of Homologous sequences which were a mixture of Orthologues and Paralogues would not be suitable as input data for Phylogenetic analysis"

Another deep one from DPJ

Consider and comment upon this further pearl of enlightenment.

In the course of the dialogue for this presentation, there was mention of "Accepted Substitutions", more formally referred to as "Accepted Point Mutations", or ... if you enjoy clumsy for the sake of a pronounceable acronym, "Point Accepted Mutation" (PAM).

How would you informally define an "Accepted Point Mutation"?

definition. ScanProsite being the program for searching the of the Prosite database. Prosite was first created way back in the 1980s and, initially, was composed exclusively of protein patterns.

There is no great value, at this stage, to be entirely familiar with this year, simple syntax.

The Extended syntax for ScanProsite is the most common syntax used for protein pattern

There is no great value, at this stage, to be entirely familiar with this very simple syntax. However, from the hints in this presentation and a quick glance at the appropriate web pages, can you interpret the pattern?

 $C\{P\}x(3,7)[FY](2)Wx(2)[VIL]$

Define both of these terms and describe simply the difference between them

Define both of these terms and describe simply the <u>difference between them</u>.

In the slide notes, there is mention of Position Weight Matrices (PWMs).

Can you say, simply, what a Position Weight Matrix might be and how it might be used?

What obvious property does a PWM possess that is lacking in a simple sequence pattern (or consensus sequence)?

The best secondary structure programs are reckoned to be around 80% accurate.

It is further suggested that 80% is about as good as it is possible to achieve.

Stated simply, why would you suppose that 100% accuracy might be unobtainable?

Hint: Do you think that two human experts, given the very best evidence of Tertian

Hint: Do you think that two human experts, given the very best evidence of Tertiary Structure, would also agree upon the exact amino acid positions where an Alpha Helix starts and finishes?

Homology Modelling is mentioned in the slides as a method for predicting tertiary structure when structure(s) of protein(s) homologous to the query protein are available. The process involves aligning the query protein with the known structure, using the known sequence as a guide.

It is also possible to predict Tertiary Structure when, known structures thought to be appropriate exist, but only for sequences that ARE NOT HOMOLOGOUS. In such cases, the Primary Sequence corresponding to the known structure will be of little assistance.

Tricky eh!? What are the name(s) for <u>those types of method</u>? ONLY if you can do so VERY simply. Say a few words to say how they over come the lack of a homologous sequence.

It was noted in the slides that often different Protein Feature searches often do not exactly agree.

It is common for two services to agree upon the presence of a domain, but not upon it precise start and end positions within a protein.

Would you find this to be worrying? Surprising? If not, why not?

End of Part 2


BREAK

More to come I fear ... but time for a swift cup of tea perchance?

Maybe time for a short jig? The whistling of a merry tune?

Or, mayhap, a delving into the melodic possibilities of youtube? There be much good stuff there ... I offer you a few of my favourites.











Once fully refreshed Click on mon braves!