

Ancestry Specific Allele Frequency Estimation (ASAFE)

Qian Sophia Zhang

Joint work with Dr. Sharon Browning and Dr. Brian Browning
Department of Biostatistics
University of Washington

July 12, 2016

① Motivation for ASAFE

② Available Data

③ Proposed Approach

④ Data Simulation

⑤ Results

Genetic Terminology

Human genomes are packaged into pairs of homologous chromosomes. 2 pairs shown below. Rows are SNPs.

- **"SNP"** := Point along a chromosome where genomes differ.
Often there are two variants or "alleles" at a SNP, labeled 0 and 1.

SNP1	0	1	SNP1	0	1	<div>Both pairs: SNP1 0/1 and SNP2 0/1. Left pair: SNP1 0 1 and SNP2 1 0. Right pair: SNP1 0 1 and SNP2 0 1.</div>
SNP2	1	0	SNP2	0	1	
SNP3	0	0	SNP3	0	0	
SNP4	1	1	SNP4	1	1	
1 pair of hom. Chr's			Another pair of hom. Chr's			

Genetic Terminology

Human genomes are packaged into pairs of homologous chromosomes. 2 pairs shown below. Rows are SNPs.

- "SNP" := Point along a chromosome where genomes differ.
Often there are two variants or "alleles" at a SNP, labeled 0 and 1.
- "Genotype" := 2 homologous chromosomes' alleles at a SNP
Ex: SNP1's genotype is 0/1, or 0|1 or 1|0. / and | denote phase.

SNP1	0	1
SNP2	1	0
SNP3	0	0
SNP4	1	1

1 pair of
hom. Chr's

SNP1	0	1
SNP2	0	1
SNP3	0	0
SNP4	1	1

Another pair of
hom. Chr's

Both pairs: SNP1 0/1 and SNP2 0/1.
Left pair: SNP1 0|1 and SNP2 1|0.
Right pair: SNP1 0|1 and SNP2 0|1.

Genetic Terminology

Human genomes are packaged into pairs of homologous chromosomes. 2 pairs shown below. Rows are SNPs.

- "Unphased genotype" (/): SNP's genotype is NOT ordered with respect to another SNP's genotype

SNP1	0	1	SNP1	0	1	<div>Both pairs: SNP1 0/1 and SNP2 0/1. Left pair: SNP1 0 1 and SNP2 1 0. Right pair: SNP1 0 1 and SNP2 0 1.</div>
SNP2	1	0	SNP2	0	1	
SNP3	0	0	SNP3	0	0	
SNP4	1	1	SNP4	1	1	
1 pair of hom. Chr's			Another pair of hom. Chr's			

Genetic Terminology

Human genomes are packaged into pairs of homologous chromosomes. 2 pairs shown below. Rows are SNPs.

- "Unphased genotype" (/): SNP's genotype is NOT ordered with respect to another SNP's genotype
- "Phased genotype" (|): SNP's genotype IS ordered with respect to another SNP's genotype

SNP1	0	1	SNP1	0	1	<div>Both pairs: SNP1 0/1 and SNP2 0/1. Left pair: SNP1 0 1 and SNP2 1 0. Right pair: SNP1 0 1 and SNP2 0 1.</div>
SNP2	1	0	SNP2	0	1	
SNP3	0	0	SNP3	0	0	
SNP4	1	1	SNP4	1	1	
1 pair of hom. Chr's			Another pair of hom. Chr's			

Genetic Terminology

Human genomes are packaged into pairs of homologous chromosomes. 2 pairs shown below. Rows are SNPs.

- "Unphased genotype" (/): SNP's genotype is NOT ordered with respect to another SNP's genotype
- "Phased genotype" (|): SNP's genotype IS ordered with respect to another SNP's genotype
- Alleles on the same side of | are on the same chromosome, but not necessarily for /


SNP1	0	1	SNP1	0	1	<div>Both pairs: SNP1 0/1 and SNP2 0/1. Left pair: SNP1 0 1 and SNP2 1 0. Right pair: SNP1 0 1 and SNP2 0 1.</div>
SNP2	1	0	SNP2	0	1	
SNP3	0	0	SNP3	0	0	
SNP4	1	1	SNP4	1	1	
1 pair of hom. Chr's			Another pair of hom. Chr's			

Hispanic Community Health Study (HCHS)

- Cohort study of 13,000 US Hispanics
- Hispanics are admixed, descended from multiple ancestral populations: Africans, Europeans, and Native Americans
- In this cohort, we test each bi-allelic SNP for association with a trait, say diabetes
- If a SNP is significantly associated with a trait, we want to perform a follow-up study on new individuals to see if we can replicate the association

The Problem that ASAFE Solves

- For a significant SNP, want **ancestry-specific allele frequencies** $\coloneqq P(\text{Allele 1} \mid \text{African}), P(\text{Allele 1} \mid \text{European}), \text{ and } P(\text{Allele 1} \mid \text{Native American})$, i.e. frequencies of allele 1 amongst chromosomes of African, European, or Native American origin at the SNP



- ASAFE \coloneqq EM algorithm for estimating these frequencies, for a SNP

How Ancestry Specific Allele Frequencies Relate to HCHS

- These frequencies inform the design of a replication study: If allele 1 were more common amongst the African chromosomes than amongst the other two ancestries' chromosomes, then one would want to recruit a population of predominantly African descent for the replication study

① Motivation for ASAFE

② Available Data

③ Proposed Approach


④ Data Simulation

⑤ Results

Available Data

- At some SNPs, the RFMix program takes phased genotypes as input, and outputs admixed individuals' phased ancestries
- Available data on admixed individuals
 - Phased ancestries, Phased genotypes: Some SNPs
 - No ancestry calls, Unphased genotypes: Other SNPs

Available Data



① Motivation for ASAFE

② Available Data

③ Proposed Approach

④ Data Simulation


⑤ Results

Proposed Approach: Fill in Ancestries

Consider a block of SNPs that have been genotyped in the admixed sample, but that do not have ancestries called. For any SNP in this block:

- Call the SNP's ancestry the nearest flanking ancestry

Then we know all SNPs' unphased genotypes and local ancestry pairs.



Proposed Approach: EM Algorithm to Deal with Unknown Phase of Genotype Relative to Ancestry Pair (ASAFE)

Consider an ancestry-specific allele (allele, ancestry) = (g,a):

- Allele $g = 0$ or 1
- Ancestry $a = \text{African (A), European (E), Native American (N)}$

There are 6 possible (g,a) alleles, so 21 values for unordered (g,a)/(g,a) genotype. We call these values unordered (g,a)-genotype categories.

1 complete observation = The (g,a)-genotype category that **an individual** belongs to at a SNP.

Complete, Unobserved Data Categories

Entry C_i is the name of the i -th complete, unobserved category.

Table: Complete Data Unordered (g,a)-genotype Categories.

$(g,a) \backslash (g',a')$	(0, A)	(0, E)	(0,N)	(1, A)	(1, E)	(1, N)
(0, A)	C_1					
(0, E)	C_2	C_4				
(0, N)	C_3	C_5	C_6			
(1, A)	C_7	C_9	C_{11}	C_{16}		
(1, E)	C_8	C_{12}	C_{14}	C_{17}	C_{19}	
(1, N)	C_{10}	C_{13}	C_{15}	C_{18}	C_{20}	C_{21}

Hardy-Weinberg Equilibrium assumed in the admixed population to get probability p_j of an individual falling into the j -th complete data category:

$$p_j = \begin{cases} p_{ga}p_{g'a'}, & \text{if } (g, a) = (g', a') \\ 2p_{ga}p_{g'a'}, & \text{otherwise} \end{cases}$$

Incomplete, Observed Data Categories

Entry O_i is the name of the i -th incomplete, observed category. Colored entries are observed data categories that map to multiple complete data categories.

Table: Incomplete, Observed Data Categories.

$(g,a) \setminus (g',a')$	(0, A)	(0, E)	(0,N)	(1, A)	(1, E)	(1, N)
(0, A)	O_1					
(0, E)	O_2	O_4				
(0, N)	O_3	O_5	O_6			
(1, A)	O_7	O_8	O_9	O_{13}		
(1, E)	O_8	O_{10}	O_{11}	O_{14}	O_{16}	
(1, N)	O_9	O_{11}	O_{12}	O_{15}	O_{17}	O_{18}

Overlaying complete and observed categories gives their correspondence. This correspondence allows us to express the probability p'_j of an individual being in observed data category $j', j' \in \{1, \dots, 18\}$ in terms of complete data category probabilities $p_j, j \in \{1, \dots, 21\}$.

Outline Approach to Estimating Ancestry-Specific Allele Frequencies

Because of the connection

- Between p'_j and p_j , and
- Between p_j and (g,a)-allele probabilities

$$\vec{p} = [p_{ga} : g \in \{0, 1\}, a \in \{A, E, N\}],$$

maximizing the observed data log likelihood (e.g. via EM algorithm ASAFE)

$$\log(P(\vec{o} = [o_1, \dots, o_n]) | \vec{p}' = [p'_1, \dots, p'_{18}]) = \sum_{j'=1}^{18} m'_{j'} \log(p'_{j'})$$

where o_i = Observed category of the i-th individual, and $m'_{j'}$ = Number of individuals in observed category j'

gives us a maximum likelihood estimate (MLE)

$\hat{\vec{p}} = [\hat{p}_{0A}, \hat{p}_{0E}, \hat{p}_{0N}, \hat{p}_{1A}, \hat{p}_{1E}, \hat{p}_{1N}]$ of \vec{p} , from which we obtain ancestry-specific allele frequency estimates:

$$\hat{p}_{1|a} = \hat{p}_{1a} / \hat{p}_a = \hat{p}_{1a} / (\hat{p}_{1a} + \hat{p}_{0a}), a \in \{A, E, N\} \leftarrow \text{THE GOAL!}$$

① Motivation for ASAFE

② Available Data

③ Proposed Approach

④ Data Simulation

⑤ Results

Simulated Genetic Data

- Used MaCS [Chen et al. (2009)] to simulate Hispanic individuals' sequence data
- For each of the 56,003 SNPs in the sequence data, ran ASAFE with inputs: Unphased admixed genotypes (ignoring known phase) and phased admixed ancestries
- Got ancestry-specific allele 1 frequencies for each ancestry (African, European, Native American), at each SNP
- For each SNP, calculated
error = Estimated $p_{1|a}$ - True $p_{1|a}$, $a \in \{A, E, N\}$

① Motivation for ASAFE

② Available Data

③ Proposed Approach

④ Data Simulation

⑤ Results

Low Error on Simulated Data

Mean and SD of errors $\{\hat{p}_{1|a} - p_{1|a} : a \in \{A, E, N\}\}$, grouped by:

- True allele frequency bin that $p_{1|a}$ falls into: Columns
- Ancestry $a \in \{A, E, N\}$: Rows

Ancestry	Statistic	True Allele 1 Frequency Bins				
		(0-0.2]	(0.2-0.4]	(0.4-0.6]	(0.6-0.8]	(0.8-1]
African	Mean	-0.0011	-0.0003	-0.0004	0.0004	-0.0004
African	SD	0.0065	0.0185	0.0233	0.0186	0.0118
European	Mean	-0.0015	-0.0004	-0.0007	-0.0010	<0.0001
European	SD	0.0077	0.0209	0.0249	0.0220	0.0122
Nat. Am.	Mean	-0.0004	-0.0017	0.0021	0.0048	0.0007
Nat. Am.	SD	0.0083	0.0235	0.0238	0.0257	0.0118

Regardless of true ancestry-specific allele frequency $p_{1|a}$ bin, errors are low:
Largest $|\text{Mean}| = 0.005$. Largest SD = 0.03.

More Results in paper supplement.

More Info: Paper and Code

- Qian S. Zhang, Brian L. Browning, and Sharon R. Browning. Asafe: ancestry-specific allele frequency estimation. *Bioinformatics*, 32(14):2227-2229, 2016.
- Package "ASAFE" on Bioconductor
- Code to reproduce analysis at <http://biostatqian.github.io/ASAFE/>

