


Putting together:

$$\beta_j^{\text{lasso}} = \begin{cases} \beta_j^{\text{ls}} - \frac{\lambda}{2} & \beta_j^{\text{ls}} - \frac{\lambda}{2} > 0 \\ 0 & \frac{\lambda}{2} \geq \beta_j^{\text{ls}} \geq -\frac{\lambda}{2} \\ \beta_j^{\text{ls}} + \frac{\lambda}{2} & \beta_j^{\text{ls}} + \frac{\lambda}{2} < 0 \end{cases}$$

Introduce the soft thresholding function

$$S_t(z) \stackrel{\text{def}}{=} \text{sgn}(z)(|z| - t)_+ = \begin{cases} z - t & z > t \\ 0 & t \geq z \geq -t \\ z + t & -t > z \end{cases} \quad (t > 0)$$

where $\text{sgn}(z) \stackrel{\text{def}}{=} \begin{cases} 1 & z > 0 \\ 0 & z = 0 \\ -1 & z < 0 \end{cases}$




$$y_+ \stackrel{\text{def}}{=} \begin{cases} y & y \geq 0 \\ 0 & y < 0 \end{cases}$$


Then $\beta_j^{\text{lasso}} = S_{\frac{\lambda}{2}}(\beta_j^{\text{ls}})$

3 Dimension Reduction.


What is the true dimensionality of this data?




What is the true dimensionality of this data?



What is the true dimensionality of this data?



What is the true dimensionality of this data?



Goed: find the most important features


for a task to reduce the number of features
(PCA)
3.1 Principal Component Analysis

Goal of PCA: find low-dimensional representation of the sample points that maximizes the data variance.

3.1.2 Math Prep.

Recall that an orthogonal matrix $U_{n \times n}$ ($U^T U = I$) preserves the Euclidean norm of a vector, i.e. $\|Ux\| = \|x\| \quad \forall x \in \mathbb{R}^n$

$$\begin{aligned} & \text{(because } \|Ux\|^2 = (Ux) \cdot (Ux) \\ & = (Ux)^T Ux = x^T \underbrace{U^T U}_{=I} x \\ & = x^T x = \|x\|^2) \end{aligned}$$



Def: For a symmetric matrix A and a non-zero vector u , the ratio

$$\frac{u^T A u}{u^T u} = \frac{u^T A u}{\|u\|^2}$$

is called the Rayleigh quotient.

Let $A \in \mathbb{R}^{n \times n}$ be symmetric with eigenvalues $\lambda_1 \geq \dots \geq \lambda_n$ and associated ^{normalized} eigenvectors $u^{(1)}, \dots, u^{(n)}$. Then the eigenvalue decomp

$$A = U \Lambda U^T = \begin{pmatrix} | & & | \\ u^{(1)} & \dots & u^{(n)} \\ | & & | \end{pmatrix} \begin{pmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{pmatrix} \begin{pmatrix} | & & | \\ u^{(1)} & \dots & u^{(n)} \\ | & & | \end{pmatrix}^T$$

Remark: Notice $\frac{u^T A u}{u^T u} = \frac{u^T A u}{\|u\|^2} = \left(\frac{u}{\|u\|}\right)^T A \left(\frac{u}{\|u\|}\right)$ so it suffices to consider the Rayleigh quotient for unit vectors.

Prop. $\lambda_n \leq \frac{u^T A u}{u^T u} \leq \lambda_1$

Each " $=$ " holds if and only if u is the associated eigenvector.