

Function:- $\arccos(x)$

Problem-2

Birva Shah (Student ID: 40070973)

12 July 2019

1 Assumptions

- In function $\arccos(x)$, x is a real number.
- Function returns the value of $\arccos(x)$ in radian.
- If the argument of the function is NaN, then the result is NaN.

2 Requirements

(a) **ID** : REQ-1

Type : Functional Requirement

Description : User shall give input value x between -1 and 1 inclusive to satisfy the constraint that the domain of the function $\arccos(x)$ is $-1 \leq x \leq 1$.

Rationale : The rationale behind this requirement is that the output of the function $\arccos(x)$ is undefined if the value of x is not between -1 and 1 inclusive.

Difficulty : Easy

(b) **ID** : REQ-2

Type : Functional Requirement

Description : The system shall take input x to give the output of the function in radian. For example: $\arccos(0.5) = 1.4719\dots$

Rationale : The rationale behind this requirement is that only one input x which is real number is required to calculate result of $\arccos(x)$.

Difficulty : Nominal

(c) **ID** : REQ-3

Type : Functional Requirement

Description : The system shall calculate the value of $\arccos(x)$ up to the precision of four decimals to get the stable output. For example: $\arccos(0.5) = 1.4719$

Rationale : The rationale behind this requirement is that the function might give an output that has infinite number of decimal points.

Difficulty : Nominal

References

[1] RapidTables,
<https://www.rapidtables.com/math/trigonometry/arccos.html>

[2] Emathhelp,
<https://www.emathhelp.net/notes/algebra-2/trigonometry/function-y-arccos-x/>

[3] Microsoft,
<https://docs.microsoft.com/en-us/powerapps/maker/canvas-apps/functions/function-trig>

[4] Mathonweb,
http://mathonweb.com/help_ebook/html/algorithms.htm*arcsinCliffsNotes*,
<https://www.cliffsnotes.com/study-guides/trigonometry/inverse-functions-and-equations/inverse-cosine-and-inverse->