




The sspm R package: spatial surplus production models for the management of northern shrimp fisheries

Valentin Lucet^{*1} and Eric Pedersen^{†1}

¹ Concordia University, Montreal, Quebec, CA

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: [Open Journals](#) 

Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Abstract

Productivity models such as Surplus Production Models (SPMs) models can be used to inform stock management of fisheries. However, those models often share three main flaws: (1) they are usually not spatially explicit, (2) fail to incorporate ecosystem predictors and therefore are ill-suited to ecosystem-based management of stocks, and (3) their deployment is often limited by code availability, quality and accessibility. To fill this gap, we developed a lag-1 autoregressive spatial SPM based on Generalized Additive Models (GAMs), broadly applicable to spatially-structured populations, and bundled into an R package. We applied this model to one of the most economically important invertebrate populations in Canadian waters, Northern Shrimp (*Pandalus borealis*) in the Newfoundland and Labrador Shelves. This stock currently lacks a population model to predict how fishing pressure and changing environmental conditions may affect future shrimp abundance in the region. Our model incorporates relevant ecosystem predictors for this stock, such as Atlantic Cod (*Gadus morhua*) density, alternate predator density, temperature, and stock biomass. In addition, the model is deployed through the R package sspm, a flexible framework aimed at making SSPMs easier to apply to spatially structured populations. The package allows for a repeatable and open workflow and improves the accessibility of SSPMs.

Summary

The R package sspm is designed to make spatially-explicit surplus production models more applicable. The package uses Generalized Additive Models (GAMs) to fit a surplus production model to biomass and harvest data. The package includes a range of features to manage biomass and harvest data. Those features are organised in a stepwise workflow, whose implementation is described in more detailed in (??).

1. Ingestion of variables as well as spatial boundaries and discretization into patches, using the user's method of choice (random or custom sampling, voronoi tessellation or delaunay triangulation).
2. Smoothing data using spatio-temporal GAMs smoothers.
3. Computation of productivity values taking into account harvest information.
4. Fitting of SPMs to smoothed data with GAMs.
5. Visualization of results, including confidence and prediction intervals.
6. One step ahead prediction of biomass.

Although it was developed in a fisheries context, the package is suitable to model spatially-structured population dynamics in general.

^{*}co-first author

[†]co-first author

Statement of need

1. Population models, in particular fisheries productivity models, rarely integrates important spatially-structured ecosystem drivers
2. The Northern Shrimp stock in the Newfoundland and Labrador Shelves currently lacks a population model
3. Current SPM models are rarely spatially explicit and usually cannot account for relevant ecosystem drivers
4. Fisheries managers lack user-friendly, flexible tools to implement and apply Spatial SPMs

Introduction

Population modelling is an exercise of interest within environmental sciences and adjacent fields. From early models that addressed simple dynamics such as exponential growth and density dependence, modern models are now acknowledging the non-stationary nature of wild populations. In addition, population models applied to resource management, such as fisheries models, are increasingly concerned with how stocks varies across time and space. Resource managers are becoming more and more interested in how ecosystem factors such as predator abundance or abiotic variables impact the spatio-temporal variability of mechanisms like productivity and density dependence.

Population models in fisheries science usually fall under two categories: process-based models and statistical models. Process based models often rely on differential equations and are based on replicating the underlying processes (predation, recruitment, dispersal) behind population dynamics. Statistical models, on the other hand, rely on fitting a model to data using distributional assumptions, and present the advantage of naturally measuring uncertainty around predictions. This is useful in a management context where uncertainty around decision-making is an important information to have on hand.

In this paper, we implement a statistical model for the population of northern shrimp of the Newfoundland and Labrador Shelves that accounts for varying productivity across time and space. The model is implemented via a R package designed for this type of spatial surplus productivity modelling, the *sspm* package. We exemplify how to successfully implement a spatial model with *sspm* and discuss the applicability of the framework to the other spatially structured populations.

Model

Surplus production models (SPMs), are simply defined as such:

$$b_{t+1} = g(b_t) * e^{\epsilon}$$

With b the biomass, e^{ϵ} an error term, and g a function of the biomass and time which, in the case of fisheries, usually involves some measure of harvest through catch.

Rho => varying prod => covariance matrix => precision => basis functions and Gams


Results

The GAM biomass estimates are consistent with those of the current tool in use for the assessment of the stock, Ogmapp, and provide valuable insights about the drivers of the rapid increase and decline of shrimp in the southern end of the shelf. Our approach demonstrates the model's ability to become a useful tool for modelling spatially-structured populations like fisheries stocks. The *sspm* package successfully modularizes each step of the modelling process

and implements a range of useful features for modeling spatially-structured populations: spatial
discretization, simplified GAM syntax, prediction intervals and scenario based forecasts for
longer-term trends. In a fisheries context, It illustrates how our model can be easily used by
managers to forecast fisheries productivity under different management regimes. The package
is also a tool to think about design choices when conceiving a user interface for managers and
on best practices when it comes to adapting research code into management tools. Finally,
our approach demonstrates how open source software tools can improve the accessibility and
reliability of models for fisheries management.

87 Package design

⁸⁸ The package follows an object oriented design, making use of the S4 class systems.



90 Application to simulated data

91 We present an example using simulated biomass and harvest data. Using real trawl and fishing
92 data provided by DFO, we generated fake data for each spatio-temporal units. ...

93 Acknowledgements

94 TBD

⁹⁵ **References**

⁹⁶ TBD

DRAFT