

An exact tidal-parity threshold in the face-on solar-sail CR3BP

Technical note — Bishwaswarup Nayak, Indian Institute of Science, Bengaluru

bishwaswarup@iisc.ac.in • September 25, 2026

Setting. A face-on sail ($\alpha = 0$) exerts a purely radial, inverse-square force, so it is conservative and rescales solar gravity: $(1 - \mu) \rightarrow (1 - \beta)(1 - \mu)$. The problem is then a photogravitational CR3BP, with pseudo-potential $U = \frac{1}{2}(x^2 + y^2) + (1 - \beta)(1 - \mu)/r_1 + \mu/r_2$. On the segment between the primaries $r_1 + r_2 = 1$ *identically*, which is what makes everything below exact rather than asymptotic.

1. A closed form for the tidal-parity threshold. Write the smaller primary's direct contribution to the collinear linearisation as the saddle strength $s = \mu/r_2^3$, and define tidal parity by $s = 1$, i.e. $r_2 = \mu^{1/3} = 3^{1/3}r_H$. At that radius the tidal acceleration equals the standoff, $\mu/r_2^2 = \mu^{1/3} = r_2$. Imposing this on the on-axis balance $x = (1 - \beta)(1 - \mu)/r_1^2 - \mu/r_2^2$, with $r_1 = 1 - \mu^{1/3}$ exact, the $\mu^{1/3}$ terms cancel and

$$\beta_{\text{crit}} = 1 - (1 - \mu^{1/3})^2 = 2\mu^{1/3} - \mu^{2/3} \quad (1)$$

for every mass ratio — no expansion in μ , no root-finding. Sun–Earth: $\beta_{\text{crit}} = 0.028646456169$, standoff 2 158 294 km. Checked against independent Brent root-finding on $s(\beta) - 1$ at machine precision (2.1×10^{-16} at Sun–Earth; $\leq 1.9 \times 10^{-15}$ over a 61-point sweep of $\mu \in [10^{-7}, 10^{-2}]$).

2. Why $s = 1$ is not an arbitrary level. Generalise to $s = \kappa$, so $r_2 = (\mu/\kappa)^{1/3} = (3/\kappa)^{1/3}r_H$. The same substitution gives a closed form for *every* κ ,

$$\beta(\kappa) = 1 - \frac{(1 - r_2)^2}{1 - \mu} \left[(1 - \mu) + \underbrace{\mu^{1/3}\kappa^{-1/3}(\kappa - 1)}_{\Delta(\kappa)} \right], \quad (2)$$

and the residue satisfies $\Delta(\kappa) = 0 \iff \kappa = 1$. Only at parity does the smaller primary drop out of the balance identically, leaving $1 - \beta = r_1^2$ and hence the pure square (1). Two markers already in the literature are members of this one family: the Hill-sphere exit $r_2 = r_H$ is exactly $\kappa = 3$, and the classical L_1 sits at $\kappa = 3.0303$.

Table 1: The hierarchy for Sun–Earth from (2); $\tau_u = 1/\lambda_u$ is the e-folding time of the residual saddle.

κ	β	r_2/r_H	standoff [km]	τ_u [d]	
3.030 3	0.000 000	0.996 7	1 491 472	23.0	classical L_1
3	0.000 298	1.000 0	1 496 477	23.1	Hill-sphere exit
1	0.028 646	1.442 2	2 158 294	38.6	tidal parity
0.1	0.087 462	3.107 2	4 649 903	109.7	

3. A universal bound on the frequency ratio. The collinear parameter $A = (1 - \beta)(1 - \mu)/r_1^3 + \mu/r_2^3$ is exactly Richardson's c_2 ; a sail detunes it continuously. With $\nu = \sqrt{c_2}$ out of plane and $\omega^2 = \frac{1}{2}[D - (c_2 - 2)]$, $D = \sqrt{c_2(9c_2 - 8)}$, in plane, the ratio depends on c_2 alone. Setting $d(\nu^2/\omega^2)/dc_2 = 0$ reduces to $D = 2c_2$, hence $9c_2^2 - 8c_2 = 4c_2^2$ and

$$c_2^* = \frac{8}{5}, \quad \lambda_u^2 : \nu^2 : \omega^2 = 7 : 8 : 9, \quad \left. \frac{\nu}{\omega} \right|_{c_2^*} = \frac{2\sqrt{2}}{3}, \quad (3)$$

so that over the whole admissible range $c_2 > 1$,

$$\frac{\nu}{\omega} \in \left[\frac{2\sqrt{2}}{3}, 1 \right), \quad \text{a band 5.72 \% wide.} \quad (4)$$

Both c_2^* and the bound are independent of μ : they are properties of the collinear linearisation, shared by every three-body system, sail or no sail. Ceccaroni, Celletti & Pucacco note that the

centre-manifold frequencies are “almost equal for all mass ratios”; (4) makes that exact and bounds the detuning $\delta = \omega - \nu$ that normal-form treatments of the collinear points rely on.

4. No low-order commensurability is available. The largest rational below unity at denominator q is $(q-1)/q$, so $p:q$ can lie in (4) only if $(q-1)/q \geq 2\sqrt{2}/3$, i.e. $q \geq (1-2\sqrt{2}/3)^{-1} = 17.485\dots$, so $q \geq 18$. *The band contains no commensurability of denominator below 18*; the lowest-order one it admits is 17:18, of order 35. Neither threshold is therefore a bifurcation — $s > 0$ for all finite r_2 , so the saddle never disappears — and I present both as exact structural landmarks rather than dynamical transitions.

5. Where this sits against flown hardware. Reducing β from primary specifications ($\beta = \sigma^*/\sigma$, $\sigma^* = 1.531 \text{ g m}^{-2}$), no sail flown and deployed reaches $\beta = 0.0062$: LightSail-2 is 0.00613, and the only *measured* value is IKAROS at 0.00062. Tidal parity is a factor 4.7 beyond the best deployed sail and 1.4 beyond Solar Cruiser as designed.

What I would most value your view on. (i) Whether the closed form (1) — or the κ family (2) — has appeared before in the photogravitational CR3BP literature, which I have not been able to search exhaustively. (ii) Whether “tidal parity” is a defensible reading of $s = 1$, or whether the κ -hierarchy framing of §2 is the honest way to present it. (iii) Whether the bound (4) is known; I have not found it stated.

The full preprint (18 pp.) and a companion document giving every derivation step and an account of each reported residual are available on request. All code and data are public at https://github.com/Bishwaswarup/SOLAR_SAIL (tag `arxiv-v1`); every number above is regenerated by the pipeline and machine-checked against the manuscript source by `paper/verify_numbers.py`, which asserts 175 numerical claims and parses the tables out of the L^AT_EX directly.