

Dissolution of the collinear structure in the solar-sail circular restricted three-body problem: an exact tidal-parity threshold and a universal frequency bound

Bishwaswarup¹

¹Indian Institute of Science, Bangalore 560012, India, bishwaswarup@iisc.ac.in

September 17, 2026

Abstract

A face-on solar sail does not displace the collinear Lagrange point of the circular restricted three-body problem; it dissolves it. Because the radiation force is radial and therefore conservative, it rescales solar gravity as $(1 - \mu) \rightarrow (1 - \beta)(1 - \mu)$, and the on-axis equilibrium degenerates smoothly into a heliocentric hovering point as the lightness number β grows. We characterise that degeneration and obtain two exact results. First, imposing *tidal parity* — the condition that the smaller primary’s contribution to the linearised dynamics equal the centrifugal background, $s \equiv \mu/r_2^3 = 1$, equivalently $r_2 = \mu^{1/3} = 3^{1/3}r_H$ — on the on-axis force balance yields the closed form $\beta_{\text{crit}} = 1 - (1 - \mu^{1/3})^2 = 2\mu^{1/3} - \mu^{2/3}$, exact for every mass ratio and verified against independent root-finding at machine precision over $\mu \in [10^{-7}, 10^{-2}]$ (2.1×10^{-16} at Sun–Earth). For Sun–Earth this gives $\beta_{\text{crit}} = 0.0286$. Second, writing c_2 for the second Legendre coefficient of the collinear expansion — the quantity a sail detunes continuously, and which is otherwise fixed by the mass ratio and the choice of point — the ratio of the out-of-plane to in-plane linear frequencies at any collinear equilibrium is confined to $\nu/\omega \in [2\sqrt{2}/3, 1)$, a band 5.72% wide containing no low-order rational. The bound is attained exactly at $c_2 = 8/5$, where the three linear rates stand in the ratio $\lambda_u^2 : \nu^2 : \omega^2 = 7 : 8 : 9$. Both $c_2 = 8/5$ and the bound are independent of μ . That the two frequencies are nearly equal at collinear points has been remarked upon [1]; we make the statement exact. We emphasise that neither result is a bifurcation: the saddle persists for all β and the degeneration is smooth, so these are exact structural landmarks rather than dynamical transitions, and we show by explicit search that no commensurability exists anywhere to make one. Reducing β from primary mission specifications, we find that no solar sail yet flown exceeds $\beta = 0.00613$, placing tidal parity a factor of 4.7 beyond demonstrated capability. Finally, linearising the sail acceleration in its two attitude angles shows that the face-on nominal is a *singular* configuration for station-keeping: $\partial \mathbf{a} / \partial \delta$ vanishes identically at $\alpha_0 = 0$, the Kalman rank drops to 4/6, and any non-zero cone angle restores controllability.

Keywords: solar sail; restricted three-body problem; artificial equilibria; collinear points; Legendre coefficient c_2 ; Hill radius; halo orbits; controllability

1 Introduction

Artificial equilibria generated by continuous low thrust have been studied since McInnes *et al.* [2] mapped the surfaces of sail-displaced equilibria in the restricted three-body problem, and the periodic orbits about them are equally well established: Baoyin and McInnes [3] constructed halo orbits at the Sun–Earth artificial L_1 , Waters and McInnes [4] treated families above the

ecliptic, and Farrés and Jorba [5, 6] analysed the quasi-periodic dynamics and station-keeping in the same region. The standard reference remains [7].

What that literature supplies is numerical: the equilibrium abscissa is obtained by root-finding on the on-axis balance for each β , and the orbit families by Lindstedt–Poincaré expansion or numerical continuation. The equilibrium coordinates enter as computed inputs. What it does not supply is an algebraic characterisation of *how the collinear structure itself degrades* as β grows — at what sail performance the smaller primary ceases to organise the dynamics at all, and whether that point admits a closed form.

This paper supplies that characterisation. Our starting observation is that a face-on sail ($\alpha = 0$) exerts a purely radial and hence conservative force, so it does not perturb the collinear point: it rescales one primary’s mass and the equilibrium migrates toward a *heliocentric hovering point* at which reduced solar gravity balances centrifugal acceleration at the synchronous rate. The three-body character of the equilibrium therefore does not shift, it decays, and the useful question is the rate and the scale of that decay.

Section 2 sets out the model and the conservative rescaling. Section 3 defines the saddle strength $s = \mu/r_2^3$ and measures its four-decade collapse. Section 4 derives the exact tidal-parity threshold, the paper’s principal result, and validates it across five decades of mass ratio. Section 5 establishes the frequency bound $\nu/\omega \in [2\sqrt{2}/3, 1)$, locates its exact extremum at $c_2 = 8/5$, and reports a systematic negative: no low-order resonance is attainable anywhere, which is what forbids interpreting either threshold as a bifurcation. Section 6 tracks the halo families across the flown sail band by pseudo-arclength continuation. Section 7 derives the sail control Jacobian and the singularity of the face-on nominal. Section 8 places β_{crit} against every sail flown to date. Section 9 states the limitations plainly.

On terminology. We use *tidal parity* for the condition $s = 1$. The term has recently appeared in translunar astrodynamics for the analogous balance between lunisolar perturbations and Earth gravity; we adopt it in that spirit and claim no coinage. Likewise, the scale $\mu^{1/3}$ is not new — it is the Hill radius up to the factor $3^{1/3}$, and is standard in tidal-stability analysis. Our contribution is not the scale but the exact sail parameter required to place a spacecraft on it.

2 Model

We work in the usual synodic frame of the circular restricted three-body problem, non-dimensionalised so that the primary separation, the total mass and the mean motion are unity. The larger primary sits at $x = -\mu$ and the smaller at $x = 1 - \mu$, with $\mu = M_2/(M_1 + M_2)$. Distances from the primaries are

$$r_1 = \|\mathbf{r} - \mathbf{r}_1\|, \quad r_2 = \|\mathbf{r} - \mathbf{r}_2\|, \quad (1)$$

and on the segment between them $r_1 = x + \mu$ and $r_2 = (1 - \mu) - x$, so that

$$r_1 + r_2 = 1 \quad \text{identically.} \quad (2)$$

Equation (2) is exact, not asymptotic, and is used in Section 4.

2.1 Sail acceleration

We model an ideal specular reflector. With α the cone angle between the Sun line and the sail normal and δ the clock angle about the Sun line,

$$\mathbf{a}_{\text{sail}} = \beta \frac{1 - \mu}{r_1^2} \cos^2 \alpha \, \hat{\mathbf{n}}, \quad \hat{\mathbf{n}} = \cos \alpha \, \hat{\mathbf{r}} + \sin \alpha \cos \delta \, \hat{\mathbf{p}} + \sin \alpha \sin \delta \, \hat{\mathbf{q}}, \quad (3)$$

where β , the lightness number, is the ratio of the peak radiation-pressure acceleration to solar gravity. The force is directed along the normal with magnitude proportional to $\cos^2 \alpha$; this is




Figure 1: Geometry of the face-on solar-sail CR3BP. **(a)** The synodic frame. The classical L_1 lies essentially on the Hill sphere ($r_2 = 0.9967 r_H$); as β grows the equilibrium migrates sunward, reaching the tidal-parity radius $r_2 = \mu^{1/3} = 3^{1/3} r_H$ at $\beta = 0.0286$. The Earth neighbourhood is exaggerated by a factor of 18; at true scale every feature would collapse onto the smaller primary. **(b)** Attitude convention and the orthonormal triad (4). At $\alpha = 0$ the normal is $\hat{r}$, the force is radial, and the sail is conservative.

the specular reflector, and it is distinct from the perfect absorber, for which the force lies along the Sun line with magnitude proportional to $\cos \alpha$. The two coincide only at $\alpha = 0$.

The triad in (3) must be orthonormal. We take

$$\hat{r} = \frac{\mathbf{r} - \mathbf{r}_1}{r_1}, \quad \hat{p} = \frac{\hat{k} \times \hat{r}}{\|\hat{k} \times \hat{r}\|}, \quad \hat{q} = \hat{r} \times \hat{p}, \quad (4)$$

with $\hat{k}$ the out-of-ecliptic unit vector. Substituting $\hat{k}$ itself for $\hat{q}$ — a natural but incorrect shortcut — gives a non-orthonormal triad, since $\hat{r} \cdot \hat{k} = z/r_1 \neq 0$ off the ecliptic, and then $\|\hat{n}\|^2 = 1 + \sin(2\alpha) \sin \delta (z/r_1)$, so that the nominal unit normal is not a unit vector and both the magnitude and the direction of the force are wrong. The error is 1.0% at $z/r_1 = 0.02$ and 4.4% at $z/r_1 = 0.09$. It vanishes identically at $\alpha = 0$, so no face-on result in this paper is affected; we record it because off-axis quantities computed on the shortcut triad are not reliable. Figure 1 shows the geometry.

2.2 The face-on case is conservative

Setting $\alpha = 0$ in (3) gives $\mathbf{a}_{\text{sail}} = \beta(1 - \mu)r_1^{-2}\hat{r}$, a central force derivable from the potential $-\beta(1 - \mu)/r_1$. It therefore merely rescales the larger primary, and the entire problem reduces to a CR3BP with modified pseudo-potential

$$U = \frac{1}{2}(x^2 + y^2) + \frac{(1 - \beta)(1 - \mu)}{r_1} + \frac{\mu}{r_2}. \quad (5)$$

A Jacobi integral survives,

$$C_{\text{sail}} = 2U - v^2, \quad (6)$$

and it is U of (5), not the unscaled gravitational pseudo-potential, that must be used: the two differ by $2\beta(1 - \mu)/r_1$, which varies along an orbit, so the gravitational Jacobi constant is not an integral for $\beta \neq 0$. On a representative family member ($\beta = 0.001$) we measure a spread of 1.8×10^{-15} in C_{sail} against 5.8×10^{-6} in its gravitational counterpart over one full period.

For $\alpha \neq 0$ the force acquires a transverse component, is no longer conservative, and no Jacobi integral exists. Every result in Sections 3–6 is therefore restricted to $\alpha = 0$; Section 7 is the sole exception and is explicitly a statement about the linearisation in the attitude angles.

3 Dissolution of the collinear structure

3.1 The hovering law

Deleting the smaller primary from (5) leaves an on-axis balance whose root is

$$x_{\text{hover}}(\beta) = [(1 - \beta)(1 - \mu)]^{1/3}, \quad (7)$$

the radius at which reduced solar gravity balances centrifugal acceleration at the synchronous rate. This is a heliocentric hovering point, not a perturbed Lagrange point. The full three-body equilibrium satisfies

$$x = \frac{(1 - \beta)(1 - \mu)}{r_1^2} - \frac{\mu}{r_2^2}, \quad (8)$$

and the difference between its root and (7) is the smaller primary's entire contribution. At $\beta = 0.5$ that contribution is 3 817 km against a standoff of 30 865 449 km, i.e. 20.6 Hill radii: the Earth is dynamically irrelevant there, and results quoted at such β describe hovering geometry rather than three-body dynamics.

3.2 Linear structure and the saddle strength

Linearising (5) on the axis gives $U_{xx} = 1 + 2A$, $U_{yy} = 1 - A$, $U_{zz} = -A$ with

$$A(\beta) = \underbrace{\frac{(1 - \beta)(1 - \mu)}{r_1^3}}_S + \underbrace{\frac{\mu}{r_2^3}}_s. \quad (9)$$

This A is not a new quantity. Setting $\beta = 0$ and using $r_1 + r_2 = 1$, (9) reduces to $\mu/\gamma^3 + (1 - \mu)/(1 - \gamma)^3$ with $\gamma = r_2$, which is exactly the second Legendre coefficient c_2 of the collinear expansion introduced by Richardson [8] and used throughout halo-orbit theory [1, 9]. We verify the identity numerically to machine zero at Sun–Earth, Earth–Moon and Sun–Jupiter. We therefore write

$$A \equiv c_2, \quad (10)$$

and use c_2 from here on. The sail's effect is to replace $(1 - \mu)$ by $(1 - \beta)(1 - \mu)$ in that coefficient: *a solar sail continuously detunes c_2* , which is otherwise fixed once the system and the libration point are chosen. This is the compact statement of what a sail contributes, and we return to it in Section 5.

The out-of-plane motion is a pure oscillator, $\nu = \sqrt{c_2}$. The in-plane characteristic polynomial is

$$\lambda^4 + (2 - A)\lambda^2 + (1 + A - 2A^2) = 0, \quad (11)$$

whose roots are $\lambda^2 = \frac{1}{2}[(A - 2) \pm D]$ with $D = \sqrt{A(9A - 8)}$, giving a saddle exponent λ_u and a centre frequency ω ,

$$\lambda_u^2 = \frac{1}{2}[(A - 2) + D], \quad \omega^2 = \frac{1}{2}[D - (A - 2)]. \quad (12)$$

A real hyperbolic pair requires $c_2 > 1$.

As the equilibrium recedes from the smaller primary, $r_1^3 \rightarrow (1 - \beta)(1 - \mu)$ by (7) and hence $S \rightarrow 1$, so

$$c_2 - 1 \longrightarrow s = \frac{\mu}{r_2^3}. \quad (13)$$

Table 1: Collapse of the saddle strength $s = \mu/r_2^3$ and of the unstable eigenvalue with sail lightness number, Sun–Earth ($\mu = 3.003 \times 10^{-6}$). Tidal parity is $s = 1$.

β	s	λ_u [nd]	note
0	3.030 3	2.532 56	classical L_1
0.010	2.123	2.138 31	
0.028646	1.000	1.505 42	tidal parity
0.050	0.401 2	0.994 49	
0.100	0.068 20	0.444 00	
0.500	3.419×10^{-4}	0.035 78	$20.6 r_H$ standoff

We call s the *saddle strength*. Because $s > 0$ for all finite r_2 , the saddle never strictly disappears: there is no bifurcation, only a smooth decay, and the physically meaningful statement is its rate. Table 1 records that rate and Figure 5 the accompanying collapse of λ_u ; s falls four decades between $\beta = 0$ and $\beta = 0.5$.

It is worth being precise about (13), because the convergence is slower than it looks. At tidal parity, $S = (1 - \mu)/(1 - \mu^{1/3}) = 1.014635$ exactly, so $c_2 = 2.014635$ rather than 2, and $\lambda_u = 1.505418$ rather than $5^{1/4} = 1.495349$. The identification $c_2 - 1 = s$ carries an $O(\mu^{1/3})$ error, 0.73 % in c_2 at the Sun–Earth threshold. Both terms in (9) are nonetheless of smaller-primary origin: with the primary deleted, $S = 1$ identically, so $S - 1$ measures its indirect effect through the displacement of the equilibrium, while s measures the direct tidal term. We therefore write $A - 1 = (S - 1) + s$ and use the exact A throughout.

Two consequences of Figure 2(d) deserve emphasis, because both have been misread in the past. As $c_2 \rightarrow 1$ the planar roots of (11) become $\lambda^2 \in \{0, -1\}$ and $\nu = \sqrt{A} \rightarrow 1$, so the in-plane period is driven to exactly 2π in non-dimensional units — one year for Sun–Earth. This is *forced*, not resonant: in an inverse square field the epicyclic frequency equals the mean motion identically. Second, $\lambda_u \rightarrow 0$ does not indicate that the sail has stabilised a saddle; the orbit has simply moved away from one, and the residual instability is precisely the smaller primary’s leftover tidal term.

4 The tidal-parity threshold in closed form

4.1 Derivation

Define tidal parity as the condition that the direct tidal term equal the centrifugal background of the synodic frame,

$$s = \frac{\mu}{r_2^3} = 1 \quad \Longleftrightarrow \quad r_2 = \mu^{1/3} = 3^{1/3} r_H, \quad (14)$$

where $r_H = (\mu/3)^{1/3}$. Below this radius the smaller primary dominates the Hessian; above it the centrifugal and rescaled-solar terms do.

Impose (14) on the balance (8). By the exact identity (2), $r_2 = \mu^{1/3}$ forces

$$r_1 = 1 - \mu^{1/3} \quad (15)$$

with no error term. Substituting, and using $x = (1 - \mu) - r_2$ and $\mu/r_2^2 = \mu^{1/3}$,

$$\underbrace{(1 - \mu) - \mu^{1/3}}_x = \frac{(1 - \beta)(1 - \mu)}{(1 - \mu^{1/3})^2} - \mu^{1/3}. \quad (16)$$




Figure 2: Dissolution of the Sun–Earth collinear structure. (a) The full three-body equilibrium is indistinguishable from the Earth-free hovering law (7); the inset shows the smaller primary’s entire contribution collapsing to a few thousand kilometres. (b) The standoff crosses the Hill sphere almost immediately (at $\beta = 2.98 \times 10^{-4}$, the Hill-exit value, not β_{crit}) and reaches $20.6 r_H$ by $\beta = 0.5$. (c) The saddle strength falls four decades; $s = 1$ locates tidal parity. (d) Both frequencies converge on the mean motion and $\lambda_u \rightarrow 0$: the Keplerian epicyclic degeneracy of an inverse square field, not a resonance with the smaller primary.

The $\mu^{1/3}$ terms cancel identically, leaving $(1 - \mu) = (1 - \beta)(1 - \mu)(1 - \mu^{1/3})^{-2}$ and hence $1 - \beta = (1 - \mu^{1/3})^2$. Therefore

$$\boxed{\beta_{\text{crit}} = 1 - (1 - \mu^{1/3})^2 = 2\mu^{1/3} - \mu^{2/3}} \quad (17)$$

This is exact for every μ ; the cancellation is algebraic and involves no expansion in μ . To leading order $\beta_{\text{crit}} \simeq 2\mu^{1/3}$.

The same substitution gives the linear structure at parity in closed form,

$$c_2^{\text{parity}} = 1 + \frac{1 - \mu}{1 - \mu^{1/3}}, \quad (18)$$

which for Sun–Earth is 2.014635, the value quoted in Section 3.

4.2 Verification

We evaluate (17) against independent Brent root-finding on $s(\beta) - 1$, where s is computed from the numerically solved equilibrium (8). Table 2 reports both across five decades of mass

Table 2: The closed form (17) against independent root-finding on $s(\beta) = 1$. Agreement is at machine precision for every mass ratio tested. $r_2/r_H = 3^{1/3} = 1.4422$ at parity by construction, independent of μ .

system	μ	β_{crit} (closed form)	β_{crit} (root-find)	$ \Delta $	r_2 [nd]
Sun–Mercury	1.660×10^{-7}	0.010 961 524 792	0.010 961 524 792	3.1×10^{-16}	0.005 496
Sun–Earth	3.003×10^{-6}	0.028 646 456 169	0.028 646 456 169	2.1×10^{-16}	0.014 427
Sun–Jupiter	9.537×10^{-4}	0.187 175 530 204	0.187 175 530 204	2.2×10^{-16}	0.098 432
Earth–Moon	1.215×10^{-2}	0.406 934 946 280	0.406 934 946 280	5.0×10^{-16}	0.229 893

ratio. Over a 61-point logarithmic sweep of $\mu \in [10^{-7}, 10^{-2}]$ the two never differ by more than 1.9×10^{-15} , i.e. a few units in the last place; at Sun–Earth the discrepancy is 2.1×10^{-16} . The two computations share no code path beyond the balance itself.

For Sun–Earth the threshold corresponds to a standoff $r_2 = 0.014\,427\,\text{nd} = 2\,158\,294\,\text{km} = 1.4422\,r_H$. For reference, the classical L_1 already sits at $r_2 = 0.9967\,r_H$, essentially on the Hill sphere, so the Hill-sphere exit criterion $r_2 = r_H$ is reached almost immediately, at $\beta = 2.9814 \times 10^{-4}$, and is not a useful discriminant.

5 The frequency ratio: a universal bound on c_2

The threshold (17) is a level crossing of a smooth monotone function. Nothing bifurcates there, and the choice of the level $s = 1$ is partly a statement about the non-dimensionalisation. The standard route to promoting such a marker into a dynamical event is a commensurability of the linear frequencies. This section searches for one and reports that none exists — together with an exact extremum that does.

5.1 The bound and its extremum

From (12), the ratio of the out-of-plane to in-plane frequency depends on c_2 alone,

$$\frac{\nu^2}{\omega^2} = \frac{2c_2}{D - (c_2 - 2)}, \quad D = \sqrt{c_2(9c_2 - 8)}. \quad (19)$$

The two frequencies in (12) are standard — they appear in this form in [9] and [1] — but their ratio is, to our knowledge, not formed anywhere. Doing so is what follows.

Differentiating (19), the stationarity condition reduces to $D = 2c_2$, i.e. $9c_2^2 - 8c_2 = 4c_2^2$, giving

$$\boxed{c_2^* = \frac{8}{5}} \quad (20)$$

and there, from (12),

$$\lambda_u = \sqrt{\frac{7}{5}}, \quad \nu = \sqrt{\frac{8}{5}}, \quad \omega = \sqrt{\frac{9}{5}} \quad \implies \quad \lambda_u^2 : \nu^2 : \omega^2 = 7 : 8 : 9, \quad (21)$$

$$\left. \frac{\nu}{\omega} \right|_{c_2^*} = \sqrt{\frac{8}{9}} = \frac{2\sqrt{2}}{3} = 0.942809041582\dots \quad (22)$$

Over the whole physically admissible range $c_2 > 1$ the ratio is confined to

$$\frac{\nu}{\omega} \in \left[\frac{2\sqrt{2}}{3}, 1 \right), \quad (23)$$

a band 5.72% wide, approaching unity in the hovering limit $c_2 \rightarrow 1$ and again as $c_2 \rightarrow \infty$. Numerically, a 4×10^5 -point scan places the minimum at $c_2 = 1.60000000$ with value

0.942809041582, agreeing with (20) and (22) to 8.1×10^{-10} and 1.1×10^{-16} respectively. Note that c_2^* , the ratios (21) and the bound (22) contain no reference to μ : they are properties of the collinear linearisation itself, shared by every three-body system.

The narrowness of (23) has been observed qualitatively. Ceccaroni *et al.* [1], sweeping all three collinear points over the full range $\mu \in (0, \frac{1}{2}]$, note that the centre-manifold dynamics is “characterized by almost equal values of the frequencies for all mass ratios”. Equations (23) and (22) make that observation exact and supply the constant it is missing.

5.2 No low-order resonance is attainable

We tested fourteen low-order rationals — 1:2, 2:3, 3:4, 4:5, 5:6, 8:9, 1:1, 9:8, 6:5, 5:4, 4:3, 3:2, 5:3, 2:1 — against the band (23). *None* lies inside it. Since $\nu < \omega$ strictly for $c_2 > 1$, even the 1:1 commensurability is approached only asymptotically, as the equilibrium degenerates into the hovering point.

The consequence is that no commensurability underwrites tidal parity, and none exists elsewhere in the face-on solar-sail CR3BP either. Equation (17) must therefore be presented as an exact *scale marker* with a clean geometric reading, not as a dynamical transition. We regard this negative as a result rather than an omission: it is the check a reader is entitled to demand, and it forecloses an over-claim.

Two caveats. First, (23) constrains the *linear* frequencies only. Halo families bifurcate from planar Lyapunov families at a commensurability of the *nonlinear* frequencies, reached at finite amplitude; the classical L_1 has $\nu/\omega = 0.9659$ and halo orbits exist there regardless. Nothing here constrains halo existence. Second, c_2^* is a stationary point, not a transition: the dynamics change character nowhere in c_2 . Its value is that a stationary point is intrinsic — it survives any smooth reparametrisation and any choice of units — whereas a level crossing at $s = 1$ does not.

5.3 What the sail contributes

It would be natural to suppose $c_2^* = 8/5$ inaccessible without continuous thrust, on the grounds that the classical collinear points have $c_2 \geq 4$. That is true only of L_1 . Computing c_2 at all three classical collinear points over $\mu \in (0, \frac{1}{2}]$ gives

$$c_2(L_1) \in [4.042, 8.000], \quad c_2(L_2) \in [1.570, 3.959], \quad c_2(L_3) \in [1.000, 1.570], \quad (24)$$

so the classical problem covers $c_2 < 4$ freely at L_2 and L_3 , and it *does* attain c_2^* : at L_2 with $\mu = 0.480187660$, where ν/ω equals (22) to twelve digits. L_3 at $\mu = \frac{1}{2}$ reaches $c_2 = 1.569787$ and misses the bound by only 3.07×10^{-5} — close enough that the near-coincidence is worth recording, since L_3 occupies precisely the interval in which the extremum lies.

The sail’s contribution is therefore *tunability*, not exclusivity, and (10) is the compact way to say it: at fixed μ a sail sweeps c_2 continuously through $(1, c_2^{\text{classical}}]$, whereas classically c_2 is fixed once the system and the point are chosen. Sun–Earth classically offers only $c_2 = 4.060819$ at L_1 , 3.940764 at L_2 and 1.000003 at L_3 ; none is at the bound, and a sail at $\beta = 0.040932$ places that system exactly on it — a factor 1.4289 beyond tidal parity, at $s = 0.5897$. Figure 3 collects all of this.

6 Halo families across the flown sail band

The band $\beta \in [0.001, 0.05]$ is where a sail both matters dynamically and still inhabits a genuine three-body structure: below it the sail is irrelevant, and above tidal parity the collinear structure has lost its meaning. We therefore walk the halo family at twelve values of β across that band.

Two methodological points are load-bearing. First, all lengths are expressed in units of the local scale $\gamma = |(1 - \mu) - x_{\text{eq}}|$, which doubles across the band (0.0101 at $\beta = 0.001$ to 0.0196 at




Figure 3: The frequency ratio at collinear equilibria. (a) ν/ω against c_2 , with the exact minimum $2\sqrt{2}/3$ at $c_2 = 8/5$ and, in the reserved strip below, the ranges of c_2 reached by the three classical collinear points over $\mu \in (0, \frac{1}{2}]$; the vertical placement of those bars is a layout choice and carries no value. (b) The three linear rates, meeting the ratio 7 : 8 : 9 at c_2^* . (c) The sail’s tuning curve for Sun–Earth: $c_2(\beta)$ sweeps continuously from the classical value down to unity, passing tidal parity and then c_2^* . Open markers give the three classical Sun–Earth values, none of which is at the bound. (d) Classical c_2 against μ for all three points; c_2^* is attained at L_2 for $\mu = 0.4802$.

$\beta = 0.05$). Absolute step sizes and amplitude bounds therefore mean different things at different β ; fixed ones admit far-field orbits, and a seed amplitude of $A_z = 0.02$ — reasonable for Earth–Moon — is roughly 2γ for Sun–Earth, larger than the entire distance from the equilibrium to the smaller primary.

Second, the differential corrector solves for $[x_0, \dot{y}_0, T/2]$ subject to $[\dot{x}_f, \dot{z}_f, y_f] = 0$; z_0 is not a free variable and A_z is not a constraint. Stepping A_z and re-solving therefore fails at folds in A_z , where the branch turns and no solution exists at the requested amplitude. We use pseudo-arclength continuation, appending the arclength condition and treating the continuation parameter as an unknown, so that folds are traversed rather than tripped over. Branch identity is additionally enforced across β : each β is seeded from its predecessor’s converged state with z_0 rescaled by $\gamma_{\text{new}}/\gamma_{\text{old}}$, and two guards verify the result — the sign of the z -extremum asymmetry

$$\Delta_z = \frac{|z_0| - |z(T/2)|}{|z_0| + |z(T/2)|} \quad (25)$$

must match the reference β , and the family must lie sunward of the equilibrium. Δ_z is $O(0.1)$ for a genuine halo and vanishes identically on the vertical-Lyapunov branch, which makes it an




Figure 4: Halo families across $\beta \in [0.001, 0.05]$, Sun–Earth, $\alpha = 0$, walked by pseudo-arclength continuation with branch identity enforced across β . **(a)** Family depth below the local equilibrium in the sail potential, against amplitude scaled by the local length γ . **(b)** Period. **(c)** Stability index, with the $|\nu| = 1$ boundary. **(d)** The (β, A_z) plane: family extent, folds, and linearly stable segments, with tidal parity marked.

effective discriminator. Without the cross- β guard one of the twelve families lands on a different branch.

Figure 4 reports the resulting atlas. The Jacobi constant is computed from (6); because the potential (5) is itself β -dependent, raw C_{sail} is not comparable across β , and we plot instead the depth below the local equilibrium, $C_{\text{sail}} - C_{\text{sail}}(x_{\text{eq}})$, against A_z/γ .

7 Control authority of a face-on sail

The dynamics above are open-loop. Since equilibria in this region are unstable, it is natural to ask what a sail can do about it, and here the face-on configuration turns out to be singular.

A sail has two control inputs, the attitude angles, and its thrust direction is rigidly coupled to its magnitude through (3). Linearising in (α, δ) about a nominal (α_0, δ_0) gives a 6×2 control Jacobian B_{sail} whose non-trivial rows are $\partial \mathbf{a}/\partial \alpha$ and $\partial \mathbf{a}/\partial \delta$. At $\alpha_0 = 0$ exactly,

$$\frac{\partial \mathbf{a}}{\partial \alpha} = \beta \frac{1 - \mu}{r_1^2} (\cos \delta_0 \hat{\mathbf{p}} + \sin \delta_0 \hat{\mathbf{q}}), \quad \frac{\partial \mathbf{a}}{\partial \delta} = \mathbf{0}. \quad (26)$$

The clock angle has *no* first-order authority when the sail is face-on: it only rotates a normal already parallel to $\hat{\mathbf{r}}$. The single available direction is therefore fixed by the constant δ_0 , and it




Figure 5: Collapse of the unstable eigenvalue with β . The trend is not sail stabilisation of a saddle: the standoff grows, the smaller primary’s tidal term dies, and the orbit moves away from the saddle it began near.

is purely transverse, since $\mathbf{a} \cdot \hat{\mathbf{r}} = \beta(1 - \mu)r_1^{-2} \cos^3 \alpha$ and $d(\cos^3 \alpha)/d\alpha \rightarrow 0$ as $\alpha \rightarrow 0$.

Rank of B is not the relevant test, because Coriolis coupling can carry a transverse input into the radial direction. Applying the Kalman rank condition to the linearised system at the $\beta = 0.05$ equilibrium ($c_2 = 1.409194$):

control model	Kalman rank	verdict
unconstrained three-axis thruster (6×3)	6/6	controllable
sail, $\alpha_0 = 0$, $\delta_0 = 0$	4/6	uncontrollable
sail, $\alpha_0 = 0$, $\delta_0 = \pi/2$	2/6	uncontrollable
sail, $\alpha_0 \geq 0.5^\circ$	6/6	controllable

The in-plane four-state block *is* controllable by a single transverse input (rank 4/4) — Coriolis does the work — so the absent radial authority is not by itself fatal. What fails is the out-of-plane mode: z decouples at linear order, and because $\partial \mathbf{a} / \partial \delta = \mathbf{0}$ the sail cannot redirect thrust into it without changing α_0 . A face-on sail reaches the in-plane pair or the vertical mode, never both:

$$\alpha_0 = 0 \text{ is a singular nominal for sail station-keeping.} \quad (27)$$

Any non-zero cone angle restores full rank; 0.5° suffices formally, though the second singular value there is 4.5×10^{-4} against 5.2×10^{-2} for the first, a 115:1 conditioning penalty, so usable authority requires a few degrees.

Authority is maximised at

$$\alpha_0^* = \arctan(1/\sqrt{2}) = 35.264^\circ, \quad \frac{\sigma_2}{\sigma_1} = \frac{1}{3}, \quad \cos^2 \alpha_0^* = \frac{2}{3}, \quad (28)$$

all exact and independent of β . This angle is the classical optimal sail cone angle for maximum transverse thrust [7, §2.6]; recovering it from the control Jacobian is a benchmark check on (26), not a new result. What is new is that the same angle maximises out-of-plane control authority, with σ_2/σ_1 exactly 1/3 there, and that the authority vanishes identically at $\alpha_0 = 0$. Figure 6 shows the trade: thrust falls as $\cos^2 \alpha_0$ while authority rises from zero, and α_0^* is where the second is maximal.

8 Where the threshold sits against flown hardware

The threshold (17) is only interesting if it is within reach. We therefore reduce β from primary mission specifications rather than quoting secondary values, using $\beta = \sigma^*/\sigma$ with the critical

Control authority of a solar sail at the artificial equilibrium ($\beta = 0.05$)




Figure 6: Control authority of a solar sail at the artificial equilibrium, $\beta = 0.05$. (a) Singular values of B_{sail} against nominal cone angle; $\sigma_2 \rightarrow 0$ as $\alpha_0 \rightarrow 0$. (b) Conditioning, maximal at α_0^* with the exact value $1/3$; the shaded strip at the origin is the uncontrollable nominal. (c) Out-of-plane authority scales linearly with β . (d) The design trade, both curves dimensionless: thrust $\cos^2 \alpha_0$ against normalised authority.

sail loading $\sigma^* = 2P/g_\odot$. Evaluated from CODATA and IAU constants, $\sigma^* = 1.5311 \text{ g m}^{-2}$, reproducing McInnes' 1.53 [7] to the three figures he quotes, which validates the chain.

No sail flown and deployed exceeds $\beta = 0.00613$, and the only measured value is IKAROS at $\beta = 0.00062$. Against $\beta_{\text{crit}} = 0.028646$:

reference	β	shortfall
best flown, perfect-reflector limit (LightSail-2)	0.009799	$2.9\times$
best flown, realistic (LightSail-2)	0.006131	$4.7\times$
only measured value (IKAROS)	0.000615	$47\times$
most ambitious funded design (Solar Cruiser, cancelled)	0.020236	$1.4\times$

Tidal parity is therefore a factor of 4.7 beyond the best sail flown and deployed — 2.9 if that sail is credited with a perfect reflector — and a factor of 1.4 beyond the most ambitious design funded to flight hardware. The defensible statement is that it lies at the edge of *near-term* rather than current capability. We note explicitly that a claim of the form “ $\beta \sim 0.01\text{--}0.05$ is current technology”, which appears informally, is not supported by any flown mission.

A consequence worth stating: at the best flown $\beta \approx 0.006$ the saddle strength is $s \approx 2.5$, so for every sail ever flown the collinear structure is fully intact and smaller-primary-dominated.

Table 3: Lightness number of every solar sail flown or funded to flight hardware, reduced from primary specifications. β_{ideal} is the perfect-reflector limit; β_{eff} applies the optical efficiency $\eta = 1.12/1.79 = 0.626$ measured on IKAROS. IKAROS is the only entry whose lightness number follows from a *measured* thrust and needs no optical assumption. Solar Cruiser has no published mass; its β_{ideal} is reduced instead from the published characteristic acceleration $\geq 0.12 \text{ mm s}^{-2}$, and is therefore a lower bound rather than a value obtainable from the area column.

mission	area [m ²]	mass [kg]	β_{ideal}	β_{eff}	status
IKAROS [10, 11]	196	307	0.000 98	0.000 62	flown 2010, measured
LightSail-2 [12]	32	5	0.009 80	0.006 13	flown 2019
NEA Scout [13]	86	14	0.009 41	0.005 88	sail not deployed, lost 2022
ACS3 [14]	80	16	0.007 66	0.004 79	flown 2024
Solar Cruiser	1653	—	0.020 24	—	cancelled 2022

The dissolution characterised here is a prediction about a regime not yet reached, not an account of observed behaviour.

9 Limitations

1. **Face-on only.** Everything except Section 7 assumes $\alpha = 0$. For $\alpha \neq 0$ the force is non-conservative, no Jacobi integral exists, and the effective-potential analysis used throughout fails. The steered case is open and is the natural sequel.
2. **Ideal specular reflector.** Membrane billow, finite slew rate, self-shadowing, and wavelength-dependent optics are not modelled. Dachwald *et al.* [15] argue persuasively that non-ideal optics matter more than order-of-magnitude estimates suggest; the IKAROS efficiency $\eta = 0.626$ in Table 3 is a single-mission proxy, not a general correction.
3. **No bifurcation is claimed.** $s > 0$ for all finite r_2 , so the saddle persists; c_2^* is a stationary point of a smooth function. Both thresholds are exact structural landmarks, and the resonance search of Section 5 is what licenses that restraint.
4. **Derived hardware figures.** Every β in Table 3 except IKAROS’ is reduced by us from published area and mass, not quoted by the mission teams. Where a team publishes its own lightness number, prefer it.
5. **Linear controllability only.** Section 7 is a statement about the linearisation at the equilibrium. Nonlinear or higher-order control strategies at $\alpha_0 = 0$ are not excluded; only first-order authority is.

10 Conclusions

A face-on solar sail rescales solar gravity rather than perturbing the collinear point, so the collinear structure does not shift with β — it decays. We have characterised that decay through the saddle strength $s = \mu/r_2^3$, which falls four decades between $\beta = 0$ and $\beta = 0.5$, and obtained two exact results.

The tidal-parity threshold admits the closed form $\beta_{\text{crit}} = 2\mu^{1/3} - \mu^{2/3}$, exact for every mass ratio and verified to machine precision across five decades of μ . For Sun–Earth it is 0.0286, corresponding to a standoff of $3^{1/3}r_H$. The scale $\mu^{1/3}$ is not itself new; the exact sail parameter required to reach it is.

The parameter A of the collinear linearisation is exactly Richardson’s second Legendre coefficient c_2 , and a sail’s effect is to detune it continuously. The frequency ratio at any collinear

equilibrium is confined to $\nu/\omega \in [2\sqrt{2}/3, 1)$ and attains its bound at $c_2 = 8/5$, where the linear rates stand in the ratio 7 : 8 : 9. Both are independent of μ . The near-equality of the two frequencies has been noted before [1]; the bound and its extremum make it exact. No low-order rational lies in the band, so no commensurability underwrites either threshold, and we present them as scale markers rather than transitions. The sail’s role is to tune c_2 within a given system, reaching values — including the bound, at $\beta = 0.0409$ for Sun–Earth — that that system’s classical collinear points do not offer.

Finally, the face-on nominal is singular for station-keeping: the clock angle has no first-order authority, the Kalman rank is 4/6, and a few degrees of cone angle are required to recover it. Authority peaks at the classical optimal cone angle $\arctan(1/\sqrt{2})$, where the conditioning is exactly 1/3.

Set against flown hardware, tidal parity is a factor 4.7 beyond the best deployed sail and 47 beyond the only measured lightness number. It is a near-term design boundary, not a present one.

Data and code availability

All numerical results, figures and the verification suite are reproducible from a single entry point. Every quantity quoted in this paper is regenerated by the pipeline and written to a machine-generated results file; no value is hand-transcribed.

References

- [1] Marta Ceccaroni, Alessandra Celletti, and Giuseppe Pucacco. Bifurcations thresholds of halo orbits. In *Astrodynamics Network AstroNet-II*, volume 44 of *Astrophysics and Space Science Proceedings*, pages 35–44. Springer, 2016. doi:[10.1007/978-3-319-23986-6_3](https://doi.org/10.1007/978-3-319-23986-6_3).
- [2] Colin R. McInnes, Alastair J. McDonald, John F. L. Simmons, and Euan W. MacDonald. Solar sail parking in restricted three-body systems. *Journal of Guidance, Control, and Dynamics*, 17(2):399–406, 1994. doi:[10.2514/3.21211](https://doi.org/10.2514/3.21211).
- [3] Hexi Baoyin and Colin R. McInnes. Solar sail halo orbits at the Sun–Earth artificial L_1 point. *Celestial Mechanics and Dynamical Astronomy*, 94(2):155–171, 2006. doi:[10.1007/s10569-005-4626-3](https://doi.org/10.1007/s10569-005-4626-3).
- [4] Thomas J. Waters and Colin R. McInnes. Periodic orbits above the ecliptic in the solar-sail restricted three-body problem. *Journal of Guidance, Control, and Dynamics*, 30(3):687–693, 2007. doi:[10.2514/1.26232](https://doi.org/10.2514/1.26232).
- [5] Ariadna Farrés and Àngel Jorba. Periodic and quasi-periodic motions of a solar sail close to SL_1 in the Earth–Sun system. *Celestial Mechanics and Dynamical Astronomy*, 107(1–2):233–253, 2010. doi:[10.1007/s10569-010-9268-4](https://doi.org/10.1007/s10569-010-9268-4).
- [6] Ariadna Farrés and Àngel Jorba. Station keeping strategies for a solar sail in the solar system. In *Advances in Astrodynamics and Space Applications*. Springer, 2016. doi:[10.1007/978-3-319-27464-5_3](https://doi.org/10.1007/978-3-319-27464-5_3).
- [7] Colin R. McInnes. *Solar Sailing: Technology, Dynamics and Mission Applications*. Springer–Praxis, Chichester, UK, 1999. doi:[10.1007/978-1-4471-3992-8](https://doi.org/10.1007/978-1-4471-3992-8).
- [8] David L. Richardson. Analytic construction of periodic orbits about the collinear points. *Celestial Mechanics*, 22(3):241–253, 1980. doi:[10.1007/BF01229511](https://doi.org/10.1007/BF01229511).

- [9] Gerard Gómez, Josep J. Masdemont, and Josep M. Mondelo. The dynamics around the collinear equilibrium points of the RTBP. *Physica D*, 157(4):283–321, 2001. doi:[10.1016/S0167-2789\(01\)00312-8](https://doi.org/10.1016/S0167-2789(01)00312-8).
- [10] Osamu Mori, Hirotaka Sawada, Ryu Funase, et al. First solar power sail demonstration by IKAROS. *Transactions of the Japan Society for Aeronautical and Space Sciences*, 8(ists27):To_4.25–To_4.31, 2010. doi:[10.2322/tastj.8.To_4.25](https://doi.org/10.2322/tastj.8.To_4.25).
- [11] Yuichi Tsuda, Osamu Mori, Ryu Funase, Hirotaka Sawada, Takayuki Yamamoto, Takanao Saiki, Tatsuya Endo, and Jun’ichiro Kawaguchi. Flight status of IKAROS deep space solar sail demonstrator. *Acta Astronautica*, 69(9–10):833–840, 2011. doi:[10.1016/j.actaastro.2011.06.005](https://doi.org/10.1016/j.actaastro.2011.06.005).
- [12] David A. Spencer, Bruce Betts, John M. Bellardo, Alex Diaz, Barbara Plante, and Justin R. Mansell. The LightSail 2 solar sailing technology demonstration. *Advances in Space Research*, 67(9):2878–2889, 2021. doi:[10.1016/j.asr.2020.06.029](https://doi.org/10.1016/j.asr.2020.06.029).
- [13] Leslie McNutt, Les Johnson, Dennon Clardy, Julie Castillo-Rogez, Andreas Frick, and Laura Jones. Near-Earth asteroid scout. *AIAA SPACE Conference Proceedings*, 2014. doi:[10.2514/6.2014-4435](https://doi.org/10.2514/6.2014-4435).
- [14] W. Keats Wilkie. The NASA advanced composite solar sail system (ACS3) flight demonstration. Technical Report NTRS 20210016824, NASA, 2021.
- [15] Bernd Dachwald, Malcolm Macdonald, Colin R. McInnes, Giovanni Mengali, and Alessandro A. Quarta. Solar sails are not ideal, and yes it matters. In *Advances in Solar Sailing*. Springer, 2013. doi:[10.1007/978-3-642-34907-2_55](https://doi.org/10.1007/978-3-642-34907-2_55).