

As a library, NLM provides access to scientific literature. Inclusion in an NLM database does not imply endorsement of, or agreement with, the contents by NLM or the National Institutes of Health.

Learn more: [PMC Disclaimer](#) | [PMC Copyright Notice](#)



*Cells*. 2025 Mar 12;14(6):417. doi: [10.3390/cells14060417](https://doi.org/10.3390/cells14060417)

## Effects of Space Flight on Inflammasome Activation in the Brain of Mice

[Upal Roy](#)<sup>1,\*</sup>, [Roey Hadad](#)<sup>2</sup>, [Angel A Rodriguez](#)<sup>1</sup>, [Alen Saju](#)<sup>1</sup>, [Deepa Roy](#)<sup>1</sup>, [Mario Gil](#)<sup>3</sup>, [Robert W Keane](#)<sup>2,4</sup>, [Ryan T Scott](#)<sup>5</sup>, [Xiao W Mao](#)<sup>6</sup>, [Juan Pablo de Rivero Vaccari](#)<sup>2,4,\*</sup>

Editor: John W Chen

[Author information](#) [Article notes](#) [Copyright and License information](#)

PMCID: PMC11941215 PMID: [40136666](#)

### Abstract

---

Space flight exposes astronauts to stressors that alter the immune response, rendering them vulnerable to infections and diseases. In this study, we aimed to determine the levels of inflammasome activation in the brains of mice that were housed in the International Space Station (ISS) for 37 days. C57BL/6 mice were launched to the ISS as part of NASA's Rodent Research 1 Mission on SpaceX-4 CRS-4 Dragon cargo spacecraft from 21 September 2014 to 25 October 2014. Dissected mouse brains from that mission were analyzed by immunoblotting of inflammasome signaling proteins and Electrochemiluminescence Immunoassay (ECLIA) for inflammatory cytokine levels. Our data indicate decreased inflammasome activation in the brains of mice that were housed in the ISS for 37 days when compared to the brains of mice that were maintained on the ground, and in mice corresponding to the baseline group that were sacrificed at the time of launching of SpaceX-4. Moreover, we did not detect any significant changes in the expression levels of the pro-inflammatory cytokines TNF- $\alpha$ , IL-2, IFN- $\gamma$ , IL-5, IL-6, IL-12p70 and IL-10 between the ground control and the flight groups. Together, these studies suggest that spaceflight results in a decrease in the levels of innate immune signaling molecules that govern inflammasome signaling in the brain of mice.

**Keywords:** inflammasome, inflammation, RR-1, space biology, International Space Station, innate immunity, inflammation

## 1. Introduction

---

Space flight exposes astronauts to a variety of stressors [1,2,3,4,5,6,7,8] ranging from psychological to physical that stem from the environment they experience [9]. These environmental factors include gravity changes, radiation exposure, altered nutrition or altered circadian rhythms, all of which affect the immune response [10,11,12,13,14]. Dysregulation of the immune response has two-fold consequences for astronauts. First, it has the potential to jeopardize the mission while in space, and it causes long-term detrimental consequences [8,15,16], not just while they are in space, but also once they return to Earth [8,17,18,19]. Therefore, it is imperative to gain a better understanding of how the immune response is altered by space flight.

Inflammasomes are supramolecular complexes comprising a sensor molecule such as NOD-like receptor 3 (NLRP3), the adaptor protein apoptosis-associated speck-like protein containing a caspase recruitment domain (ASC), and the cysteine aspartase caspase-1 [20]. The inflammasome plays a significant role in the regulation of the inflammatory response [21,22,23,24,25] by modulating the innate immune response through the activation of caspase-1, the processing of the pro-inflammatory cytokines interleukin (IL)-1 $\beta$  and IL-18, as well as the programmed cell death mechanism of pyroptosis [26]. The inflammasome has been shown to contribute to a variety of diseases including neurodegenerative diseases such as multiple sclerosis [27,28,29,30,31], Parkinson's disease [32,33], Alzheimer's disease [34,35], inflammaging [36,37] and metabolic syndrome [38,39]. Several of these diseases, with an inflammasome etiology, have also been suggested to be relevant in the context of space flight such as the development of dementia [40], altered aging and inflammation [41] or diabetes [42]. Hence, it is important to understand the effects of space flight on inflammasome signaling to develop therapies against specific targets of the immune response to protect and improve the performance of astronauts not only during a mission, but also their health on Earth [43,44].

It has been shown that space radiation affects hippocampus-dependent cognitive tasks [45], spatial navigation and episodic memory formation [46,47], which is very important for the overall health of astronauts [43,48,49,50,51,52,53,54,55,56,57]. In addition, it has been suggested that space radiation affects the cortex and causes altered cognitive flexibility and synaptic plasticity, which contribute to cognitive deficits [58]. A recent study used the anti-oxidant LGM2605 to inhibit the NLRP3 inflammasome in an in vitro model of lung vascular networks that were exposed to  $\gamma$ -rays and mixed-field radiation sources to mimic some of the space flight conditions in vitro [59]. The study found that 3–4 keV/ $\mu$ m and 8–10 keV/ $\mu$ m linear energy transfer (LET) proton radiation elevated the levels of NLRP3, and that this increase can be targeted with anti-oxidant approaches [59]. In this study, we aimed to characterize the effect of space flight on the expression of inflammasome proteins and inflammatory cytokines in mice that were part of NASA's Rodent Research 1 Mission (RR-1) on SpaceX-4 CRS-4 Dragon cargo spacecraft that spent 37 to 38 days at the International Space Station (ISS).

## 2. Materials and Methods


---

Animal procedures were approved by IACUC numbers NASA: NAS-13-002-Y1 and CASIS: CAS-13-001-Y1 for flight at the NASA Ames Research Center and the Kennedy Space Center in accordance with relevant guidelines and regulations.

### 2.1. Flight Animals

This animal study was approved by the NASA Ames Research Center and The Kennedy Space Center Institutional Animal Care and Use Committees. Brain tissue samples were obtained by the authors through an award from the NASA Biological Institutional Scientific Collection. C57BL/6 female mice (16-week-old, Jackson Labs) were flown to the ISS as part of RR-1 on SpaceX-4 CRS-4 Dragon cargo spacecraft [60]. During the flight, animals were kept at an ambient temperature of 26 to 28 °C in a 12 h light and dark cycle. Flight animals received water and a food bar diet specifically designed by NASA ad libitum. RR-1 mission animals were on board the Dragon capsule for 4 days (transit time) and then spent 37 days in the ISS ([Figure 1](#)).

Figure 1.



[Open in a new tab](#)

Study timeline.

RR-1 (<https://osdr.nasa.gov/bio/repo/data/payloads/RR-1> , accessed on 9 March 2025) lasted 37 days, and according to the NASA Open Science Data Repository's (OSDR) Environmental Data App, between 21 September and 23 September, these mice experienced temperature changes ranging from 18.3 °C to 26.69 °C. On 24 September, the temperature stabilized between 21 and 23 °C. Importantly, mice in the ISS (flight group) and in the ground control group experienced the same temperature conditions. Overall, the lowest temperature in the ISS during RR-1 was 18.50 °C and the highest temperature was 26.80 °C, with a mean temperature of 22.76 °C, whereas in the ground control group, the lowest temperature was 18.30 °C and the maximum was 27.10 °C (mean temperature 22.76 °C). Moreover,

telemetry data indicated that mice in the flight group experienced initial higher CO<sub>2</sub> levels at around 2621 ppm on 20 September, whereas mice in the ground group experienced much lower CO<sub>2</sub> levels at approximately 476 ppm. By 27 September, both groups of mice experienced similar CO<sub>2</sub> conditions through the rest of the flight, approximately between 1868 ppm and 4690 ppm. The minimum CO<sub>2</sub> level in the ISS group was 1862 ppm and the maximum was 4667 ppm (mean 3197.27, median 3192 ppm), whereas in the ground group, the minimum was 438 ppm, the maximum was 4690 ppm and the mean was 2853.74. The median was 3188 ppm. Moreover, by October 25th, mice in the ISS experienced an accumulated radiation dose of 7.17 mGy/day. In the ISS group, the minimum galactic cosmic radiation (GCR) dose was 0.13 mGy/day, and the maximum was 0.14 mGy/day (mean 0.13 mGy/day). Furthermore, the radiation experienced as a result of the South Atlantic Anomaly (SAA) ranged between 0.06 and 0.10 mGy/day (mean 0.07 mGy/day). Together, these accounted for a minimum total dose of 0.19 mGy/day, a maximum of 0.23 mGy/day and an average of 0.20 mGy/day.

## 2.2. Ground Control Animals

Animals were kept in similar conditions to flight animals (ambient temperature of 26 to 28 °C in a 12 h light and dark cycle). Telemetry data (with a 4-day delay) from the flight group were used to keep the ground control animals under similar conditions (housed in RR-1 flight hardware within an environmental simulator under the same conditions as the ISS). Water and food conditions for this group were the same as for the flight animal group as described.

## 2.3. Vivarium and Baseline Animals

Animals in the vivarium group were animals that were housed under standard vivarium conditions and remained alive for the duration of study. Animals in the baseline group consisted of mice that were euthanized at the time when the animals in the flight group were launched.

## 2.4. Brain Dissection

Within 4 h of splashdown, flight and ground control animals were sacrificed and brains were removed and stored at –80 °C until further analysis. For the study focusing on the forebrain, microdissections of the entire hippocampus and primary motor cortex were performed following well-established rodent brain microdissection protocols [[61,62,63](#)], and a brain atlas served as a reference to ensure accuracy of brain region identification [[64](#)]. Following brain dissection, samples were stored at –80 °C until further analysis. Tissues used in this study consisted of brain cortex, hippocampus and rest of the tissue from the brain areas are referred as “other than cortex and hippocampus”.

## 2.5. Immunoblotting

Analysis of inflammasome signaling proteins in brain protein lysates by immunoblotting was carried out as described in [65]. Accordingly, brain lysates (25 µg) were run and resolved in 4–20% Criterion TGX Stain-Free precast gels (Bio-Rad, Hercules, CA, USA) prior to transfer to PVDF membranes (Bio-Rad). Membranes were then incubated with primary antibodies against caspase-1 (Novus Biologicals Centennial, CO, USA), ASC (Santa Cruz Biotechnology, Inc., Dallas, TX, USA), and IL-1β (Cell Signaling, Technology, Inc., Danvers, MA, USA) at a dilution of 1:1000 for 1 h followed by HRP-linked secondary antibodies (1:1000) for 45 min. β-actin (1:5000, Sigma Aldrich, St. Louis, MO, USA) was used as a standard and protein loading control. Membranes were imaged with the ChemiDoc Touch Imaging System (Bio-Rad) following enhanced chemiluminescence. Densitometric analysis for relative protein quantification was carried out with UN-SCAN-IT gel 6.3 Software (Silk Scientific Corporation, Vineyard, UT, USA).

## 2.6. Electrochemiluminescence Immunoassay (ECLIA)

Using protein lysates, a V-PLEX ECLIA kit (MSD) was used to measure the inflammatory profile in the different brain sections as described in [66,67] using the MESO-QuickPlex SQ-120MM (MSD, Rockville, MD, USA). Briefly, to measure the protein levels of TNF-α, IL-2, IFN-γ, IL-5, IL-6, IL-12p70 and IL-10, after washing steps, the plate was incubated overnight with 50 µL of samples. The following day, after washing steps, samples were incubated in detection antibodies for 2 h followed by 3 more washing steps and the addition of the 2X Read Buffer (MSD, Rockville, MD, USA). After obtaining the quantity of protein in pg/mL for each analyte using the MESO-QuickPlex SQ-120MM, samples were then normalized to total protein concentration per sample.

## 2.7. Statistical Analyses

Statistical analyses were carried out with Prism 10.0 software (GraphPad Software, San Diego, CA, USA). Descriptive statistics were first calculated. Data were tested for normality with the Shapiro–Wilk Test. Comparisons between the flight and ground groups were carried out by a one-tail *t*-test when data met normality and by the Mann–Whitney test when data were not normally distributed.

# 3. Results


---

## 3.1. Levels of Inflammasome Signaling Molecules Are Decreased in the Brain of Mice That Experienced Spaceflight

It has been previously shown that space travel affects the immune response [11]. To test the effects of spaceflight on inflammasome activation in the brain, we carried out an immunoblot analysis of brain areas other than cortex and hippocampus from mice that were in the ISS for approximately 1 month (Figure 1). Our findings indicate that caspase-1 (95% CI = Ground [1.06–5.9], Flight [0.3–2.3]) (Figure 2A), ASC (Ground [1.3–7.6], Flight [0.6–3.5]) (Figure 2B) and

IL-1 $\beta$  (Ground [1.8–5.4], Flight [0.8–2.7]) (Figure 2C) were significantly decreased in the mice that were in the ISS when compared to the ground control group. In addition, the expression levels for all the inflammasome proteins analyzed in these studies in the vivarium group and those in the baseline group were similar to the spaceflight group (Figure S1). Together, these findings indicate that the expression levels of inflammasome signaling proteins in the brain of mice are lower in the mice that were in the ISS (flight) than the mice in the ground control group.

Figure 2.



[Open in a new tab](#)

Inflammasome signaling proteins in the brain of mice in the ISS. Immunoblot of inflammasome proteins for the expression of (A) caspase-1, (B) ASC and (C) IL-1 $\beta$  in an area of the brain of mice other than cortex and hippocampus after 37 days of spaceflight. Data were normalized to  $\beta$ -actin as a protein loading control. Data presented as box-plots with dots as all data points. \*  $p < 0.05$ . N = 5 per group.


### 3.2. Inflammasome Proteins Are Decreased in the Cortex and Hippocampus of Mice After Spaceflight

We then aimed to determine the levels of inflammasome signaling proteins present in the cortex (Figure 3A) and hippocampus (Figure 3B). Accordingly, immunoblot analysis of cortical and hippocampal protein lysates suggested that similar to our previously described findings (Figure 2), there was a trend towards decreased inflammasome activation in the cortex (Figure 3A) and the hippocampus (Figure 3B) of mice after spaceflight when compared to the ground control group. In addition, the brains of mice in the baseline group also had significantly lower levels of inflammasome activation in the cortex and hippocampus, which were similar to the protein expression levels in the flight group. Moreover, animals in the vivarium group presented much higher inflammasome protein levels than under baseline and



spaceflight conditions. These findings suggest that in the cortex and hippocampus, inflammasome activation is lower after spaceflight.

Figure 3.



[Open in a new tab](#)

Inflammasome signaling proteins in the (A) cortex and (B) hippocampus of mice in the ISS. Immunoblot of inflammasome signaling proteins caspase-1, ASC and IL-1 $\beta$  in the cortex of mice after 37 days of spaceflight. Viv = Vivarium, Gr = Ground, Bsl = Baseline and Flgt = Flight. Data were normalized to  $\beta$ -actin as a protein loading control. N = 2 per group.


### 3.3. Levels of Tnf- $\alpha$ and Il12p70 Are Decreased in the Brain of Mice After Spaceflight

Gene expression changes associated with inflammation have been described in the brains of mice who experienced spaceflight conditions and remained in the ISS for over 30 days [61]. To determine the protein levels of inflammatory cytokines in the brain of mice in the ground control and spaceflight conditions, we carried out an ECLIA to measure the protein levels of TNF- $\alpha$  (Figure 4A), IL-2 (Figure 4B), IFN- $\gamma$  (Figure 4C), IL-5 (Figure 4D), IL-6 (Figure 4E), IL-12p70 (Figure 4F) and IL-10 (Figure 4G) in brain areas other than the cortex and hippocampus. Accordingly, we found similar levels of these cytokines between the ground and flight groups (Figure 4). Moreover, there was a trend towards lower levels of TNF- $\alpha$  and IL-12p70 in the baseline group when compared to the other groups (Figure S2). Together, these trends suggest that these inflammatory cytokines were not altered in regions outside the cortex and hippocampus as a



result of mission RR1.

Figure 4.



[Open in a new tab](#)

Inflammatory cytokine profiles in the brains of mice in the ISS. Protein levels of inflammatory cytokines (**A**) TNF- $\alpha$ , (**B**) IL-2, (**C**) IFN- $\gamma$  (**D**) IL-5, (**E**) IL-6, (**F**) IL-12p70 and (**G**) IL-10 in an area of the brain of mice other than cortex and hippocampus after 37 days of spaceflight and measured by ECLIA. Data were normalized to total protein. Data presented as box-plots with dots as all data points. N = 5 per group.

### 3.4. Inflammatory Cytokine Levels in the Cortex and Hippocampus of Mice After Spaceflight

To preliminarily determine the inflammatory profile in the cortex ([Figure S3](#)) and hippocampus ([Figure S4](#)) of mice, we used ECLIA to measure the levels of TNF- $\alpha$ , IL-10, IFN- $\gamma$ , IL-5, IL-6 and IL-12p70 in protein lysates. However, due to the low sample size ( $N = 2$ ), we were unable to carry out any statistical analyses in these tissues for the expression of these cytokines, yet the preliminary findings suggest that in the brain, the inflammatory response mediated by different cytokines is different in the cortex ([Figure S3](#)) than in the hippocampus ([Figure S4](#)) as a result of space flight.

## 4. Discussion

---

Previous studies have shown that the adaptive immune response is compromised as a result of space flight missions, resulting in the reactivation of viruses [\[68\]](#). In addition, a previous study has shown that following 13 days of spaceflight on board of space shuttle mission STS-118, mice experienced a decrease in liver, spleen and thymus mass compared to the ground control group, which was consistent with the decreased number of lymphocytes, monocyte/macrophage and granulocytes [\[69\]](#). However, how space flight affects the inflammatory response mediated by the inflammasome in the brain of mice has yet to be fully understood. In this study, we determined the levels of the key inflammasome signaling proteins caspase-1, ASC and IL-1 $\beta$  in the brain of mice after a stay of 37 days on the ISS. In addition, we measured the levels of cytokines that contribute significantly to the inflammatory response in the brain such as TNF- $\alpha$ , IL-10, IFN- $\gamma$ , IL-5, IL-6 and IL-12p70 [\[70,71,72\]](#). We used immunoblotting and ECLIA approaches to determine the inflammatory protein profile in the brains of mice that were in the ISS as part of the RR-1 mission that was launched on SpaceX-4 CRS-4 on 21 September 2014 and returned to earth on October 25th of the same year. Consistent with results from the STS-118 study, here we demonstrate that conditions experienced during a mission to low Earth orbit result in a significant decrease in levels of inflammasome proteins in the brain.

Previous studies from RR-1 have shown that these mice experienced increased expression of genes associated with non-shivering thermogenesis in interscapular brown adipose tissue as well as increased expression of genes associated with glucose metabolism and adipogenesis in gonadal white adipose tissue [\[73\]](#), consistent with other studies that showed altered liver [\[74\]](#) and muscle metabolism in these mice, which suggested that these mice experienced muscle atrophy [\[75\]](#) and that dietary changes might have been needed as a countermeasure [\[76\]](#).

During the RR-1 mission, mice in the flight group presented higher levels of CO<sub>2</sub> than the ground control group (2621 ppm vs. 476 ppm). However, approximately a week later, mice in both groups experienced similar CO<sub>2</sub> conditions. Increased CO<sub>2</sub> levels in neutrophils for a couple of hours have been shown to increase IL-1 $\beta$  as well as the expression of NLRP3 and caspase-1 without affecting the levels of ASC [\[77\]](#). Thus, it is possible that the changes seen between the ground control group and the flight group are due to the fluctuations in the CO<sub>2</sub> levels. Furthermore, another variable

that needs to be taken into account is that there was a difference between the vivarium group and the ground control group, bringing up the possibility of an effect due to the habitat. Notably, the CO<sub>2</sub> levels in the vivarium group were the same as ambient air (~419 ppm), which were much lower than the ground group (median: 3188 ppm). In addition, some differences have been described between the vivarium and the animal enclosure module in which water and food consumption were higher in the animal enclosure module without a significant effect on overall body mass but some differences in the weight of certain organs were reported; however, that study did not report the weight of the brain [78]. Alternatively, taken together, it is possible that the effects seen in the flight group are in part due to the effects of the CO<sub>2</sub> levels in the brain and the stress associated with space flight. Therefore, part of the elevated levels seen in the ground control group compared to the vivarium group may be due in part to the habitat environment and the CO<sub>2</sub> levels. Importantly, the CO<sub>2</sub> levels between the ground and the flight groups eventually stabilized after 7 days (ISS median: 3192 ppm; ground median: 3188 ppm).

During the RR-1, mice in the flight group experienced an average of 0.20 mGy/day. Previous studies have shown that whole body exposure to higher doses from a <sup>137</sup>CsCl radiation source (0.5 to 4 Gy) resulted in a dose-dependent increase in inflammasome activation as determined by the cleavage of caspase-1 at 1 day after exposure, an effect that was not present at 1 and 4 h; moreover, caspase-1 cleavage was significantly decreased by day 14 in immune cells [79]. Following this radiation protocol, inflammasome activation was mainly identified in macrophages, dendritic cells, and natural killer cells [79]. In addition, UV radiation has been shown to also activate inflammasomes [80]. Furthermore, it has been shown that exposure of mice to silicon (0.5 Gy) or iron (0.15 Gy) ions results in increased cytokine production that is associated with infiltration of immune cells [81,82]. A recent study in a model of the lung vascular network exposed to low levels of radiation ranging from 0.25 to 1 Gy [83] showed increased expression of the inflammasome sensor NLRP3, which was decreased by treatment with LGM2605, a scavenger of radiation-induced reactive oxygen species (ROS) and active chlorine species (ACS), which is capable of reducing ionizing species due to radiation [59]. Similarly, in a study using a clinostat to model microgravity (10<sup>-3</sup> gravitational force) in human umbilical vein endothelial cells, the authors showed that microgravity also increased the expression of NLRP3 consistent with increased expression of IL-1β and increased stress in the endoplasmic reticulum [84]. Moreover, a recent study showed that 8 Gy of radiation can inhibit activation of the NLRP3 inflammasome in macrophages exposed to ATP and LPS by a mechanism that involves inhibiting TWIK2 expression [85]. However, whether the 7.17 mGy/day of accumulated radiation is also capable of causing lower levels of inflammasome protein expression experienced by mice in the flight group is yet to be tested.

In this study, we have shown that in the brain of mice who were in the ISS, there were lower levels of the inflammasome signaling proteins caspase-1, ASC and IL-1β when compared to the ground control group. Interestingly, the levels of these proteins were similar to those levels present in the brains of the animals corresponding to the baseline group, which were sacrificed at the time of launch, suggesting that the inflammasome proteins continued to increase over time in the ground control group; however, this increase in inflammasome protein expression was arrested or stalled in the ISS group. We have previously shown that inflammasome activation in the brain increases as part of the inflammaging

response [36,37,67]. In contrast to those inflammaging studies, the brain data from the mice in RR1 presented lower levels of these inflammasome proteins similar to those levels that were present in mice that were sacrificed at the time of launch. These data would suggest that the innate immune response mediated by the inflammasome is not weakened by space flight, but that it does not develop as it normally does on Earth. Consistent with these findings on decreased inflammation in the Central Nervous System (CNS), animals in RR-1 have also been shown to present younger epigenetic age in the retina than the ground control group [86]. Similarly, in the NASA Twin Study, the spaceflight twin presented a longer telomere length than his twin brother who remained on Earth, suggesting that in space, the aging process is slowed down [41]. Together, these findings support our hypothesis that spaceflight decreases brain inflammaging mediated by the inflammasome.

It is possible that the decrease in the levels of inflammasome signaling proteins after spaceflight may be the result of a protective attempt to dampen the innate immune response in the brain following the stress experienced in space in the ISS. Stalled replication forks have been previously shown to limit the innate immune response mediated by cGAS–STING following replication stress and in order to maintain genome stability [87]. Thus, it is possible that the stress of exposure associated with radiation and microgravity [88,89,90,91,92,93] in the ISS mice resulted in stalled replication forks responsible for regulating inflammasome expression in the brain, which ultimately is responsible for the levels of inflammasome activation in the brain similar to those seen in mice that were sacrificed at the time of launch (baseline group).

Furthermore, the inflammatory cytokines tested presented a similar pattern of expression between the baseline and flight groups. Moreover, our findings on the cortex and hippocampus suggest a differential expression of different cytokines in different parts of the brain, even with a very limited sample size. As anticipated, these findings suggest diverse roles in the inflammatory response mounted by different cytokines in the brain.

In general, we would have anticipated that the inflammasome would be increased in the ISS group when compared to the baseline group based on previous studies looking at the peripheral immune response as a result of space flight. However, in this study, we have measured the inflammatory response in the CNS, suggesting that the immune response mounted in the brain as a result of space flight differs between the CNS and the periphery, an effect that is perhaps due to the relatively immune-privileged status of the brain owing to the characteristics of the blood–brain barrier. Moreover, similar observations have been previously reported in which spaceflight conditions resulted in immune dysfunctions and dysregulated inflammation [94,95]. Future studies are needed to confirm these results in other missions. However, consistent with these results, a previous study in the brain of mice that were in the ISS for 35 days and were flown on board SpaceX-12 as part of RR9 showed downregulation of immune genes [61]. Furthermore, in this study, only female mice were used. Thus, future studies are needed to better understand the sex differences associated with space flight in the inflammatory response in the brain, considering that previous studies have shown that brain cells respond differently in males vs. females to space-related radiation conditions [96,97,98,99]. Another limitation of this study is the low sample size characteristic of this type of space biology studies. In addition, although the trends are similar across

different brain areas, it is not clear what areas of the brain are included in those that did not include the cortex and hippocampus. Moreover, since we have only analyzed samples from RR-1, we were not able to compare the results of RR-1 to other missions. Thus, we do not know if the results of this study could be generalized to space flight in general or if these findings are particular to events/conditions that took place during the RR-1 mission. Future, studies should aim at analyzing mouse brains from other missions with larger sample sizes, as well as studies with more accurately described dissected brain areas, and with considerations for sex differences. Furthermore, it would be important to compare these findings in the rodent brain to blood-based biomarkers obtained from astronauts in missions of different durations.

In conclusion, this study offers for the first time information on the effects of spaceflight on the innate immune response mediated by the inflammasome in the mouse brain after inhabiting the ISS for approximately one month. Thus, it is important to expand on these findings in the future to develop effective countermeasures against the detrimental effects associated with space flight.

## Acknowledgments

---

Figures created with BioRender.

## Supplementary Materials

---

The following supporting information can be downloaded at <https://www.mdpi.com/article/10.3390/cells14060417/s1> : Figure S1: Inflammasome proteins in the brain. Figure S2: Inflammatory cytokines in the brain. Figure S3: Inflammatory cytokine profile in the cortex of mice in the ISS; Figure S4: Inflammatory cytokine profile in the hippocampus of mice in the ISS.

[cells-14-00417-s001.zip](#) (930KB, zip)

## Author Contributions

---

Conceptualization, U.R. and J.P.d.R.V.; methodology, J.P.d.R.V., A.A.R., A.S. and M.G.; software, A.A.R., M.G. and J.P.d.R.V.; validation, J.P.d.R.V., D.R. and U.R.; formal analysis, J.P.d.R.V., R.W.K. and R.H.; investigation, J.P.d.R.V., R.W.K., R.H., A.A.R. and D.R.; resources, J.P.d.R.V., R.W.K., X.W.M., R.T.S. and U.R.; data curation, J.P.d.R.V., U.R. and R.T.S.; writing—original draft preparation, J.P.d.R.V.; writing—review and editing, all authors, visualization,

J.P.d.R.V. and U.R.; supervision, U.R., D.R and J.P.d.R.V.; project administration, J.P.d.R.V. and U.R.; funding acquisition, U.R. All authors have read and agreed to the published version of the manuscript.

## Institutional Review Board Statement

---

Animal procedures were approved by IACUC numbers NASA: NAS-13-002-Y1 and CASIS: CAS-13-001-Y1 for flight at the NASA Ames Research Center and the Kennedy Space Center in accordance with relevant guidelines and regulations.

## Informed Consent Statement

---

Not applicable.

## Data Availability Statement

---

Data produced for this publication were deposited in the NASA Open Science Data Repository (OSDR) under accession numbers OSD-738 (<https://doi.org/10.26030/y2bh-8d88>), OSD-751 (<https://doi.org/10.26030/wa2p-cn71>), and OSD-752 (<https://doi.org/10.26030/3zzw-1v11>). Data used in this paper were downloaded from NASA's Environmental Data Application (EDA) at the Open Science Data Repository (<https://visualization.osdr.nasa.gov/eda/>, accessed on 9 March 2025). Data were derived from the Rodent Research-1 (RR1) payload.

## Conflicts of Interest

---

J.P.d.R.V. and R.W.K. are co-founders and managing members of InflamaCORE and LLC and have licensed patents on inflammasome proteins as biomarkers of injury and disease as well as on targeting inflammasome proteins for therapeutic purposes. J.P.d.R.V. and R.W.K. are Scientific Advisory Board Members of ZyVersa Therapeutics.

## Funding Statement

---

This project was supported by an award from NASA (NASA-MUREP) and UTRGV internal funding to UR.

## Footnotes

---

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI

and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

## References

---

1. Buchheim J.I., Matzel S., Rykova M., Vassilieva G., Ponomarev S., Nichiporuk I., Horl M., Moser D., Biere K., Feuerecker M., et al. Stress Related Shift Toward Inflammaging in Cosmonauts After Long-Duration Space Flight. *Front. Physiol.* 2019;10:85. doi: 10.3389/fphys.2019.00085. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
2. Carpo N., Tran V., Biancotti J.C., Cepeda C., Espinosa-Jeffrey A. Space Flight Enhances Stress Pathways in Human Neural Stem Cells. *Biomolecules.* 2024;14:65. doi: 10.3390/biom14010065. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
3. Kvetnansky R., Ksinantova L., Koska J., Noskov V.B., Vigas M., Grigoriev A.I., Macho L. Effect of space flight and head-down bedrest on neuroendocrine response to metabolic stress in physically trained subjects. *J. Gravitational Physiol.* 2004;11:P57–P60. [[PubMed](#)] [[Google Scholar](#)]
4. Macho L., Koska J., Ksinantova L., Pacak K., Hoff T., Noskov V.B., Grigoriev A.I., Vigas M., Kvetnansky R. The response of endocrine system to stress loads during space flight in human subject. *Adv. Space Res.* 2003;31:1605–1610. doi: 10.1016/S0273-1177(03)00097-8. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
5. Rizzo A.M., Corsetto P.A., Montorfano G., Milani S., Zava S., Tavella S., Cancedda R., Berra B. Effects of long-term space flight on erythrocytes and oxidative stress of rodents. *PLoS ONE.* 2012;7:e32361. doi: 10.1371/journal.pone.0032361. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
6. Sonnenfeld G. Space flight, microgravity, stress, and immune responses. *Adv. Space Res.* 1999;23:1945–1953. doi: 10.1016/S0273-1177(99)00455-X. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
7. Stein T.P. Space flight and oxidative stress. *Nutrition.* 2002;18:867–871. doi: 10.1016/S0899-9007(02)00938-3. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
8. Verhaar A.P., Hoekstra E., Tjon A.S., Utomo W.K., Deuring J.J., Bakker E.R., Muncan V., Peppelenbosch M.P. Dichotomal effect of space flight-associated microgravity on stress-activated protein kinases in innate immunity. *Sci. Rep.* 2014;4:5468. doi: 10.1038/srep05468. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
9. Afshinnekoo E., Scott R.T., MacKay M.J., Pariset E., Cekanaviciute E., Barker R., Gilroy S., Hassane D., Smith S.M., Zwart S.R., et al. Fundamental Biological Features of Spaceflight: Advancing the Field to Enable Deep-Space Exploration. *Cell.* 2020;183:1162–1184. doi: 10.1016/j.cell.2020.10.050. [[DOI](#)] [[PMC free article](#)]



[article](#)] [[PubMed](#)] [[Google Scholar](#) ]

10. Chapes S.K., Woods K.M., Armstrong J.W. Ground-Based experiments complement microgravity flight opportunities in the investigation of the effects of space flight on the immune response: Is protein kinase C gravity sensitive? *Trans. Kans. Acad. Sci.* 1993;96:74–79. doi: 10.2307/3628318. [[DOI](#) ] [[PubMed](#)] [[Google Scholar](#) ]

11. Crucian B.E., Chouker A., Simpson R.J., Mehta S., Marshall G., Smith S.M., Zwart S.R., Heer M., Ponomarev S., Whitmire A., et al. Immune System Dysregulation During Spaceflight: Potential Countermeasures for Deep Space Exploration Missions. *Front. Immunol.* 2018;9:1437. doi: 10.3389/fimmu.2018.01437. [[DOI](#) ] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#) ]

12. Marcos A., Nova E., Montero A. Changes in the immune system are conditioned by nutrition. *Eur. J. Clin. Nutr.* 2003;57((Suppl. S1)):S66–S69. doi: 10.1038/sj.ejcn.1601819. [[DOI](#) ] [[PubMed](#)] [[Google Scholar](#) ]

13. Sephton S., Spiegel D. Circadian disruption in cancer: A neuroendocrine-immune pathway from stress to disease? *Brain Behav. Immun.* 2003;17:321–328. doi: 10.1016/S0889-1591(03)00078-3. [[DOI](#) ] [[PubMed](#)] [[Google Scholar](#) ]

14. Suman S., Fornace A.J., Jr. Countermeasure development against space radiation-induced gastrointestinal carcinogenesis: Current and future perspectives. *Life Sci. Space Res.* 2022;35:53–59. doi: 10.1016/j.lssr.2022.09.005. [[DOI](#) ] [[PubMed](#)] [[Google Scholar](#) ]

15. Ruff G.E. Experimental studies of stress in space flight. *Am. J. Psychiatry.* 1959;115:1109–1110. [[PubMed](#)] [[Google Scholar](#) ]

16. Tigranyan R.A., Macho L., Kvetnansky R., Nemeth S., Kalita N.F. Stress in space flight: Metabolic aspects. *Physiologist.* 1980;23:S45–S50. [[PubMed](#)] [[Google Scholar](#) ]

17. Cogoli A. The effect of space flight on human cellular immunity. *Environ. Med.* 1993;37:107–116. [[PubMed](#)] [[Google Scholar](#) ]

18. Mandel A.D., Balish E. Effect of space flight on cell-mediated immunity. *Aviat. Space Environ. Med.* 1977;48:1051–1057. [[PubMed](#)] [[Google Scholar](#) ]

19. Sonnenfeld G., Miller E.S. Space flight and humoral and cellular immunity of animals. *Physiologist.* 1993;36:S68–S70. [[PubMed](#)] [[Google Scholar](#) ]

20. de Rivero Vaccari J.P., Dietrich W.D., Keane R.W. Therapeutics targeting the inflammasome after central nervous system injury. *Transl. Res.* 2016;167:35–45. doi: 10.1016/j.trsl.2015.05.003. [[DOI](#) ] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#) ]

21. Lang Y., Chu F., Shen D., Zhang W., Zheng C., Zhu J., Cui L. Role of Inflammasomes in Neuroimmune and Neurodegenerative Diseases: A Systematic Review. *Mediat. Inflamm.* 2018;2018:1549549. doi: 10.1155/2018/1549549. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
22. Patel S. Inflammasomes, the cardinal pathology mediators are activated by pathogens, allergens and mutagens: A critical review with focus on NLRP3. *Biomed. Pharmacother.* 2017;92:819–825. doi: 10.1016/j.biopha.2017.05.126. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
23. Platnich J.M., Muruve D.A. NOD-like receptors and inflammasomes: A review of their canonical and non-canonical signaling pathways. *Arch. Biochem. Biophys.* 2019;670:4–14. doi: 10.1016/j.abb.2019.02.008. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
24. Singhal G., Jaehne E.J., Corrigan F., Toben C., Baune B.T. Inflammasomes in neuroinflammation and changes in brain function: A focused review. *Front. Neurosci.* 2014;8:315. doi: 10.3389/fnins.2014.00315. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
25. Yang C.A., Chiang B.L. Inflammasomes and human autoimmunity: A comprehensive review. *J. Autoimmun.* 2015;61:1–8. doi: 10.1016/j.jaut.2015.05.001. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
26. de Rivero Vaccari J.P., Dietrich W.D., Keane R.W. Activation and regulation of cellular inflammasomes: Gaps in our knowledge for central nervous system injury. *J. Cereb. Blood Flow Metab.* 2014;34:369–375. doi: 10.1038/jcbfm.2013.227. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
27. Brint A., Greene S., Fennig-Victor A.R., Wang S. Multiple sclerosis: The NLRP3 inflammasome, gasdermin D, and therapeutics. *Ann. Transl. Med.* 2024;12:62. doi: 10.21037/atm-23-1960. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
28. Fan H., Fu Q., Du G., Qin L., Shi X., Wang D., Yang Y. Microglial Mayhem NLRP3 Inflammasome's Role in Multiple Sclerosis Pathology. *CNS Neurosci. Ther.* 2024;30:e70135. doi: 10.1111/cns.70135. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
29. Malhotra S., Hurtado-Navarro L., Pappolla A., Villar L.M.M., Rio J., Montalban X., Pelegrin P., Comabella M. Increased NLRP3 Inflammasome Activation and Pyroptosis in Patients With Multiple Sclerosis With Fingolimod Treatment Failure. *Neurol. Neuroimmunol. Neuroinflammation.* 2023;10:e200100. doi: 10.1212/NXI.0000000000200100. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
30. Otalora-Alcaraz A., Reilly T., Oro-Nolla M., Sun M.C., Costelloe L., Kearney H., Patra P.H., Downer E.J. The NLRP3 inflammasome: A central player in multiple sclerosis. *Biochem. Pharmacol.* 2025;232:116667. doi: 10.1016/j.bcp.2024.116667. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
31. Shadab A., Abbasi-Kolli M., Saharkhiz M., Ahadi S.H., Shokouhi B., Nahand J.S. The interplay between

mitochondrial dysfunction and NLRP3 inflammasome in multiple sclerosis: Therapeutic implications and animal model studies. *Biomed. Pharmacother.* 2024;175:116673. doi: 10.1016/j.biopha.2024.116673. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

32. Cabrera Ranaldi E., Nuytemans K., Martinez A., Luca C.C., Keane R.W., de Rivero Vaccari J.P. Proof-of-Principle Study of Inflammasome Signaling Proteins as Diagnostic Biomarkers of the Inflammatory Response in Parkinson's Disease. *Pharmaceuticals.* 2023;16:883. doi: 10.3390/ph16060883. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

33. Su Q., Ng W.L., Goh S.Y., Gulam M.Y., Wang L.F., Tan E.K., Ahn M., Chao Y.X. Targeting the inflammasome in Parkinson's disease. *Front. Aging Neurosci.* 2022;14:957705. doi: 10.3389/fnagi.2022.957705. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

34. McManus R.M., Latz E. NLRP3 inflammasome signalling in Alzheimer's disease. *Neuropharmacology.* 2024;252:109941. doi: 10.1016/j.neuropharm.2024.109941. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

35. Vontell R.T., Gober R., Dallmeier J., Brzostowicki D., Barreda A., Blennow K., Zetterberg H., Kvartsberg H., Gultekin S.H., de Rivero Vaccari J.P., et al. Association of region-specific hippocampal reduction of neurogranin with inflammasome proteins in post mortem brains of Alzheimer's disease. *Alzheimers Dement.* 2024;10:e12444. doi: 10.1002/trc2.12444. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

36. Cyr B., Hadad R., Keane R.W., de Rivero Vaccari J.P. The Role of Non-canonical and Canonical Inflammasomes in Inflammaging. *Front. Mol. Neurosci.* 2022;15:774014. doi: 10.3389/fnmol.2022.774014. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

37. Mejias N.H., Martinez C.C., Stephens M.E., de Rivero Vaccari J.P. Contribution of the inflammasome to inflammaging. *J. Inflamm.* 2018;15:23. doi: 10.1186/s12950-018-0198-3. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

38. Lamkanfi M., Kanneganti T.D. The inflammasome: A remote control for metabolic syndrome. *Cell Res.* 2012;22:1095–1098. doi: 10.1038/cr.2012.55. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

39. Pahwa R., Singh A., Adams-Huet B., Devaraj S., Jialal I. Increased inflammasome activity in subcutaneous adipose tissue of patients with metabolic syndrome. *Diabetes Metab. Res. Rev.* 2021;37:e3383. doi: 10.1002/dmrr.3383. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

40. Miller K.B., Mi K.L., Nelson G.A., Norman R.B., Patel Z.S., Huff J.L. Ionizing radiation, cerebrovascular disease, and consequent dementia: A review and proposed framework relevant to space radiation exposure. *Front. Physiol.* 2022;13:1008640. doi: 10.3389/fphys.2022.1008640. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

41. Garrett-Bakelman F.E., Darshi M., Green S.J., Gur R.C., Lin L., Macias B.R., McKenna M.J., Meydan C., Mishra T., Nasrini J., et al. The NASA Twins Study: A multidimensional analysis of a year-long human spaceflight. *Science*. 2019;364:eaau8650. doi: 10.1126/science.aau8650. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
42. Tobin B.W., Uchakin P.N., Leeper-Woodford S.K. Insulin secretion and sensitivity in space flight: Diabetogenic effects. *Nutrition*. 2002;18:842–848. doi: 10.1016/S0899-9007(02)00940-1. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
43. Childress S.D., Williams T.C., Francisco D.R. NASA Space Flight Human-System Standard: Enabling human spaceflight missions by supporting astronaut health, safety, and performance. *NPJ Microgravity*. 2023;9:31. doi: 10.1038/s41526-023-00275-2. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
44. Smith S.M., McCoy T., Gazda D., Morgan J.L., Heer M., Zwart S.R. Space flight calcium: Implications for astronaut health, spacecraft operations, and Earth. *Nutrients*. 2012;4:2047–2068. doi: 10.3390/nu4122047. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
45. Yasuda H., Komiyama T., Fujitaka K. Probability of hippocampus cell hits by high-LET space radiation in a low-Earth-orbit mission (STS-91) *Phys. Med.* 2001;17((Suppl. S1)):166–169. [[PubMed](#)] [[Google Scholar](#)]
46. Robinson N.T.M., Descamps L.A.L., Russell L.E., Buchholz M.O., Bicknell B.A., Antonov G.K., Lau J.Y.N., Nutbrown R., Schmidt-Hieber C., Hausser M. Targeted Activation of Hippocampal Place Cells Drives Memory-Guided Spatial Behavior. *Cell*. 2020;183:1586–1599.e1510. doi: 10.1016/j.cell.2020.09.061. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
47. Tidmore A., Dutta S.M., Fesshaye A.S., Russell W.K., Duncan V.D., Britten R.A. Space Radiation-Induced Alterations in the Hippocampal Ubiquitin-Proteome System. *Int. J. Mol. Sci.* 2021;22:7713. doi: 10.3390/ijms22147713. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
48. Blaber E., Marcal H., Burns B.P. Bioastronautics: The influence of microgravity on astronaut health. *Astrobiology*. 2010;10:463–473. doi: 10.1089/ast.2009.0415. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
49. Bondar R.L. Astronaut aids health research. Interview by Marilyn Rubin and Sarah Gueldner. *Reflections*. 1995;21:12. [[PubMed](#)] [[Google Scholar](#)]
50. Chancellor J.C., Scott G.B., Sutton J.P. Space Radiation: The Number One Risk to Astronaut Health beyond Low Earth Orbit. *Life*. 2014;4:491–510. doi: 10.3390/life4030491. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
51. Green D.A. SpaceX must share astronaut health data to boost space biomedicine. *Nature*. 2024;634:782.

doi: 10.1038/d41586-024-03438-7. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

52. Hamm P.B., Nicogossian A.E., Pool S.L., Wear M.L., Billica R.D. Design and current status of the longitudinal study of astronaut health. *Aviat. Space Environ. Med.* 2000;71:564–570. [[PubMed](#)] [[Google Scholar](#)]

53. Oluwafemi F.A., Abdelbaki R., Lai J.C., Mora-Almanza J.G., Afolayan E.M. A review of astronaut mental health in manned missions: Potential interventions for cognitive and mental health challenges. *Life Sci. Space Res.* 2021;28:26–31. doi: 10.1016/j.lssr.2020.12.002. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

54. Peterson L.E., Pepper L.J., Hamm P.B., Gilbert S.L. Longitudinal study of astronaut health: Mortality in the years 1959–1991. *Radiat. Res.* 1993;133:257–264. doi: 10.2307/3578364. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

55. Smith K., Mercuri J. Microgravity and Radiation Effects on Astronaut Intervertebral Disc Health. *Aerosp. Med. Hum. Perform.* 2021;92:342–352. doi: 10.3357/AMHP.5713.2021. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

56. Vafapour H., Rafiepour P., Moradgholi J., Mortazavi S. Evaluating the biological impact of shelters on astronaut health during different solar particle events: A Geant4-DNA simulation study. *Radiat. Environ. Biophys.* 2025:1–14. doi: 10.1007/s00411-025-01111-9. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

57. Waisberg E., Ong J., Masalkhi M., Cools B., Lee R., Vinken M., Lee A.G. Hepatotoxicity in spaceflight: Intrinsic and extrinsic risks for astronaut metabolic health. *Ir. J. Med. Sci.* 2025:1–4. doi: 10.1007/s11845-025-03884-w. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

58. Kiffer F., Carr H., Groves T., Anderson J.E., Alexander T., Wang J., Seawright J.W., Sridharan V., Carter G., Boerma M., et al. Effects of (1)H + (16)O Charged Particle Irradiation on Short-Term Memory and Hippocampal Physiology in a Murine Model. *Radiat. Res.* 2018;189:53–63. doi: 10.1667/RR14843.1. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

59. Chatterjee S., Pietrofesa R.A., Park K., Tao J.Q., Carabe-Fernandez A., Berman A.T., Koumenis C., Sielecki T., Christofidou-Solomidou M. LGM2605 Reduces Space Radiation-Induced NLRP3 Inflammasome Activation and Damage in In Vitro Lung Vascular Networks. *Int. J. Mol. Sci.* 2019;20:176. doi: 10.3390/ijms20010176. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

60. Choi S.Y., Saravia-Butler A., Shirazi-Fard Y., Leveson-Gower D., Stodieck L.S., Cadena S.M., Beegle J., Solis S., Ronca A., Globus R.K. Validation of a New Rodent Experimental System to Investigate Consequences of Long Duration Space Habitation. *Sci. Rep.* 2020;10:2336. doi: 10.1038/s41598-020-58898-4. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

61. Holley J.M., Stanbouly S., Pecaut M.J., Willey J.S., Delp M., Mao X.W. Characterization of gene expression profiles in the mouse brain after 35 days of spaceflight mission. *NPJ Microgravity*. 2022;8:35. doi: 10.1038/s41526-022-00217-4. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
62. Aboghazleh R., Boyajian S.D., Atiyat A., Udwan M., Al-Helalat M., Al-Rashaideh R. Rodent brain extraction and dissection: A comprehensive approach. *MethodsX*. 2024;12:102516. doi: 10.1016/j.mex.2023.102516. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
63. Meyerhoff J., Muhie S., Chakraborty N., Naidu L., Sowe B., Hammamieh R., Jett M., Gautam A. Microdissection of Mouse Brain into Functionally and Anatomically Different Regions. *J. Vis. Exp.* 2021;168:e61941. doi: 10.3791/61941. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
64. Franklin K.B. The Mouse Brain: In Stereotaxic Coordinates. 3rd ed. Elsevier; Amsterdam, The Netherlands: 2007. [[Google Scholar](#)]
65. Cyr B., de Rivero Vaccari J.P. Methods to Study Inflammasome Activation in the Central Nervous System: Immunoblotting and Immunohistochemistry. *Methods Mol. Biol.* 2023;2696:223–238. doi: 10.1007/978-1-0716-3350-2\_15. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
66. Scott X.O., Chen S.H., Hadad R., Yavagal D., Peterson E.C., Starke R.M., Dietrich W.D., Keane R.W., de Rivero Vaccari J.P. Cohort study on the differential expression of inflammatory and angiogenic factors in thrombi, cerebral and peripheral plasma following acute large vessel occlusion stroke. *J. Cereb. Blood Flow Metab.* 2022;42:1827–1839. doi: 10.1177/0271678X221106956. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
67. Cyr B., de Rivero Vaccari J.P. Sex Differences in the Inflammatory Profile in the Brain of Young and Aged Mice. *Cells*. 2023;12:1372. doi: 10.3390/cells12101372. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
68. Crucian B., Stowe R.P., Mehta S., Quiriarte H., Pierson D., Sams C. Alterations in adaptive immunity persist during long-duration spaceflight. *NPJ Microgravity*. 2015;1:15013. doi: 10.1038/npjmgrav.2015.13. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
69. Baqai F.P., Gridley D.S., Slater J.M., Luo-Owen X., Stodieck L.S., Ferguson V., Chapes S.K., Pecaut M.J. Effects of spaceflight on innate immune function and antioxidant gene expression. *J. Appl. Physiol.* 2009;106:1935–1942. doi: 10.1152/japplphysiol.91361.2008. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
70. Carregosa D., Loncarevic-Vasiljkovic N., Feliciano R., Moura-Louro D., Mendes C.S., Dos Santos C.N. Locomotor and gait changes in the LPS model of neuroinflammation are correlated with inflammatory cytokines in blood and brain. *J. Inflamm.* 2024;21:39. doi: 10.1186/s12950-024-00412-y. [[DOI](#)] [[PMC free](#)]



[article](#)] [[PubMed](#)] [[Google Scholar](#) ]

71. Yang J., Ran M., Li H., Lin Y., Ma K., Yang Y., Fu X., Yang S. New insight into neurological degeneration: Inflammatory cytokines and blood-brain barrier. *Front. Mol. Neurosci.* 2022;15:1013933. doi: 10.3389/fnmol.2022.1013933. [[DOI](#) ] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#) ]

72. Zhou L., Li Q., Liu S., Wang L., Yu M., Lu X., Yang L., Lei W., Chen G. Association of inflammatory cytokines with magnetic resonance imaging features of the brain in patients with depression. *Brain Res. Bull.* 2024;219:111131. doi: 10.1016/j.brainresbull.2024.111131. [[DOI](#) ] [[PubMed](#)] [[Google Scholar](#) ]

73. Wong C.P., Iwaniec U.T., Turner R.T. Evidence for increased thermogenesis in female C57BL/6J mice housed aboard the international space station. *NPJ Microgravity.* 2021;7:23. doi: 10.1038/s41526-021-00150-y. [[DOI](#) ] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#) ]

74. Beheshti A., Chakravarty K., Fogle H., Fazelinia H., Silveira W.A.D., Boyko V., Polo S.L., Saravia-Butler A.M., Hardiman G., Taylor D., et al. Multi-omics analysis of multiple missions to space reveal a theme of lipid dysregulation in mouse liver. *Sci. Rep.* 2019;9:19195. doi: 10.1038/s41598-019-55869-2. [[DOI](#) ] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#) ]

75. Li K., Desai R., Scott R.T., Steele J.R., Machado M., Demharter S., Hoarfrost A., Braun J.L., Fajardo V.A., Sanders L.M., et al. Explainable machine learning identifies multi-omics signatures of muscle response to spaceflight in mice. *NPJ Microgravity.* 2023;9:90. doi: 10.1038/s41526-023-00337-5. [[DOI](#) ] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#) ]

76. Vitry G., Finch R., McStay G., Behesti A., Dejean S., Larose T., Wotring V., da Silveira W.A. Muscle atrophy phenotype gene expression during spaceflight is linked to a metabolic crosstalk in both the liver and the muscle in mice. *iScience.* 2022;25:105213. doi: 10.1016/j.isci.2022.105213. [[DOI](#) ] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#) ]

77. Thom S.R., Bhopale V.M., Hu J., Yang M. Increased carbon dioxide levels stimulate neutrophils to produce microparticles and activate the nucleotide-binding domain-like receptor 3 inflammasome. *Free Radic. Biol. Med.* 2017;106:406–416. doi: 10.1016/j.freeradbiomed.2017.03.005. [[DOI](#) ] [[PubMed](#)] [[Google Scholar](#) ]

78. Moyer E.L., Dumars P.M., Sun G.S., Martin K.J., Heathcote D.G., Boyle R.D., Skidmore M.G. Evaluation of rodent spaceflight in the NASA animal enclosure module for an extended operational period (up to 35 days) *NPJ Microgravity.* 2016;2:16002. doi: 10.1038/npjmgrav.2016.2. [[DOI](#) ] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#) ]

79. Stoecklein V.M., Osuka A., Ishikawa S., Lederer M.R., Wanke-Jellinek L., Lederer J.A. Radiation exposure induces inflammasome pathway activation in immune cells. *J. Immunol.* 2015;194:1178–1189. doi:



10.4049/jimmunol.1303051. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

80. Feldmeyer L., Keller M., Niklaus G., Hohl D., Werner S., Beer H.D. The inflammasome mediates UVB-induced activation and secretion of interleukin-1beta by keratinocytes. *Curr. Biol.* 2007;17:1140–1145. doi: 10.1016/j.cub.2007.05.074. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

81. Tungjai M., Whorton E.B., Rithidech K.N. Persistence of apoptosis and inflammatory responses in the heart and bone marrow of mice following whole-body exposure to (2)(8)Silicon ((2)(8)Si) ions. *Radiat. Environ. Biophys.* 2013;52:339–350. doi: 10.1007/s00411-013-0479-4. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

82. Yan X., Sasi S.P., Gee H., Lee J., Yang Y., Mehrzad R., Onufrak J., Song J., Enderling H., Agarwal A., et al. Cardiovascular risks associated with low dose ionizing particle radiation. *PLoS ONE*. 2014;9:e110269. doi: 10.1371/journal.pone.0110269. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

83. Zeitlin C., Hassler D.M., Cucinotta F.A., Ehresmann B., Wimmer-Schweingruber R.F., Brinza D.E., Kang S., Weigle G., Bottcher S., Bohm E., et al. Measurements of energetic particle radiation in transit to Mars on the Mars Science Laboratory. *Science*. 2013;340:1080–1084. doi: 10.1126/science.1235989. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

84. Jiang M., Wang H., Liu Z., Lin L., Wang L., Xie M., Li D., Zhang J., Zhang R. Endoplasmic reticulum stress-dependent activation of iNOS/NO-NF-kappaB signaling and NLRP3 inflammasome contributes to endothelial inflammation and apoptosis associated with microgravity. *FASEB J.* 2020;34:10835–10849. doi: 10.1096/fj.202000734R. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

85. Huang X., Niu M., Sun T., Li M., Jiang X., Duan H., Zhang T., Zhang J., Xie F., Song R., et al. X-ray irradiation reduces ATP-dependent activation of NLRP3 inflammasome by inhibiting TWIK2 activity in macrophages. *Immunol. Lett.* 2024;272:106967. doi: 10.1016/j.imlet.2024.106967. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

86. Chen Z., Stanbouly S., Nishiyama N.C., Chen X., Delp M.D., Qiu H., Mao X.W., Wang C. Spaceflight decelerates the epigenetic clock orchestrated with a global alteration in DNA methylome and transcriptome in the mouse retina. *Precis. Clin. Med.* 2021;4:93–108. doi: 10.1093/pcmedi/pbab012. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

87. Emam A., Wu X., Xu S., Wang L., Liu S., Wang B. Stalled replication fork protection limits cGAS-STING and P-body-dependent innate immune signalling. *Nat. Cell Biol.* 2022;24:1154–1164. doi: 10.1038/s41556-022-00950-8. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

88. Bellone J.A., Gifford P.S., Nishiyama N.C., Hartman R.E., Mao X.W. Long-term effects of simulated microgravity and/or chronic exposure to low-dose gamma radiation on behavior and blood-brain barrier

integrity. NPJ Microgravity. 2016;2:16019. doi: 10.1038/npjmgrav.2016.19. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

89. Pani G., Verslegers M., Quintens R., Samari N., de Saint-Georges L., van Oostveldt P., Baatout S., Benotmane M.A. Combined Exposure to Simulated Microgravity and Acute or Chronic Radiation Reduces Neuronal Network Integrity and Survival. PLoS ONE. 2016;11:e0155260. doi: 10.1371/journal.pone.0155260. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

90. Radstake W.E., Gautam K., Miranda S., Vermeesen R., Tabury K., Rehnberg E., Buset J., Janssen A., Leysen L., Neefs M., et al. The Effects of Combined Exposure to Simulated Microgravity, Ionizing Radiation, and Cortisol on the In Vitro Wound Healing Process. Cells. 2023;12:246. doi: 10.3390/cells12020246. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

91. Sanzari J.K., Romero-Weaver A.L., James G., Krigsfeld G., Lin L., Diffenderfer E.S., Kennedy A.R. Leukocyte activity is altered in a ground based murine model of microgravity and proton radiation exposure. PLoS ONE. 2013;8:e71757. doi: 10.1371/journal.pone.0071757. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

92. Shanmugarajan S., Zhang Y., Moreno-Villanueva M., Clanton R., Rohde L.H., Ramesh G.T., Sibonga J.D., Wu H. Combined Effects of Simulated Microgravity and Radiation Exposure on Osteoclast Cell Fusion. Int. J. Mol. Sci. 2017;18:2443. doi: 10.3390/ijms18112443. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

93. Yamanouchi S., Rhone J., Mao J.H., Fujiwara K., Saganti P.B., Takahashi A., Hada M. Simultaneous Exposure of Cultured Human Lymphoblastic Cells to Simulated Microgravity and Radiation Increases Chromosome Aberrations. Life. 2020;10:187. doi: 10.3390/life10090187. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

94. Hicks J., Olson M., Mitchell C., Juran C.M., Paul A.M. The Impact of Microgravity on Immunological States. Immunohorizons. 2023;7:670–682. doi: 10.4049/immunohorizons.2200063. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

95. Spatz J.M., Fulford M.H., Tsai A., Gaudilliere D., Hedou J., Ganio E., Angst M., Aghaeepour N., Gaudilliere B. Human immune system adaptations to simulated microgravity revealed by single-cell mass cytometry. Sci. Rep. 2021;11:11872. doi: 10.1038/s41598-021-90458-2. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

96. Krukowski K., Grue K., Frias E.S., Pietrykowski J., Jones T., Nelson G., Rosi S. Female mice are protected from space radiation-induced maladaptive responses. Brain Behav. Immun. 2018;74:106–120. doi: 10.1016/j.bbi.2018.08.008. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

97. Liu B., Hinshaw R.G., Le K.X., Park M.A., Wang S., Belanger A.P., Dubey S., Frost J.L., Shi Q., Holton P., et al. Space-like (56)Fe irradiation manifests mild, early sex-specific behavioral and neuropathological changes in wildtype and Alzheimer's-like transgenic mice. *Sci. Rep.* 2019;9:12118. doi: 10.1038/s41598-019-48615-1. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
98. Raber J., Yamazaki J., Torres E.R.S., Kirchoff N., Stagaman K., Sharpton T., Turker M.S., Kronenberg A. Combined Effects of Three High-Energy Charged Particle Beams Important for Space Flight on Brain, Behavioral and Cognitive Endpoints in B6D2F1 Female and Male Mice. *Front. Physiol.* 2019;10:179. doi: 10.3389/fphys.2019.00179. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
99. Rola R., Fishman K., Baure J., Rosi S., Lamborn K.R., Obenaus A., Nelson G.A., Fike J.R. Hippocampal neurogenesis and neuroinflammation after cranial irradiation with (56)Fe particles. *Radiat. Res.* 2008;169:626–632. doi: 10.1667/RR1263.1. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

## Associated Data

---

*This section collects any data citations, data availability statements, or supplementary materials included in this article.*

## Supplementary Materials

[cells-14-00417-s001.zip](#) (930KB, zip)

## Data Availability Statement

Data produced for this publication were deposited in the NASA Open Science Data Repository (OSDR) under accession numbers OSD-738 (<https://doi.org/10.26030/y2bh-8d88>), OSD-751 (<https://doi.org/10.26030/wa2p-cn71>), and OSD-752 (<https://doi.org/10.26030/3zzw-1v11>). Data used in this paper were downloaded from NASA's Environmental Data Application (EDA) at the Open Science Data Repository (<https://visualization.osdr.nasa.gov/eda/>, accessed on 9 March 2025). Data were derived from the Rodent Research-1 (RR1) payload.

