## Problem 1

x-axis:  $[0,L_x]$  in m-1 subintervals of length h ,  $0=x_1 < x_2 < ... < x_{m-1} < x_m = L_x$  y-axis:  $[0,L_y]$  in n-1 subintervals of length k ,  $0=y_1 < y_2 < ... < y_{n-1} < y_n = L_y$  Robin boundary method


$$\frac{\delta^2 u}{\delta x^2} + \frac{\delta^2 u}{\delta y^2} = \frac{2H}{K\delta} u \qquad , \qquad \int (x,y) = \frac{2H}{K\delta} u$$

## 1 Discretize the equation

$$\frac{\partial^{2} u}{\partial x^{2}} \approx \frac{u(x+h,y) - 2u(x,y) + u(x-h,y)}{h^{2}} \longrightarrow \frac{u_{i+1,j} - 2u_{i,j} + u_{i-1,j}}{h^{2}}$$

 $\frac{\partial^{2} u}{\partial y^{2}} \approx \frac{u(x, y+h) - 2u(x, y) + u(x, y+h)}{k^{2}} \longrightarrow \frac{u_{i,j+1} - 2u_{i,j} + u_{i,j-1}}{k^{2}}$   $Eq: \frac{u_{i+1,j} - 2u_{i,j} + u_{i-1,j}}{h^{2}} + \frac{u_{i,j+1} - 2u_{i,j} + u_{i,j-1}}{k^{2}} = \frac{2H}{K\delta} U_{i,j}$ 

works on the inner points 2 = i = m-1 2 = j = n-1



## Depends on L

$$\frac{u_{i,1,j}}{h^2} - \frac{2u_{i,j}}{h^2} + \frac{u_{i-1,j}}{h^2} + \frac{u_{i,j+1}}{k^2} - \frac{2u_{i,j}}{k^2} + \frac{u_{i,j-1}}{k^2} - \frac{2Hu_{i,j}}{K\delta} = 0$$

$$\frac{1}{h^2} u_{inj} - \frac{2}{h^2} u_{i,j} + \frac{1}{h^2} u_{i-1,j} + \frac{1}{k^2} u_{i,j+1} - \frac{2}{k^2} u_{i,j} + \frac{1}{k^2} u_{i,j-1} - \frac{2H}{K8} u_{i,j} = 0$$

$$\frac{1}{h^2} u_{i+1,j} - \frac{1}{h^2} u_{i-1,j} + \frac{1}{k^2} u_{i,j+1} + \frac{1}{k^2} u_{i,j-1} - \left(\frac{2}{h^2} + \frac{2}{k^2} + \frac{2H}{K\delta}\right) u_{i,j} = 0$$

$$\frac{u_{i,1,j} - u_{i-1,j}}{h^2} + \frac{u_{i,j+1} + u_{i,j-1}}{k^2} - \left(\frac{2}{h^2} + \frac{2}{k^2} + \frac{2H}{K\delta}\right)u_{i,j} = 0$$

works on the inner points 25 j ≤ n-1

Bottom: 
$$\frac{\partial u}{\partial n} = -\frac{\partial u}{\partial u}$$
  
Top:  $\frac{\partial u}{\partial n} = \frac{\partial u}{\partial v}$   
Left:  $\frac{\partial u}{\partial n} = -\frac{\partial u}{\partial x}$   
Right:  $\frac{\partial u}{\partial n} = \frac{\partial u}{\partial x}$   
Heat BC (left):  $\frac{\partial u}{\partial n} = -\frac{\partial u}{\partial x} = \frac{P}{L\delta K}$ 

Right: 
$$\frac{\partial u}{\partial n} = \frac{\partial u}{\partial x}$$

Heat BC (left): 
$$\frac{\partial u}{\partial n} = -\frac{\partial u}{\partial x} = \frac{P}{L\delta K}$$

$$-\frac{2kH}{K}u_{i,j} = -3u_{i,j} + 4u_{i,j+1} - u_{i,j+2}$$

$$0 = \left(\frac{2kH}{K} - 3\right)u_{i,j} + 4u_{i,j+1} - u_{i,j+2}$$

$$0 = \left(\frac{2kH}{K} - 3\right) u_{i,1} + 4 u_{i,2} - u_{i,3}$$

Top: 
$$\frac{H}{K}u_{i,j} \approx \frac{-3u_{i,j} + 4u_{i,j-1} - u_{i,j-2}}{-2k}$$

$$0 = \left(\frac{2kH}{K} - 3\right) u_{i,j} + 4u_{i,j-1} - u_{i,j-2}$$

$$0 = \left(\frac{2kH}{K} - 3\right) u_{i,n} + 4u_{i,n-1} - u_{i,n-2}$$

$$-\frac{2hH}{K}uij = -3uij + 4uinj - uinij$$

$$0 = \left(\frac{2hH}{K} - 3\right)u_{i,j} + 4u_{i+1,j} - u_{i+2,j}$$

Right: 
$$\frac{H}{K}$$
 uij  $\approx \frac{-3ui,j + 4ui-1,j - ui-2,j}{-2h}$ 

$$0 = \left(\frac{2hH}{K} - 3\right) u_{m,j} + 4 u_{m-1,j} - u_{m-2,j}$$