

# BEZIER SURFACES

**Rodrigo Silveira**

Curve and Surface Design  
Facultat d'Informàtica de Barcelona  
Universitat Politècnica de Catalunya

# INTRO TO BEZIER SURFACES

# INTRO TO BEZIER SURFACES

Idea: use grid of control points

# INTRO TO BEZIER SURFACES


Idea: use grid of control points

Consider  $(n + 1) \times (m + 1)$  control points arranged in a rectangular grid

# INTRO TO BEZIER SURFACES

Idea: use grid of control points


Consider  $(n + 1) \times (m + 1)$  control points arranged in a rectangular grid



# INTRO TO BEZIER SURFACES

Idea: use grid of control points


Consider  $(n + 1) \times (m + 1)$  control points arranged in a rectangular grid



# INTRO TO BEZIER SURFACES

Idea: use grid of control points


Consider  $(n + 1) \times (m + 1)$  control points arranged in a rectangular grid



# INTRO TO BEZIER SURFACES

Idea: use grid of control points

Consider  $(n + 1) \times (m + 1)$  control points arranged in a rectangular grid



$$S(u, v) = \sum_{i=0}^m \sum_{j=0}^n B_{m,i}(u) B_{n,j}(v) P_{i,j}$$

$$0 \leq u, v \leq 1$$

where the terms  $B_{m,i}$  and  $B_{n,j}$  are the Bernstein polynomials, same as in Bézier curves



# INTRO TO BEZIER SURFACES

Idea: use grid of control points

Consider a grid of  $(m + 1) \times (n + 1)$  control points arranged in a rectangular grid


$$S(u, v) = \sum_{i=0}^m \sum_{j=0}^n B_{m,i}(u) B_{n,j}(v) P_{i,j} \quad 0 \leq u, v \leq 1$$

In matrix form:

$$S(u, v) = (B_{m,0}(u), B_{m,1}(u), \dots, B_{m,m}(u)) \begin{pmatrix} P_{0,0} & P_{0,1} & \dots & P_{0,n} \\ P_{1,0} & P_{1,1} & \dots & P_{1,n} \\ \vdots & \vdots & \ddots & \vdots \\ P_{m,0} & P_{m,1} & \dots & P_{m,n} \end{pmatrix} \begin{pmatrix} B_{n,0}(v) \\ B_{n,1}(v) \\ \vdots \\ B_{n,n}(v) \end{pmatrix}$$

# BEZIER SURFACES


## Example



# BEZIER SURFACES

## Example



$(0,2,0)$	$(1,2,1)$	$(2,2,0)$
$(0,1,1)$	$(1,1,4)$	$(2,1,1)$
$(0,0,0)$	$(1,0,1)$	$(2,0,0)$



Biquadratic Bézier surface patch [Salomon, Fig 6.20]

# BEZIER SURFACES

## Example



Biquadratic Bézier surface patch [Salomon, Fig 6.20]

# PROPERTIES OF BEZIER SURFACES

Properties of Bézier surface (on rectangular grid)  $0 \leq u, v \leq 1$


$$S(u, v) = \sum_{i=0}^m \sum_{j=0}^n B_{m,i}(u) B_{n,j}(v) P_{i,j}$$

# PROPERTIES OF BEZIER SURFACES

## Properties of Bézier surface (on rectangular grid) $0 \leq u, v \leq 1$

- Endpoints (patch corners)

$$S(u, v) = \sum_{i=0}^m \sum_{j=0}^n B_{m,i}(u) B_{n,j}(v) P_{i,j}$$



# PROPERTIES OF BEZIER SURFACES


## Properties of Bézier surface (on rectangular grid) $0 \leq u, v \leq 1$

- Endpoints (patch corners)

$$S(u, v) = \sum_{i=0}^m \sum_{j=0}^n B_{m,i}(u) B_{n,j}(v) P_{i,j}$$

Recall:  $S(u, v) = \sum_{i=0}^m \sum_{j=0}^n \binom{m}{i} \binom{n}{j} u^i (1-u)^{m-i} v^j (1-v)^{n-j} P_{i,j}$  (And  $0^0 = 1$ !)

$S(0, 0) = P_{0,0}$ ,  $S(1, 0) = P_{m,0}$ ,  $S(0, 1) = P_{0,n}$  and  $S(1, 1) = P_{m,n}$



# PROPERTIES OF BEZIER SURFACES

## Properties of Bézier surface (on rectangular grid) $0 \leq u, v \leq 1$


- Endpoints (patch corners)

$$S(u, v) = \sum_{i=0}^m \sum_{j=0}^n B_{m,i}(u) B_{n,j}(v) P_{i,j}$$

Recall:  $S(u, v) = \sum_{i=0}^m \sum_{j=0}^n \binom{m}{i} \binom{n}{j} u^i (1-u)^{m-i} v^j (1-v)^{n-j} P_{i,j}$  (And  $0^0 = 1$ !)

$S(0, 0) = P_{0,0}$ ,  $S(1, 0) = P_{m,0}$ ,  $S(0, 1) = P_{0,n}$  and  $S(1, 1) = P_{m,n}$

- Boundary curves





# PROPERTIES OF BEZIER SURFACES

## Properties of Bézier surface (on rectangular grid) $0 \leq u, v \leq 1$

- Endpoints (patch corners)



$$S(u, v) = \sum_{i=0}^m \sum_{j=0}^n B_{m,i}(u) B_{n,j}(v) P_{i,j}$$

Recall:  $S(u, v) = \sum_{i=0}^m \sum_{j=0}^n \binom{m}{i} \binom{n}{j} u^i (1-u)^{m-i} v^j (1-v)^{n-j} P_{i,j}$  (And  $0^0 = 1!$ )

$S(0, 0) = P_{0,0}$ ,  $S(1, 0) = P_{m,0}$ ,  $S(0, 1) = P_{0,n}$  and  $S(1, 1) = P_{m,n}$

- Boundary curves

$S(u, 0)$  is the Bézier curve defined by  $P_{0,0}, P_{1,0}, \dots, P_{n,0}$



# PROPERTIES OF BEZIER SURFACES

## Properties of Bézier surface (on rectangular grid) $0 \leq u, v \leq 1$

- Endpoints (patch corners)

$$S(u, v) = \sum_{i=0}^m \sum_{j=0}^n B_{m,i}(u) B_{n,j}(v) P_{i,j}$$

Recall:  $S(u, v) = \sum_{i=0}^m \sum_{j=0}^n \binom{m}{i} \binom{n}{j} u^i (1-u)^{m-i} v^j (1-v)^{n-j} P_{i,j}$  (And  $0^0 = 1!$ )

$S(0, 0) = P_{0,0}$ ,  $S(1, 0) = P_{m,0}$ ,  $S(0, 1) = P_{0,n}$  and  $S(1, 1) = P_{m,n}$



- Boundary curves

$S(u, 0)$  is the Bézier curve defined by  $P_{0,0}, P_{1,0}, \dots, P_{n,0}$

$S(u, 1)$  is the Bézier curve defined by  $P_{0,n}, P_{1,n}, \dots, P_{m,n}$

$S(v, 0)$  is the Bézier curve defined by  $P_{0,0}, P_{0,1}, \dots, P_{0,n}$

$S(1, v)$  is the Bézier curve defined by  $P_{m,0}, P_{m,1}, \dots, P_{m,n}$



# PROPERTIES OF BEZIER SURFACES

## Properties of Bézier surface (on rectangular grid)

- Uniparametric curves  
(i.e., fixed  $u$  or fixed  $v$ )

$$S(u, v) = \sum_{i=0}^m \sum_{j=0}^n B_{m,i}(u) B_{n,j}(v) P_{i,j}$$


# PROPERTIES OF BEZIER SURFACES

## Properties of Bézier surface (on rectangular grid)

- Uniparametric curves  
(i.e., fixed  $u$  or fixed  $v$ )

$$S(u, v) = \sum_{i=0}^m \sum_{j=0}^n B_{m,i}(u) B_{n,j}(v) P_{i,j}$$

Consider the curve obtained for a fix parameter, e.g.,  $u = u_0$



# PROPERTIES OF BEZIER SURFACES

## Properties of Bézier surface (on rectangular grid)


- Uniparametric curves  
(i.e., fixed  $u$  or fixed  $v$ )

$$S(u, v) = \sum_{i=0}^m \sum_{j=0}^n B_{m,i}(u) B_{n,j}(v) P_{i,j}$$

Consider the curve obtained for a fix parameter, e.g.,  $u = u_0$

Then we have:

$$\begin{aligned} S(u_0, v) &= \sum_{i=0}^m \sum_{j=0}^n B_{m,i}(u_0) B_{n,j}(v) P_{i,j} \\ &= \sum_{j=0}^n B_{n,j}(v) \left( \sum_{i=0}^m B_{m,i}(u_0) P_{i,j} \right) \end{aligned}$$



# PROPERTIES OF BEZIER SURFACES

## Properties of Bézier surface (on rectangular grid)


- Uniparametric curves  
(i.e., fixed  $u$  or fixed  $v$ )

$$S(u, v) = \sum_{i=0}^m \sum_{j=0}^n B_{m,i}(u) B_{n,j}(v) P_{i,j}$$

Consider the curve obtained for a fix parameter, e.g.,  $u = u_0$

Then we have:

$$\begin{aligned} S(u_0, v) &= \sum_{i=0}^m \sum_{j=0}^n B_{m,i}(u_0) B_{n,j}(v) P_{i,j} \\ &= \sum_{j=0}^n B_{n,j}(v) \underbrace{\left( \sum_{i=0}^m B_{m,i}(u_0) P_{i,j} \right)}_{\text{new control point } Q_j} \end{aligned}$$



# PROPERTIES OF BEZIER SURFACES

## Properties of Bézier surface (on rectangular grid)

- Uniparametric curves  
(i.e., fixed  $u$  or fixed  $v$ )

$$S(u, v) = \sum_{i=0}^m \sum_{j=0}^n B_{m,i}(u) B_{n,j}(v) P_{i,j}$$

Consider the curve obtained for a fix parameter, e.g.,  $u = u_0$


Then we have:

$$\begin{aligned} S(u_0, v) &= \sum_{i=0}^m \sum_{j=0}^n B_{m,i}(u_0) B_{n,j}(v) P_{i,j} \\ &= \sum_{j=0}^n B_{n,j}(v) \underbrace{\left( \sum_{i=0}^m B_{m,i}(u_0) P_{i,j} \right)}_{\text{new control point } Q_j} \end{aligned}$$

Example on the right, for  $u = 0.25$ ,  $n = m = 2$

$$S(0.25, v) = \sum_{j=0}^2 B_{n,j}(v) \left( \sum_{i=0}^2 B_{m,i}(u_0) P_{i,j} \right)$$

$$S(0.25, v) = \sum_{j=0}^2 B_{n,j}(v) \underbrace{\left( B_{m,0}(0.25) P_{0,j} + B_{m,1}(0.25) P_{1,j} + B_{m,2}(0.25) P_{2,j} \right)}_{\text{new control points, for } j = 0, 1, 2 \text{ (i.e., } Q_0, Q_1, Q_2)}$$



# PROPERTIES OF BEZIER SURFACES

## Properties of Bézier surface (on rectangular grid)

- Uniparametric curves  
(i.e., fixed  $u$  or fixed  $v$ )

$$S(u, v) = \sum_{i=0}^m \sum_{j=0}^n B_{m,i}(u) B_{n,j}(v) P_{i,j}$$

Consider the curve obtained for a fix parameter, e.g.,  $u = u_0$


Then we have:

$$\begin{aligned} S(u_0, v) &= \sum_{i=0}^m \sum_{j=0}^n B_{m,i}(u_0) B_{n,j}(v) P_{i,j} \\ &= \sum_{j=0}^n B_{n,j}(v) \underbrace{\left( \sum_{i=0}^m B_{m,i}(u_0) P_{i,j} \right)}_{\text{new control point } Q_j} \end{aligned}$$

Example on the right, for  $u = 0.25$ ,  $n = m = 2$

$$S(0.25, v) = \sum_{j=0}^2 B_{n,j}(v) \left( \sum_{i=0}^2 B_{m,i}(u_0) P_{i,j} \right)$$

$$S(0.25, v) = \sum_{j=0}^2 B_{n,j}(v) \underbrace{\left( B_{m,0}(0.25) P_{0,j} + B_{m,1}(0.25) P_{1,j} + B_{m,2}(0.25) P_{2,j} \right)}_{\text{new control points, for } j = 0, 1, 2 \text{ (i.e., } Q_0, Q_1, Q_2)}$$





# PROPERTIES OF BEZIER SURFACES

## Properties of Bézier surface (on rectangular grid)

- Uniparametric curves  
(i.e., fixed  $u$  or fixed  $v$ )

$$S(u, v) = \sum_{i=0}^m \sum_{j=0}^n B_{m,i}(u) B_{n,j}(v) P_{i,j}$$

Consider the curve obtained for a fix parameter, e.g.,  $u = u_0$


Then we have:

$$\begin{aligned} S(u_0, v) &= \sum_{i=0}^m \sum_{j=0}^n B_{m,i}(u_0) B_{n,j}(v) P_{i,j} \\ &= \sum_{j=0}^n B_{n,j}(v) \underbrace{\left( \sum_{i=0}^m B_{m,i}(u_0) P_{i,j} \right)}_{\text{new control point } Q_j} \end{aligned}$$

Example on the right, for  $u = 0.25$ ,  $n = m = 2$

$$S(0.25, v) = \sum_{j=0}^2 B_{n,j}(v) \left( \sum_{i=0}^2 B_{m,i}(u_0) P_{i,j} \right)$$

$$S(0.25, v) = \sum_{j=0}^2 B_{n,j}(v) \underbrace{\left( B_{m,0}(0.25) P_{0,j} + B_{m,1}(0.25) P_{1,j} + B_{m,2}(0.25) P_{2,j} \right)}_{\text{new control points, for } j = 0, 1, 2 \text{ (i.e., } Q_0, Q_1, Q_2)}$$



# PROPERTIES OF BEZIER SURFACES

## Properties of Bézier surface (on rectangular grid)

- Uniparametric curves  
(i.e., fixed  $u$  or fixed  $v$ )

$$S(u, v) = \sum_{i=0}^m \sum_{j=0}^n B_{m,i}(u) B_{n,j}(v) P_{i,j}$$

Consider the curve obtained for a fix parameter, e.g.,  $u = u_0$


Then we have:

$$\begin{aligned} S(u_0, v) &= \sum_{i=0}^m \sum_{j=0}^n B_{m,i}(u_0) B_{n,j}(v) P_{i,j} \\ &= \sum_{j=0}^n B_{n,j}(v) \underbrace{\left( \sum_{i=0}^m B_{m,i}(u_0) P_{i,j} \right)}_{\text{new control point } Q_j} \end{aligned}$$

Example on the right, for  $u = 0.25$ ,  $n = m = 2$

$$S(0.25, v) = \sum_{j=0}^2 B_{n,j}(v) \left( \sum_{i=0}^2 B_{m,i}(u_0) P_{i,j} \right)$$

$$S(0.25, v) = \sum_{j=0}^2 B_{n,j}(v) \underbrace{\left( B_{m,0}(0.25) P_{0,j} + B_{m,1}(0.25) P_{1,j} + B_{m,2}(0.25) P_{2,j} \right)}_{\text{new control points, for } j = 0, 1, 2 \text{ (i.e., } Q_0, Q_1, Q_2)}$$



# PROPERTIES OF BEZIER SURFACES

## Properties of Bézier surface (on rectangular grid)


- Uniparametric curves  
(i.e., fixed  $u$  or fixed  $v$ )

$$S(u, v) = \sum_{i=0}^m \sum_{j=0}^n B_{m,i}(u) B_{n,j}(v) P_{i,j}$$

All uniparametric curves are Bézier curves

- Affine invariance

$$\sum_{i=0}^m \sum_{j=0}^n B_{m,i}(u) B_{n,j}(v) = 1 \quad \text{for all } (u, v)$$



# PROPERTIES OF BEZIER SURFACES

## Properties of Bézier surface (on rectangular grid)

- Uniparametric curves  
(i.e., fixed  $u$  or fixed  $v$ )


$$S(u, v) = \sum_{i=0}^m \sum_{j=0}^n B_{m,i}(u) B_{n,j}(v) P_{i,j}$$

All uniparametric curves are Bézier curves

- Affine invariance

$$\sum_{i=0}^m \sum_{j=0}^n B_{m,i}(u) B_{n,j}(v) = 1 \quad \text{for all } (u, v)$$

- Convex hull property



# PROPERTIES OF BEZIER SURFACES

## Properties of Bézier surface (on rectangular grid)

- Uniparametric curves  
(i.e., fixed  $u$  or fixed  $v$ )

$$S(u, v) = \sum_{i=0}^m \sum_{j=0}^n B_{m,i}(u) B_{n,j}(v) P_{i,j}$$


All uniparametric curves are Bézier curves

- Affine invariance

$$\sum_{i=0}^m \sum_{j=0}^n B_{m,i}(u) B_{n,j}(v) = 1 \quad \text{for all } (u, v)$$

- Convex hull property

- No variation diminishing property



# CONNECTING BEZIER SURFACES

Smooth connection of rectangular Bézier patches

# CONNECTING BEZIER SURFACES


## Smooth connection of rectangular Bézier patches

- Same idea as for curves

# CONNECTING BEZIER SURFACES

## Smooth connection of rectangular Bézier patches

- Same idea as for curves






# CONNECTING BEZIER SURFACES

## Smooth connection of rectangular Bézier patches


- Same idea as for curves



# CONNECTING BEZIER SURFACES

## Smooth connection of rectangular Bézier patches


- Same idea as for curves



# CONNECTING BEZIER SURFACES

## Smooth connection of rectangular Bézier patches

- Same idea as for curves




- Continuity ( $C^0$ -cont)

$$P_{i,n} = Q_{i,0} \quad \forall i = 0, \dots, m$$

# CONNECTING BEZIER SURFACES

## Smooth connection of rectangular Bézier patches

- Same idea as for curves




- Continuity ( $C^0$ -cont)
- Smoothness ( $C^1$ -cont)

$$P_{i,n} = Q_{i,0} \quad \forall i = 0, \dots, m$$

# CONNECTING BEZIER SURFACES

## Smooth connection of rectangular Bézier patches

- Same idea as for curves



- Continuity ( $C^0$ -cont)
- Smoothness ( $C^1$ -cont)

$$P_{i,n} = Q_{i,0} \quad \forall i = 0, \dots, m$$

$$\left. \frac{\partial P(u, v)}{\partial u} \right|_{u=1} = \left. \frac{\partial Q(u, v)}{\partial u} \right|_{u=0}$$

# CONNECTING BEZIER SURFACES

Smoothness condition ( $C^1$ -continuity)

$$\left. \frac{\partial P(u,v)}{\partial u} \right|_{u=1} = \left. \frac{\partial Q(u,v)}{\partial u} \right|_{u=0}$$

# CONNECTING BEZIER SURFACES

## Smoothness condition ( $C^1$ -continuity)

$$\left. \frac{\partial P(u,v)}{\partial u} \right|_{u=1} = \left. \frac{\partial Q(u,v)}{\partial u} \right|_{u=0}$$

$$P(u,v) = \sum_{i=0}^m \sum_{j=0}^n B_{m,i}(u) B_{n,j}(v) P_{i,j} = \sum_{i=0}^m \sum_{j=0}^n \binom{m}{i} u^i (1-u)^{m-i} \binom{n}{j} v^j (1-v)^{n-j} P_{i,j}$$

# CONNECTING BEZIER SURFACES

## Smoothness condition ( $C^1$ -continuity)

$$\left. \frac{\partial P(u,v)}{\partial u} \right|_{u=1} = \left. \frac{\partial Q(u,v)}{\partial u} \right|_{u=0}$$

$$P(u,v) = \sum_{i=0}^m \sum_{j=0}^n B_{m,i}(u) B_{n,j}(v) P_{i,j} = \sum_{i=0}^m \sum_{j=0}^n \binom{m}{i} u^i (1-u)^{m-i} \binom{n}{j} v^j (1-v)^{n-j} P_{i,j}$$

$$\left. \frac{\partial P(u,v)}{\partial u} \right|_{u=1} = \sum_{i=0}^m \binom{m}{i} n v^i (1-v)^{m-i} P_{i,n} - \sum_{i=0}^m \binom{m}{i} n v^i (1-v)^{m-i} P_{i,n-1}$$



# CONNECTING BEZIER SURFACES

## Smoothness condition ( $C^1$ -continuity)

$$\left. \frac{\partial P(u,v)}{\partial u} \right|_{u=1} = \left. \frac{\partial Q(u,v)}{\partial u} \right|_{u=0}$$

$$P(u,v) = \sum_{i=0}^m \sum_{j=0}^n B_{m,i}(u) B_{n,j}(v) P_{i,j} = \sum_{i=0}^m \sum_{j=0}^n \binom{m}{i} u^i (1-u)^{m-i} \binom{n}{j} v^j (1-v)^{n-j} P_{i,j}$$

$$\begin{aligned} \left. \frac{\partial P(u,v)}{\partial u} \right|_{u=1} &= \sum_{i=0}^m \binom{m}{i} n v^i (1-v)^{m-i} P_{i,n} - \sum_{i=0}^m \binom{m}{i} n v^i (1-v)^{m-i} P_{i,n-1} \\ &= n \sum_{i=0}^m \binom{m}{i} v^i (1-v)^{m-i} (P_{i,n} - P_{i,n-1}) \end{aligned}$$

# CONNECTING BEZIER SURFACES

## Smoothness condition ( $C^1$ -continuity)

$$\left. \frac{\partial P(u,v)}{\partial u} \right|_{u=1} = \left. \frac{\partial Q(u,v)}{\partial u} \right|_{u=0}$$

$$P(u,v) = \sum_{i=0}^m \sum_{j=0}^n B_{m,i}(u) B_{n,j}(v) P_{i,j} = \sum_{i=0}^m \sum_{j=0}^n \binom{m}{i} u^i (1-u)^{m-i} \binom{n}{j} v^j (1-v)^{n-j} P_{i,j}$$

$$\begin{aligned} \left. \frac{\partial P(u,v)}{\partial u} \right|_{u=1} &= \sum_{i=0}^m \binom{m}{i} n v^i (1-v)^{m-i} P_{i,n} - \sum_{i=0}^m \binom{m}{i} n v^i (1-v)^{m-i} P_{i,n-1} \\ &= n \sum_{i=0}^m \binom{m}{i} v^i (1-v)^{m-i} (P_{i,n} - P_{i,n-1}) \end{aligned}$$

Analogously,

$$\left. \frac{\partial Q(u,v)}{\partial u} \right|_{u=0} = k \sum_{i=0}^m \binom{m}{i} v^i (1-v)^{m-i} (Q_{i,1} - Q_{i,0})$$

# CONNECTING BEZIER SURFACES

## Smoothness condition ( $C^1$ -continuity)

$$\left. \frac{\partial P(u,v)}{\partial u} \right|_{u=1} = \left. \frac{\partial Q(u,v)}{\partial u} \right|_{u=0}$$

$$P(u,v) = \sum_{i=0}^m \sum_{j=0}^n B_{m,i}(u) B_{n,j}(v) P_{i,j} = \sum_{i=0}^m \sum_{j=0}^n \binom{m}{i} u^i (1-u)^{m-i} \binom{n}{j} v^j (1-v)^{n-j} P_{i,j}$$

$$\begin{aligned} \left. \frac{\partial P(u,v)}{\partial u} \right|_{u=1} &= \sum_{i=0}^m \binom{m}{i} n v^i (1-v)^{m-i} P_{i,n} - \sum_{i=0}^m \binom{m}{i} n v^i (1-v)^{m-i} P_{i,n-1} \\ &= n \sum_{i=0}^m \binom{m}{i} v^i (1-v)^{m-i} (P_{i,n} - P_{i,n-1}) \end{aligned}$$

Analogously,

$$\left. \frac{\partial Q(u,v)}{\partial u} \right|_{u=0} = k \sum_{i=0}^m \binom{m}{i} v^i (1-v)^{m-i} (Q_{i,1} - Q_{i,0})$$

Therefore, the condition for  $C^1$ -continuity is:  $n(P_{i,n} - P_{i,n-1}) = k(Q_{i,1} - Q_{i,0}) \quad \forall i = 0, \dots, m$

# CONNECTING BEZIER SURFACES

## Smoothness condition ( $C^1$ -continuity)

$$\left. \frac{\partial P(u,v)}{\partial u} \right|_{u=1} = \left. \frac{\partial Q(u,v)}{\partial u} \right|_{u=0}$$

$$P(u,v) = \sum_{i=0}^m \sum_{j=0}^n B_{m,i}(u) B_{n,j}(v) P_{i,j} = \sum_{i=0}^m \sum_{j=0}^n \binom{m}{i} u^i (1-u)^{m-i} \binom{n}{j} v^j (1-v)^{n-j} P_{i,j}$$

$$\begin{aligned} \left. \frac{\partial P(u,v)}{\partial u} \right|_{u=1} &= \sum_{i=0}^m \binom{m}{i} n v^i (1-v)^{m-i} P_{i,n} - \sum_{i=0}^m \binom{m}{i} n v^i (1-v)^{m-i} P_{i,n-1} \\ &= n \sum_{i=0}^m \binom{m}{i} v^i (1-v)^{m-i} (P_{i,n} - P_{i,n-1}) \end{aligned}$$

Analogously,

$$\left. \frac{\partial Q(u,v)}{\partial u} \right|_{u=0} = k \sum_{i=0}^m \binom{m}{i} v^i (1-v)^{m-i} (Q_{i,1} - Q_{i,0})$$


Therefore, the condition for  $C^1$ -continuity is:  $n(P_{i,n} - P_{i,n-1}) = k(Q_{i,1} - Q_{i,0}) \quad \forall i = 0, \dots, m$

If we just want  $G^1$ -cont, it is enough with  $P_{i,n} - P_{i,n-1} = \alpha(Q_{i,1} - Q_{i,0})$ , for some  $\alpha \neq 0 \in \mathbb{R}$

# DE CASTELJAU'S ALGORITHM

## Applying De Casteljau to each dimension



- For surfaces: apply it in two phases (along  $u$ , and along  $v$ )



# DE CASTELJAU'S ALGORITHM

## Applying De Casteljau to each dimension



- For surfaces: apply it in two phases (along  $u$ , and along  $v$ )



# DE CASTELJAU'S ALGORITHM

## Applying De Casteljau to each dimension



- For surfaces: apply it in two phases (along  $u$ , and along  $v$ )



# DE CASTELJAU'S ALGORITHM

## Applying De Casteljau to each dimension

- For surfaces: apply it in two phases (along  $u$ , and along  $v$ )








# DE CASTELJAU'S ALGORITHM

## Applying De Casteljau to each dimension




- For surfaces: apply it in two phases (along  $u$ , and along  $v$ )



# DE CASTELJAU'S ALGORITHM

## Applying De Casteljau to each dimension




- For surfaces: apply it in two phases (along  $u$ , and along  $v$ )



# DE CASTELJAU'S ALGORITHM

## Applying De Casteljau to each dimension




- For surfaces: apply it in two phases (along  $u$ , and along  $v$ )



# DE CASTELJAU'S ALGORITHM

## Applying De Casteljau to each dimension




- For surfaces: apply it in two phases (along  $u$ , and along  $v$ )



# DE CASTELJAU'S ALGORITHM

## Applying De Casteljau to each dimension




- For surfaces: apply it in two phases (along  $u$ , and along  $v$ )



# DE CASTELJAU'S ALGORITHM

## Applying De Casteljau to each dimension




- For surfaces: apply it in two phases (along  $u$ , and along  $v$ )



# DE CASTELJAU'S ALGORITHM

## Applying De Casteljau to each dimension




- For surfaces: apply it in two phases (along  $u$ , and along  $v$ )



# DE CASTELJAU'S ALGORITHM

## Applying De Casteljau to each dimension

- For surfaces: apply it in two phases (along  $u$ , and along  $v$ )



- Applications of De Casteljau, e.g., to curve subdivision, also extend to surfaces



# INTERPOLATING SURFACE

## Interpolating Bézier surface patch

Problem: given  $(m + 1) \times (n + 1)$  data points  $Q_{k,l}$ , compute a set of  $(m + 1) \times (n + 1)$  control points  $P_{i,j}$  such that the Bézier surface  $S$  defined by the points  $P_{i,j}$  goes through the points  $Q_{k,l}$

# INTERPOLATING SURFACE

## Interpolating Bézier surface patch

Problem: given  $(m + 1) \times (n + 1)$  data points  $Q_{k,l}$ , compute a set of  $(m + 1) \times (n + 1)$  control points  $P_{i,j}$  such that the Bézier surface  $S$  defined by the points  $P_{i,j}$  goes through the points  $Q_{k,l}$

Same approach as for curves:

# INTERPOLATING SURFACE

## Interpolating Bézier surface patch

Problem: given  $(m + 1) \times (n + 1)$  data points  $Q_{k,l}$ , compute a set of  $(m + 1) \times (n + 1)$  control points  $P_{i,j}$  such that the Bézier surface  $S$  defined by the points  $P_{i,j}$  goes through the points  $Q_{k,l}$

Same approach as for curves:

- 1) Select  $m + 1$  values  $u_k$  (for  $k = 0, \dots, m$ )

# INTERPOLATING SURFACE

## Interpolating Bézier surface patch

Problem: given  $(m + 1) \times (n + 1)$  data points  $Q_{k,l}$ , compute a set of  $(m + 1) \times (n + 1)$  control points  $P_{i,j}$  such that the Bézier surface  $S$  defined by the points  $P_{i,j}$  goes through the points  $Q_{k,l}$

Same approach as for curves:

- 1) Select  $m + 1$  values  $u_k$  (for  $k = 0, \dots, m$ )
- 2) Select  $n + 1$  values  $v_l$  (for  $l = 0, \dots, n$ )

# INTERPOLATING SURFACE

## Interpolating Bézier surface patch

Problem: given  $(m + 1) \times (n + 1)$  data points  $Q_{k,l}$ , compute a set of  $(m + 1) \times (n + 1)$  control points  $P_{i,j}$  such that the Bézier surface  $S$  defined by the points  $P_{i,j}$  goes through the points  $Q_{k,l}$

Same approach as for curves:

- 1) Select  $m + 1$  values  $u_k$  (for  $k = 0, \dots, m$ )
- 2) Select  $n + 1$  values  $v_l$  (for  $l = 0, \dots, n$ )
- 3) Require  $S(u_k, v_l) = Q_{k,l}$  for all  $k, l$

# INTERPOLATING SURFACE

## Interpolating Bézier surface patch

Problem: given  $(m + 1) \times (n + 1)$  data points  $Q_{k,l}$ , compute a set of  $(m + 1) \times (n + 1)$  control points  $P_{i,j}$  such that the Bézier surface  $S$  defined by the points  $P_{i,j}$  goes through the points  $Q_{k,l}$

Same approach as for curves:

- 1) Select  $m + 1$  values  $u_k$  (for  $k = 0, \dots, m$ )
- 2) Select  $n + 1$  values  $v_l$  (for  $l = 0, \dots, n$ )
- 3) Require  $S(u_k, v_l) = Q_{k,l}$  for all  $k, l$

Defines system of  $(m + 1) \times (n + 1)$  equations that always has a solution

# INTERPOLATING SURFACE

## Interpolating Bézier surface patch

Problem: given  $(m + 1) \times (n + 1)$  data points  $Q_{k,l}$ , compute a set of  $(m + 1) \times (n + 1)$  control points  $P_{i,j}$  such that the Bézier surface  $S$  defined by the points  $P_{i,j}$  goes through the points  $Q_{k,l}$

Same approach as for curves:

- 1) Select  $m + 1$  values  $u_k$  (for  $k = 0, \dots, m$ )
- 2) Select  $n + 1$  values  $v_l$  (for  $l = 0, \dots, n$ )
- 3) Require  $S(u_k, v_l) = Q_{k,l}$  for all  $k, l$

Defines system of  $(m + 1) \times (n + 1)$  equations that always has a solution

$$S(u_k, v_l) = B_m^t(u_k) P B_n(v_l) = Q_{k,l}, \text{ for } k = 0, \dots, m \text{ and } l = 0, \dots, n$$

# INTERPOLATING SURFACE

## Interpolating Bézier surface patch

Problem: given  $(m + 1) \times (n + 1)$  data points  $Q_{k,l}$ , compute a set of  $(m + 1) \times (n + 1)$  control points  $P_{i,j}$  such that the Bézier surface  $S$  defined by the points  $P_{i,j}$  goes through the points  $Q_{k,l}$

Same approach as for curves:

- 1) Select  $m + 1$  values  $u_k$  (for  $k = 0, \dots, m$ )
- 2) Select  $n + 1$  values  $v_l$  (for  $l = 0, \dots, n$ )
- 3) Require  $S(u_k, v_l) = Q_{k,l}$  for all  $k, l$

Defines system of  $(m + 1) \times (n + 1)$  equations that always has a solution

$$S(u_k, v_l) = \underbrace{B_m^t(u_k) P B_n(v_l)}_{\text{matricial form of Bézier surface formula}} = Q_{k,l}, \text{ for } k = 0, \dots, m \text{ and } l = 0, \dots, n$$

matricial form of Bézier surface formula



# INTERPOLATING SURFACE

## Interpolating Bézier surface patch

Example from [Salomon, page 232]:

**Example:** We choose  $m = 3$  and  $n = 2$ . The system of equations becomes

$$\left[ (1 - u_k)^3, 3u_k(1 - u_k)^2, 3u_k^2(1 - u_k), u_k^3 \right] \begin{bmatrix} \mathbf{P}_{00} & \mathbf{P}_{01} & \mathbf{P}_{02} \\ \mathbf{P}_{10} & \mathbf{P}_{11} & \mathbf{P}_{12} \\ \mathbf{P}_{20} & \mathbf{P}_{21} & \mathbf{P}_{22} \\ \mathbf{P}_{30} & \mathbf{P}_{31} & \mathbf{P}_{32} \end{bmatrix} \begin{bmatrix} (1 - w_l)^2 \\ 2w_l(1 - w_l) \\ w_l^2 \end{bmatrix} = \mathbf{Q}_{kl}$$

# INTERPOLATING SURFACE

## Interpolating Bézier surface patch

Example from [Salomon, page 232]:

**Example:** We choose  $m = 3$  and  $n = 2$ . The system of equations becomes

$$\left[ (1 - u_k)^3, 3u_k(1 - u_k)^2, 3u_k^2(1 - u_k), u_k^3 \right] \begin{bmatrix} \mathbf{P}_{00} & \mathbf{P}_{01} & \mathbf{P}_{02} \\ \mathbf{P}_{10} & \mathbf{P}_{11} & \mathbf{P}_{12} \\ \mathbf{P}_{20} & \mathbf{P}_{21} & \mathbf{P}_{22} \\ \mathbf{P}_{30} & \mathbf{P}_{31} & \mathbf{P}_{32} \end{bmatrix} \begin{bmatrix} (1 - w_l)^2 \\ 2w_l(1 - w_l) \\ w_l^2 \end{bmatrix} = \mathbf{Q}_{kl}$$

12 equations, 12 unknowns

# RATIONAL BÉZIER SURFACES

Rational rectangular Bézier surface patch

# RATIONAL BÉZIER SURFACES

## Rational rectangular Bézier surface patch

Definition analogous to the one for curves

# RATIONAL BÉZIER SURFACES

## Rational rectangular Bézier surface patch

Definition analogous to the one for curves

$$P(u, w) = \frac{\sum_{i=0}^m \sum_{j=0}^n w_{i,j} B_{m,i}(u) B_{n,j}(w) P_{i,j}}{\sum_{i=0}^m \sum_{j=0}^n w_{i,j} B_{m,i}(u) B_{n,j}(w)} \quad \begin{array}{l} 0 \leq u, w \leq 1 \\ w_{i,j} \in \mathbb{R}_{>0} \text{ for all } i, j \end{array}$$

If all weights are  $w_{i,j} = 1$ , it reduces to the ordinary Bézier surface

# TRIANGULAR PATCHES


Surface patches don't need to be rectangular

$$\begin{array}{ccccccc} & & & & & & \mathbf{P}_{040} \\ & & & & & \mathbf{P}_{031} & \mathbf{P}_{130} \\ & & & & \mathbf{P}_{020} & \mathbf{P}_{121} & \mathbf{P}_{220} \\ & & \mathbf{P}_{010} & \mathbf{P}_{011} & \mathbf{P}_{110} & \mathbf{P}_{022} & \mathbf{P}_{121} & \mathbf{P}_{220} \\ & \mathbf{P}_{001} & \mathbf{P}_{100} & \mathbf{P}_{002} & \mathbf{P}_{101} & \mathbf{P}_{200} & \mathbf{P}_{013} & \mathbf{P}_{112} & \mathbf{P}_{211} & \mathbf{P}_{310} \\ & & & & & & \mathbf{P}_{004} & \mathbf{P}_{103} & \mathbf{P}_{202} & \mathbf{P}_{301} & \mathbf{P}_{400} \end{array}$$

# TRIANGULAR PATCHES

Surface patches don't need to be rectangular


Control points arranged as triangular array



# TRIANGULAR PATCHES

Surface patches don't need to be rectangular

Control points arranged as triangular array





Bézier formula needs version based on three variables

$$\mathbf{P}(u, v, w) = \sum_{i+j+k=n} \mathbf{P}_{ijk} \frac{n!}{i!j!k!} u^i v^j w^k = \sum_{i+j+k=n} \mathbf{P}_{ijk} B_{ijk}^n(u, v, w)$$



# TRIANGULAR PATCHES

## Example



$$n = 3$$