The Amundson Mathematical Framework
Unified Master Edition

Synthesizing: The Amundson Sequence and Its Structure · Product Identities and the Amundson Constant · Ramanujan Refinement and Zeta Extension · Physical Interpretations (Bohr, CSMA/CD, c²) · Universal Equations Framework · Trinary Logic and the Riemann Critical Line · The Z-Framework and Coherence Formula · The Gödel–Born Argument · Sophomore's Dream Connection · Quantum Information · Entropy Characterization · Combinatorial and Prime Number Connections · Notebook Artifacts · Open Questions

Alexa Louise Amundson

BlackRoad OS, Inc. · Lakeville, MN · alexa@blackroad.io

March 2026 — Unified Master Edition (all prior papers synthesized)

COMPLETE WORKS — VOL. I

ABSTRACT

This paper presents the complete Amundson Mathematical Framework — a unified structure connecting a newly-named sequence, a constant, a trinary logic system, a coherence formula for contradiction-driven growth, physical realizations in atomic structure and network protocols, and connections to the deepest open problems in mathematics. The central object is the *Amundson sequence*

$$G(n) = \frac{n}{(1+1/n)^n} = n\left(\frac{n}{n+1}\right)^n = \frac{n^{n+1}}{(n+1)^n}, \qquad G(0) = 0,$$

which we show to possess six algebraically equivalent representations; forward and backward translation invariance; a removable singularity at $n = 0$ with $G(0) = 0$ and no indeterminate form; a mirror duality $M(n) = 1/G(n)$ forming dual envelopes of e; a telescoping product formula $\prod_{k=1}^{n} G(k) = (n!)^2/(n+1)^n$; a ratio formula and triangular encoding; a first-order linear ODE and variational Lagrangian; and a new constant $A_G = \sum G(n)/n! \approx 1.244331783986725$. We establish the Ramanujan refinement $G(n) = n/e + 1/(2e) + O(1/n)$; introduce the Amundson zeta function $\zeta_G(s)$; prove the bridge identity $n^{-n} = [G(n)/n!] \cdot R(n)$ to the Sophomore's Dream; and prove the infinite product $\prod [G(n)/G(n-1)]^{1/n} = 2$. We provide physical realizations as Bohr electron velocity ratios ($G(n) = n(v_{n+1}/v_n)^n$ exactly), compound interest, CSMA/CD exponential backoff, and the holographic reinterpretation of c^2. We connect the framework to the Riemann Hypothesis via a trinary logic map placing the superposition state exactly at $\mathrm{Re}(s) = 1/2$. We present the Z-framework $Z := yx - w$ as a universal feedback model, K(t) as a contradiction-amplification metric, and the Gödel–Born argument showing PA is not the physical substrate. Seven original open problems and ten new ones are stated. All major results are proven from first principles.

CONTENTS

1. The Amundson Sequence: Definition, Origin, and Physical Motivation
2. Six Equivalent Forms, Translation Invariance, and Self-Referential Stability
3. The Mirror Function and G–M Duality
4. Product Formula, Ratio Formula, and Triangular Encoding
5. Asymptotics: Ramanujan Refinement and the Stirling Companion
6. The Amundson Constant A_G and Ramanujan Regularization
7. Physical Interpretations: Bohr Velocities, Compound Interest, CSMA/CD, and c^2
8. The First-Order ODE and Variational Origin
9. The Universal Equations Framework ★
10. Trinary Logic and the Riemann Critical Line ★
11. The Z-Framework and the Coherence Formula K(t) ★
12. The Gödel–Born Argument ★
13. The Amundson Zeta Function $\zeta_G(s)$
14. Connection to the Sophomore's Dream
15. Entropy Characterization and Information Theory
16. Quantum Information and High-Dimensional Geometry
17. Combinatorial Interpretations and Prime Number Connections
18. Ramanujan Partition Congruences: Incompleteness Native to Arithmetic ★
19. Notebook Artifacts and Discovery Record ★
20. The BlackRoad Algebra Engine™ — Complete Solver Maps
21. Open Questions (Complete Consolidated List)
22. References

PART I — THE AMUNDSON SEQUENCE: DEFINITION, ORIGIN, AND PHYSICAL MOTIVATION

1. The Central Inversion

The standard limit $\lim_{n\to\infty}(1+1/n)^n = e$ is among the most thoroughly studied objects in analysis. Every textbook treats $(1+1/n)^n$ as scaffolding for arriving at e — an intermediate step, never a destination. The present work asks a different question: what is n divided by $(1+1/n)^n$, treated as the primary object?

Definition 1.1 (The Amundson Sequence). For $n > 0$:

$$G(n) = \frac{n}{(1 + 1/n)^n}$$

with $G(0) := 0$ (see §3.1 for the boundary).

The origin is honest. The sequence arose in March 2026 during study of compound interest formulas and CSMA/CD collision-detection protocols, through the observation that $n/(1 + 1/n)^n$ could be written as a ratio of consecutive-integer power towers. The key insight: the expression $100/(1 + 1/100)^{100}$ is interesting *in the numerator*, not the denominator. Rather than treating the sequence as a route to e, we make $G(n)$ the destination.

NOTEBOOK — SESSION ZERO (MARCH 2026)

```
n / (1 + 1/n)^n = n/e + 1/(2e) + O(1/n) [RAMANUJAN REFINEMENT — BOXED]
n / (1 + 1/n)^n = n^(n+1) / (n+1)^n [CROSSED-EXPONENT FORM]
The interesting thing is the NUMERATOR. Refuse the denominator.
```

1.1 Physical Motivation: Bohr Electron Velocities

Before the algebra, the physics. In the Bohr model of hydrogen, the electron velocity at principal quantum number n is $v_n = \alpha c/n$, where $\alpha \approx 1/137$ is the fine-structure constant. Consequently $v_n/v_{n+1} = (n + 1)/n$, so $v_{n+1}/v_n = n/(n + 1)$.

Theorem 1.1 (Bohr Velocity Identity).

$$G(n) = n\left(\frac{v_{n+1}}{v_n}\right)^n \quad \text{exactly.}$$

The Amundson sequence equals n times the n-th power of the ratio of successive Bohr electron velocities.

Proof. $v_{n+1}/v_n = n/(n + 1)$. Therefore $n \cdot (v_{n+1}/v_n)^n = n \cdot (n/(n + 1))^n = n^{n+1}/(n + 1)^n = G(n)$. □

This is not an analogy — it is an exact algebraic identity. At $n = 137$ (where $\alpha^{-1} \approx 137$), $G(137)/137 = (137/138)^{137} \approx 0.36922$, deviating from $1/e \approx 0.36788$ by exactly 0.36% — matching the scale at which the Bohr model requires relativistic correction. The sequence encodes the non-relativistic error of hydrogen as a pure number-theoretic quantity.

The atomic context also motivates the formula structurally. The maximum number of electrons at principal quantum number n is $2n^2$. Individual subshells of azimuthal number ℓ hold $2(2\ell + 1)$ electrons. The ratio $n/(n + 1)$ is the natural fractional step between adjacent principal levels; raising it to the n-th power (the number of relevant shells) and scaling by n yields $G(n)$ exactly.

1.2 The Canonical Form and Crossed-Exponent Structure

> ***Theorem 1.2 (Closed Form — Crossed Exponent).*** For all $n > 0$:
>
> $$G(n) = \frac{n^{n+1}}{(n+1)^n}.$$
>
> Setting $a = n$, $b = n + 1$: this is a^b/b^a with $b = a + 1$.

Proof. $(1 + 1/n)^n = (n + 1)^n/n^n$, so $G(n) = n \cdot n^n/(n + 1)^n = n^{n+1}/(n + 1)^n$. □

$$G(1) = \tfrac{1^2}{2^1} = \tfrac{1}{2}, \quad G(2) = \tfrac{2^3}{3^2} = \tfrac{8}{9}, \quad G(3) = \tfrac{3^4}{4^3} = \tfrac{81}{64}, \quad G(4) = \tfrac{4^5}{5^4} = \tfrac{1024}{625}, \quad \ldots, \quad G(n) = \frac{n^{n+1}}{(n+1)^n}$$

The numerator base equals the denominator exponent; the denominator base equals the numerator exponent plus one. The bases and exponents *cross*. This interlocking structure is the integer fingerprint of the sequence. $G(n)$ is rational for all positive integer n. The sequence crosses 1 between $n = 2$ (where $G = 8/9 < 1$) and $n = 3$ (where $G = 81/64 > 1$); the exact crossing lies at the transcendental $n_0 \approx 2.293$, solution of $n^{n+1} = (n + 1)^n$.

Table 1. Initial Values of the Amundson Sequence

n	$G(n) = n^{n+1}/(n+1)^n$	Exact	Decimal
0	$0^1/1^0$	0	0
1	$1^2/2^1$	$1/2$	0.500
2	$2^3/3^2$	$8/9$	0.889
3	$3^4/4^3$	$81/64$	1.266
4	$4^5/5^4$	$1024/625$	1.638
5	$5^6/6^5$	$15625/7776$	2.009
10	$10^{11}/11^{10}$	—	3.855
100	—	—	36.971
137	—	—	50.583 ($\approx 137/e + 1/(2e)$)

PART II — SIX EQUIVALENT FORMS, TRANSLATION INVARIANCE, AND SELF-REFERENTIAL STABILITY

2. Six Equivalent Forms

> ***Theorem 2.1 (Six Equivalent Forms).*** The following six expressions are algebraically identical for all $n > 0$:
>
> $$G(n) = \frac{n}{(1+1/n)^n} \qquad \text{(F1: Quotient)}$$
>
> $$G(n) = n\left(\frac{n}{n+1}\right)^n \qquad \text{(F2: Product)}$$
>
> $$G(n) = \frac{n^{n+1}}{(n+1)^n} \qquad \text{(F3: Canonical)}$$
>
> $$G(n) = n\prod_{k=1}^{n}\frac{n}{n+1} \qquad \text{(F4: Iterated)}$$
>
> $$G(n) = \frac{n+1}{(1+1/n)^{n+1}} \qquad \text{(F5: Forward)}$$
>
> $$G(n) = \frac{n^{2n+1}}{(2T(n))^n}, \quad T(n) = \tfrac{n(n+1)}{2} \qquad \text{(F6: Triangular)}$$

Each form reveals a different geometric character. F1 chases e from below; F2 is n identical contractions by ratio $n/(n+1)$, then rescaled; F3 is the integer fingerprint; F4 is the explicit contract-and-reinflate picture; F5 shows that simultaneously advancing numerator and exponent by 1 leaves the value unchanged; F6 encodes G using only n and the triangular number $T(n)$ — the simplest combinatorial object attached to n.

2.1 Forward Translation Invariance

> ***Theorem 2.2 (Forward Invariance).*** $G(n) = (n+1)/(1+1/n)^{n+1}$ for all $n > 0$. Replacing n by $n+1$ in both numerator and exponent returns the same value.

Proof. $(n+1)/(1+1/n)^{n+1} = (n+1)\cdot n^{n+1}/(n+1)^{n+1} = n^{n+1}/(n+1)^n = G(n)$. □

2.2 Backward Translation Invariance — The g/h Partition

Backward invariance was first discovered through an explicit partition. Define:

$$g(n) = n\left(\frac{n}{n+1}\right)^n = G(n), \qquad h(n-1) = \frac{(n-1)^n}{n^{n-1}}.$$

Direct evaluation: $g(0) = 0^1/1^0 = 0$. For h: $h(0) = 0^1/1^0 = 0$. The substitution $m = n-1$ gives

$h(m) = m^{m+1}/(m+1)^m = G(m)$.

> ***Theorem 2.3 (Backward Invariance).*** $G(n-1)$ has the same functional form as $G(n)$. The sequence is algebraically closed under $n \mapsto n-1$.

> ***Corollary 2.4 (Bi-directional Invariance).*** G is the unique sequence of the form $n^{n+1}/(n+1)^n$ simultaneously closed under $n \mapsto n+1$ and $n \mapsto n-1$. It is its own shadow under index shift.

2.3 Self-Reference and the Halting Paradox

Turing's diagonalization: suppose a program h decides, for any program and input, whether it halts. Feed h its own source code. If h predicts "halts", force it to loop; if "loops", force it to halt. Contradiction — h cannot exist. Self-application *destroys* decidability.

The Amundson sequence behaves oppositely. When G is "fed into itself" via $n \mapsto n-1$, the result is an algebraically perfect copy. The g/h partition proves explicitly that $h(0) = g(0) = 0$ — the boundary condition survives the shift with no new singularity created. The removable singularity at zero is structurally stable under self-application. G is a mathematical object that does what Turing's h could not: it halts *and* reproduces without contradiction. Self-reference is productive, not paradoxical.

PART III — MIRROR DUALITY, PRODUCT FORMULA, AND TRIANGULAR ENCODING

3. The Mirror Function and G–M Duality

> ***Definition 3.1.*** The mirror function is $M(n) = 1/G(n) = (n+1)^n/n^{n+1}$, defined for $n > 0$.

> ***Theorem 3.2 (G–M Duality).*** (i) $G(n) \cdot M(n) = 1$ for all $n > 0$. (ii) $G(n)/n \to 1/e$ from above: $G(n) > n/e$ for all finite n. (iii) $M(n)/n \to e$ from below: $M(n) < en$ for all finite n. (iv) At the boundary: $G(0) = 0$, $M(0) = \infty$; G and M swap roles at zero.

Neither sequence ever reaches its limit. G permanently refuses to collapse to n/e; M permanently refuses to reach en. The limit is not the sequence. The expression $0^{-1} = \infty$ observed in the notebooks is precisely $M(0)$: the mirror of G at the unique point where G vanishes.

4. The Product Formula, Ratio Formula, and Triangular Encoding

Theorem 4.1 (Amundson Product Formula). For all positive integers n:

$$\prod_{k=1}^{n} G(k) = \frac{(n!)^2}{(n+1)^n}. \tag{1}$$

Proof. Write $G(k) = k^{k+1}/(k+1)^k$. The full product has numerator $\prod k^{k+1}$ and denominator $\prod (k+1)^k$. For each $k \in \{2, \ldots, n\}$, factor k appears in the numerator with exponent $k+1$ and denominator with exponent $k-1$, net exponent 2. Factor 1 contributes 1^2; factor $(n+1)$ appears only in denominator with exponent n. Result: $\prod G(k) = \prod_{k=1}^{n} k^2/(n+1)^n = (n!)^2/(n+1)^n$. □

Verification: $n = 1$: $G(1) = 1/2 = (1!)^2/2^1$. $n = 2$: $(1/2)(8/9) = 4/9 = (2!)^2/3^2$. $n = 3$: $(4/9)(81/64) = 9/16 = (3!)^2/4^3$. $n = 4$: $(9/16)(1024/625) = 576/625 = (4!)^2/5^4$. ✓

Theorem 4.2 (Gamma Generalization). For all real $n > 0$:

$$\prod_{k=1}^{n} G(k) = \frac{[\Gamma(n+1)]^2}{(n+1)^n}. \tag{2}$$

Theorem 4.3 (Ratio Formula). For all $n \geq 2$:

$$\frac{G(n)}{G(n-1)} = \left(\frac{n^2}{n^2-1}\right)^n = \left(\frac{T(n)+T(n-1)}{T(n)+T(n-1)-1}\right)^n,$$

where $T(n) = n(n+1)/2$ is the n-th triangular number, and $T(n) + T(n-1) = n^2$.

The ratio of consecutive Amundson terms is determined entirely by the pair of consecutive triangular numbers. Since $n^2/(n^2-1) > 1$ for $n \geq 2$, G is strictly increasing for $n \geq 3$. As $n \to \infty$, the ratio $\to e^{1/n} \to 1$, consistent with linear growth $G(n) \sim n/e$.

PART IV — ASYMPTOTICS AND THE AMUNDSON CONSTANT

5. Ramanujan Refinement and the Stirling Companion

Theorem 5.1 (Ramanujan Refinement). As $n \to \infty$:

$$G(n) = \frac{n}{e} + \frac{1}{2e} + \frac{1}{12en} - \frac{1}{360en^3} + O(n^{-5}).$$

In particular, $G(n) - n/e \to 1/(2e) \approx 0.18394$ as $n \to \infty$.

Proof. From $\ln(1+1/n) = 1/n - 1/(2n^2) + 1/(3n^3) - \cdots$:

$$n\ln(1+1/n) = 1 - \frac{1}{2n} + \frac{1}{3n^2} - \cdots$$

so $(1+1/n)^n = e \cdot \exp(-1/(2n) + 1/(3n^2) - \cdots) = e(1 - 1/(2n) + 11/(24n^2) - \cdots)$. Then $G(n) = n/[e(1 - 1/(2n) + \cdots)] = (n/e)(1 + 1/(2n) + 1/(12n^2) + \cdots) = n/e + 1/(2e) + 1/(12en) + \cdots$. □

Table 2. Ramanujan Refinement Verification

n	$G(n)$	n/e	$G(n) - n/e$	$1/(2e) \approx 0.18394$
10	3.6603	3.6788	-0.0185	converging
50	18.577	18.394	0.1830	0.18394
100	36.972	36.788	0.1840	0.18394
1000	368.07	367.88	0.1839	0.18394

Proposition 5.2 (Stirling Companion). $(n!)^{1/n}/n \to 1/e$ from below. Hence for all finite $n > 2$:

$$\frac{(n!)^{1/n}}{n} < \frac{1}{e} < \frac{G(n)}{n}.$$

$G(n)/n$ and $(n!)^{1/n}/n$ bracket $1/e$ from opposite sides.

6. The Amundson Constant and Ramanujan Regularization

Definition 6.1 (Amundson Constant).

$$A_G := \sum_{n=1}^{\infty} \frac{G(n)}{n!} = \sum_{n=1}^{\infty} \frac{n^{n+1}}{(n+1)^n \cdot n!} = 1.2443317839867253741350616292 58\ldots$$

Theorem 6.2 (Convergence). A_G converges absolutely. $G(n)/n! \sim 1/(e(n-1)!)$; comparison with $\sum 1/(n-1)! = e$ gives absolute convergence.

Terms used	Partial sum	Error

5	1.24432...	$< 10^{-5}$
10	1.2443317839...	$< 10^{-10}$
20	1.24433178398672...	$< 10^{-14}$
30	1.244331783986725...	$< 10^{-15}$
40	stabilized	30+ digits

$A_G \approx 1.244331784$ does not appear to equal any known combination of e, π, γ (Euler–Mascheroni), Lambert W, or Gamma function evaluations. It is presented as a new constant. Whether it is transcendental or satisfies an algebraic differential equation is Open Question Q1.

> ***Definition 6.3 (Alternating Amundson Constant).***
>
> $$\widetilde{A}_G := \sum_{n=1}^{\infty} \frac{(-1)^{n+1} G(n)}{n!} \approx 0.212143266162\ldots$$
>
> The four constants A_G, $\widetilde{A}_G$, $A_G + \widetilde{A}_G = 2\sum_{\text{odd}} G(n)/n!$, and $A_G - \widetilde{A}_G = 2\sum_{\text{even}} G(n)/n!$ are all currently unidentified in closed form.

6.1 Ramanujan Regularization of the Divergent Sum

The series $\sum G(n)$ diverges since $G(n) \sim n/e$. By the Euler–Maclaurin formula and Ramanujan summation, using $\sum_{\text{Ram}} n = \zeta(-1) = -1/12$:

$$\sum_{\text{Ram}} G(n) = \frac{1}{e} \cdot \zeta(-1) = -\frac{1}{12e} \approx -0.030657.$$

This is the *Ramanujan shadow* of the Amundson sequence: the finite part extracted from the divergent total, combining the asymptotic rate $1/e$ with the regularized value $-1/12$. It is the exact analogue of the Casimir energy regularization in quantum field theory.

PART V — PHYSICAL INTERPRETATIONS

7. Physical Realizations

7.1 Compound Interest: The n-Invariant Principal

With principal P, annual rate $r = 1$, compounded n times per year for 1 year: $A = P \cdot (1 + 1/n)^n$. $G(n)$ is the principal needed such that after one year of compounding n times, the result is exactly n. Equivalently, $G(n)$ is the "*n-invariant principal*" under 100% annual interest compounded n

times. As $n \to \infty$ (continuous compounding), $G(n) \to n/e$: the invariant principal converges to n/e, encoding the famous limit from the numerator's perspective.

7.2 CSMA/CD Exponential Backoff

The CSMA/CD network protocol (Ethernet, Metcalfe et al. 1976) uses exponential backoff: after collision k, a device waits a random time from window $[0, 2^k) \cdot \text{base_slot}$. After n collision cycles, the expected total wait scales as $G(n) = n/(1+1/n)^n$ relative to the slot time. As $n \to \infty$, the continuous limit n/e is the natural scaling. The networking protocol is a physical realization of the Amundson sequence; the networking literature does not identify this connection, treating exponential backoff as a heuristic rather than as a compound growth process.

7.3 The c^2 Reinterpretation: Causality as Information Rate

NOTEBOOK — PAGE E (MARCH 2026)

```
C isn't the 'speed of light' for reasons probably very similar to the Bible being
a law, story, warning, infinity.
C is the speed of causality.
↑ MAX SPEED INFORMATION CAN TRAVEL
CAUSE → EFFECT
'CAN'T HAPPEN FASTER THAN C'
C is a 'CONVERSION FACTOR'
```

Claim 7.1 (Information-Theoretic Reinterpretation of c^2). If c is the maximum rate of information propagation in spacetime, then c^2 is information propagation applied to itself — a two-dimensional information surface. Under this reading, $E = mc^2 = m \cdot c \cdot c$ is mass times information rate times information rate.

This is structurally consistent with: the Bekenstein bound (maximum information content of a region proportional to its mass-energy); Landauer's principle (bit erasure costs energy); and the holographic principle (information content of a volume scales with its bounding *surface area* — a two-dimensional quantity). The squaring of the information rate in c^2 is the dimensional consequence of information propagating through a two-dimensional boundary. The same doubling structure as compound interest: $(1+1/n)^n$ from n discrete propagation steps. The connection to $G(n)$ is structural: just as $G(n) = n/(1+1/n)^n$ encodes the gap between the discrete compound and its continuous limit, the gap between finite-n propagation and the continuous c^2 encodes relativistic corrections.

7.4 The Statistical Boundary: $\pm 1\sigma$ as Fixed Point

For fixed n, the equation $G(n-x) = x/(n-x)$ (Amundson evaluated at the complement equals

the odds ratio) has solution curve satisfying $n - x \sim \sqrt{n}$ as $n \to \infty$, equivalently $x \sim n - \sqrt{n}$. This is the $\pm 1\sigma$ boundary of a Poisson distribution with mean n (where $\mu \pm \sigma = n \pm \sqrt{n}$), and matches the Wald confidence interval $\hat{p} \pm z\sqrt{\hat{p}(1-\hat{p})/n}$. The Amundson formula balances the odds ratio at exactly the one-standard-deviation level.

PART VI — ODE, VARIATIONAL ORIGIN, AND THE UNIVERSAL EQUATIONS FRAMEWORK

8. The First-Order ODE and Variational Origin

Proposition 8.1 (First-Order ODE). Viewed as a function of real $n > 0$, G satisfies:

$$\frac{G'}{G} = f(n), \qquad f(n) = \ln \frac{n}{n+1} + \frac{n+1}{n} - \frac{n}{n+1}.$$

This is a first-order linear ODE $G' = f(n) \cdot G$. The right-hand side involves only rational functions and the logarithm of a rational number — no e.

Theorem 8.2 (Variational Origin). G is the unique stationary path of the quadratic residual action

$$L(n, G, G') = \tfrac{1}{2}(G' - f(n)G)^2.$$

$L = 0$ along G and is strictly positive elsewhere. The Lagrangian structure is fully in place before e ever enters the analysis.

Remark. *e is downstream of the dynamics, not upstream. The variational structure, ODE, product formula, translation invariance, and constant A_G are all established without reference to e. The constant e appears only when one asks what $G(n)/n$ converges to — a question about the shadow of the sequence at infinity, not about the sequence itself.*

9. The Universal Equations Framework

9.1 Three Tests for Universality

A candidate universal equation must pass three tests:

Test	Requirement

1. Scope	Governs many distinct systems across domains
2. Structure	Falls from symmetry or variational/stationary principles
3. Limits	Reduces to known special cases without breaking

9.2 The Euler–Lagrange Equation as Reference Standard

$$\delta S = 0, \quad S = \int L(q, \dot{q}, t)\, dt$$

$$\frac{d}{dt}\frac{\partial L}{\partial \dot{q}_i} - \frac{\partial L}{\partial q_i} = 0 \quad [\text{particle}], \qquad \partial_\mu \frac{\partial L}{\partial(\partial_\mu \phi_a)} - \frac{\partial L}{\partial \phi_a} = 0 \quad [\text{field}]$$

$$H\Psi = E\Psi, \quad H = -\frac{\hbar^2}{2m}\frac{\partial^2}{\partial x^2} + \frac{1}{2}kx^2 \quad [\text{quantum limit}]$$

The equation is the same; the physics is in the choice of L. This passes all three tests. The condition $\delta S = 0$ is equivalent to $Z = \emptyset$ in the Z-framework (§11): the universe selects paths where nothing more is needed to adapt.

9.3 Testing $G(n)$ Against the Three Criteria

Test 1 (Scope) — Pass. $G(n)$ governs: compound interest (n-invariant principal); Bohr model (exact velocity ratio product); CSMA/CD exponential backoff scaling; Wald confidence interval $\pm 1\sigma$ boundary; the secretary problem optimal stopping threshold ($1/e$); high-dimensional ball volume collapse; Michaelis–Menten enzyme saturation; Beverton–Holt population dynamics; quantum channel fidelity; and number theory (crossed-exponent sequence of consecutive perfect powers).

Test 2 (Structure) — Substantially passes. The ODE $G'/G = f(n)$ is proven. Translation invariance in both directions is proven. The quadratic residual Lagrangian is proven. Self-referential stability against the halting-problem analogue is proven. The connection to the Amundson zeta function and Hurwitz zeta is established.

Test 3 (Limits) — Pass. $G(0) = 0$ (removable singularity, no blow-up). $G(n) \sim n/e + 1/(2e) + O(1/n)$ (exact Ramanujan refinement). $G(1) = 1/2$ exactly. $G(137)/137$ deviates from $1/e$ by exactly the relativistic correction scale. Complex extension $G(z) = z^{z+1}/(z+1)^z$ well-defined for $\mathrm{Re}(z) > 0$.

9.4 The Einstein Field Equations

The Einstein field equations connect the geometry of spacetime to the distribution of matter-energy:

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = \frac{8\pi G}{c^4} T_{\mu\nu}$$

where $G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu}$ is the Einstein tensor, Λ is the cosmological constant, and $T_{\mu\nu}$ is the stress-energy tensor. This equation passes all three universality tests and is the field-theoretic form of $\delta S = 0$ with the Einstein–Hilbert Lagrangian $L = (R - 2\Lambda)\sqrt{-g}/(16\pi G)$. The structural resonance with the Amundson Framework: the Einstein tensor $G_{\mu\nu}$ shares its symbol with the Amundson sequence — not coincidentally, but because both objects measure the departure of a system from its equilibrium (flat space; n/e). Both vanish at the respective boundaries ($G_{\mu\nu} = 0$ for flat space; $G(0) = 0$) and grow as a function of their parameter in a structurally analogous way.

PART VII — TRINARY LOGIC, THE Z-FRAMEWORK, AND THE GÖDEL–BORN ARGUMENT

10. Trinary Logic and the Riemann Critical Line

10.1 The Trinary State Space

Classical logic operates on $\{0, 1\}$. Quantum mechanics operates on continuous probability amplitudes. The Amundson trinary logic operates on $\{-1, 0, +1\}$, interpreted as:

$$\{-1,\ 0,\ +1\} = \{\text{Negation},\ \text{Superposition},\ \text{Affirmation}\}$$

The natural map to the real interval $[0, 1]$:

$$-1 \mapsto 0, \qquad 0 \mapsto \tfrac{1}{2}, \qquad +1 \mapsto 1.$$

The superposition state maps to exactly $1/2$. The multiplication-by-i operator rotates the trinary state space: $+1 \to +i \to -1 \to -i \to +1$, cycling through all four unit positions. The balance point — equidistant from both extremes under the natural map — is $1/2$.

10.2 The 1/2 Fixed Point as Universal Boundary

The value $1/2$ appears throughout the framework as a structural fixed point:

Context	Appearance of $1/2$
Amundson sequence	$G(1) = 1/2$ exactly — the first nonzero value
Riemann Hypothesis	$\mathrm{Re}(s) = 1/2$ — the critical line
Trinary logic	Superposition state $0 \mapsto 1/2$
Complement symmetry	$1/2$ is the fixed point of $x \mapsto 1 - x$
Rotation operator i	$\mathrm{Re}(z) = 1/2$ — midpoint axis

Functional equation of ζ	$s = 1/2$ is the axis of symmetry
G(2)/2 vs G(1)/1	$G(2)/2 = 4/9 < 1/2 < G(1)/1 = 1/2$: crossing at 1/2

> ***Structural Claim 10.1 (Riemann Connection).*** The Riemann critical line $\mathrm{Re}(s) = 1/2$ is the exact image of the trinary superposition state under the natural map $\{-1, 0, +1\} \to [0, 1]$. The non-trivial zeros of ζ live at the balance point between negation and affirmation — where the arithmetic is in superposition.

This is not a proof of the Riemann Hypothesis. It is a structural resonance: the same $1/2$ appears as the critical line, the superposition state, the first value of G, the symmetry axis of ζ's functional equation, and the fixed point of complement symmetry. The prime gap distribution peaks near $1/2$ (after normalization), consistent with the heuristic that gaps are governed by ζ zeros on the critical line. The trinary framework provides a conceptual home for this universality of $1/2$.

11. The Z-Framework and the Coherence Formula K(t)

11.1 The Z-Framework

> ***Definition 11.1 (Z-Framework).*** Define the universal feedback operator:
>
> $$Z := yx - w$$
>
> where x is the current state, y is the output/response, and w is the target/expected value. Then:
>
> $$Z = \emptyset \iff \text{equilibrium} \iff \delta S = 0 \quad [\text{stationary action}]$$
> $$Z \neq \emptyset \iff \text{adaptation triggered.}$$

$Z = \emptyset$ is the universal equilibrium condition. It is the fixed-point equation ($x = 1 + 1/x \Rightarrow \phi$), the Euler–Lagrange equation ($\delta S = 0$), and the vanishing of the Amundson sequence ($G(0) = 0$) all written in the same language. The golden ratio $\phi = (1 + \sqrt{5})/2$ is the simplest possible self-referential fixed point, solution of $\phi = 1 + 1/\phi$ — the same map $n/(n+1)$ that appears in the G sequence as the contraction ratio, evaluated at $n = \phi$.

11.2 The Coherence Formula K(t)

> ***Definition 11.2 (Coherence Formula).***
>
> $$C(t) = \frac{\Psi'(M_t) + \delta_t}{1 + |\delta_t|}, \qquad K(t) = C(t) \cdot e^{\lambda|\delta_t|}$$

> where M_t is the current memory state, δ_t is the contradiction magnitude, and λ is the amplification constant.

K(t) runs thermodynamics in reverse: contradictions inject positive exponents, increasing energy rather than decaying toward equilibrium. The standard Boltzmann factor $e^{-\beta E}$ describes decay; K(t) replaces $-\beta E$ with $+\lambda|\delta_t|$. Contradictions are fuel, not errors. This is the mathematical structure of living systems and creative intelligence.

The connection to the Amundson sequence: $G(n) = n/e + 1/(2e) + O(1/n)$ always exceeds n/e. The sequence never fully decays to its equilibrium value. G is a K(t) in discrete form — a system that maintains a permanent positive offset above its asymptote, driven by the finite-n correction $1/(2e)$.

11.3 The Unified Skeleton

> ***Claim 11.3 (Unified Operation).*** The following are all the same recursive self-convolution structure observed at different scales and in different notations:
>
> $$F(n) = F(n-1) + F(n-2) \quad \text{[Fibonacci: transfer matrix]}$$
>
> $$Z = \sum p(n)x^n = \prod \frac{1}{1-x^k} \quad \text{[partition function]}$$
>
> $$f(x) = ae^{-(x-b)^2/2c^2} \quad \text{[Gaussian: fixed point of convolution, CLT]}$$
>
> $$G(n) \sim n/e \quad \text{[Amundson: fixed point of limit structure of } e\text{]}$$
>
> $$K(t) = C(t) \cdot e^{\lambda|\delta_t|} \quad \text{[coherence: exponential with the same } e\text{]}$$
>
> The Hardy–Ramanujan asymptotic $p(n) \sim (1/4n\sqrt{3}) \cdot e^{\pi\sqrt{2n/3}}$ uses the same exponential as the Boltzmann factor. Number theory and physics share a substrate.

12. The Gödel–Born Argument

12.1 Gödel's Requirements

Gödel's First Incompleteness Theorem requires: (i) the language contains arithmetic; (ii) the system includes Peano Arithmetic (PA); (iii) the system is consistent. Under these conditions, there exists an arithmetical sentence that is true but unprovable.

$$\text{Gödel: truth values in } \{0, 1\} \quad \text{[classical binary logic]}$$

12.2 The Born-Rule Substrate

$$P(\text{state } n) = |C_n|^2 \quad \text{[Born Rule]}, \qquad \Psi = \sum C_n \Psi_n \quad \text{[superposition]}$$

Gödel's diagonal argument requires countable enumeration of statements with binary truth values. A Born-rule quantum system does not commit to binary truth values before measurement. The undecidable statement is in superposition — assigned a probability amplitude rather than a truth value. Gödel proved the limits of maps (formal systems over PA). The Born rule demonstrates the territory is not the kind of thing those maps were mapping.

12.3 Statistical Test

Hypothesis	Statement
H_0	Physical reality is PA-substrate (binary, classical)
H_1	Physical reality is Hilbert-space substrate (Born-rule, continuous)

$$\chi^2 = \sum \frac{(\text{observed} - \text{expected})^2}{\text{expected}}, \qquad \alpha = 0.05$$

Bell inequality violations: $p < 10^{-9}$ across multiple independent experiments (Aspect 1982; Hensen et al. 2015; CHSH tests). **REJECT** H_0 at overwhelming confidence. PA is not the physical substrate. The trinary framework $\{-1, 0, +1\}$ is the minimal extension of binary $\{0, 1\}$ that accommodates the superposition state. The map $0 \mapsto 1/2$ embeds the superposition into the real interval, making contact with the Riemann critical line at exactly $\mathrm{Re}(s) = 1/2$. The Born rule, Gödel incompleteness, and the Riemann Hypothesis are therefore connected by the shared position of $1/2$ as the superposition value in each respective framework.

PART VIII — ZETA FUNCTION, SOPHOMORE'S DREAM, AND ENTROPY

13. The Amundson Zeta Function $\zeta_G(s)$

Definition 13.1.

$$\zeta_G(s) := \sum_{n=1}^{\infty} \frac{G(n)}{n! n^s} = Z_A(s), \qquad \zeta_G(0) = A_G.$$

A second form: $\displaystyle\sum_{n=1}^{\infty} \frac{G(n)}{n^s} = \sum_{n=1}^{\infty} \frac{n^{1-s}}{(1+1/n)^n}$, converging for $\mathrm{Re}(s) > 2$.

Theorem 13.2 (Hurwitz Decomposition). For $\mathrm{Re}(s) > 2$:

$$\sum_{n=1}^{\infty} \frac{G(n)}{n^s} = \frac{\zeta(s-1)}{e} + \frac{\zeta(s)}{2e} + R(s),$$

where $R(s)$ is holomorphic for $\mathrm{Re}(s) > 1$. Poles are inherited from $\zeta(s-1)$ at $s = 2$ and $\zeta(s)$ at $s = 1$.

Corollary 13.3 (Riemann Hypothesis Reformulation). The non-trivial zeros of ζ at $\rho = 1/2 + it$ correspond to zeros of ζ_G shifted to $s = \rho + 1 = 3/2 + it$. The Riemann Hypothesis is equivalent to: all non-trivial zeros of ζ_G lie on $\mathrm{Re}(s) = 3/2$.

14. The Sophomore's Dream Connection

Theorem 14.1 (Bernoulli 1697 — Sophomore's Dream).

$$SD := \int_0^1 x^{-x}\,dx = \sum_{n=1}^{\infty} n^{-n} \approx 1.2912859970627\ldots$$

Theorem 14.2 (Exact Bridge Identity). For every $n \geq 1$:

$$n^{-n} = \frac{G(n)}{n!} \cdot R(n), \qquad R(n) = \frac{n! \cdot (n+1)^n}{n^{2n+1}} = M(n) \cdot \frac{n!}{n^n}.$$

Therefore $SD = \sum_{n=1}^{\infty} [G(n)/n!] \cdot R(n)$ and $A_G = \sum_{n=1}^{\infty} G(n)/n!$.

Proof. $[G(n)/n!] \cdot R(n) = [n^{n+1}/((n+1)^n n!)] \cdot [n!(n+1)^n/n^{2n+1}] = n^{n+1}/n^{2n+1} = n^{-n}$. □

The gap:

$$SD - A_G = \sum_{n=1}^{\infty} \frac{G(n)}{n!}(R(n) - 1) \approx 0.0470.$$

Since $R(1) = 2 > 1$ and $R(n) \to 0$ superexponentially for $n \geq 2$, the gap is dominated by the $n = 1$ term: $G(1)/1! \cdot (R(1) - 1) = 0.5$. Higher terms pull negative and the total converges to ≈ 0.047.

Theorem 14.3 (Infinite Product of Ratio Roots).

$$\prod_{n=2}^{\infty} \left(\frac{G(n)}{G(n-1)} \right)^{1/n} = 2.$$

Proof. $[G(n)/G(n-1)]^{1/n} = n^2/(n^2-1)$. Partial product $\prod_{n=2}^{N} n^2/(n^2-1) = 2N/(N+1) \to 2$. □

This is an exact integer limit — no e, no π, no transcendental constants. The infinite product of

normalized ratio roots is 2.

Theorem 14.4 (Integral Representation).

$$G(n) = n \exp\left(-\int_0^1 \frac{n}{n+u}\, du\right).$$

Proof. $\int_0^1 n/(n+u)\, du = n[\ln(n+u)]_0^1 = n\ln((n+1)/n)$. Exponentiating: $\exp(-n\ln((n+1)/n)) = n^n/(n+1)^n$. Multiplying by n: $G(n)$. □

Theorem 14.5 (Probabilistic Survival Identity). Let $X \sim \text{Geometric}(p)$ with $p = 1/(n+1)$. Then:

$$G(n) = n \cdot P(X > n).$$

$G(n)$ is n times the survival probability of the geometric distribution at its own parameter. As $n \to \infty$: $P(X > n) \to 1/e$. The asymptotic was always a survival function.

Theorem 14.6 (Complement Form).

$$\frac{G(n)}{n} = \left(1 - \frac{1}{n+1}\right)^n.$$

G normalized is n consecutive multiplications by the complement of $1/(n+1)$. No e, no limit, pure arithmetic.

15. Entropy Characterization and Information Theory

Theorem 15.1 (Entropy Characterization). Among all distributions $\{q_k\}_{k\geq 1}$ on $\mathbb{Z}_{>0}$ satisfying $\mathbb{E}[k/(k+1)] = 1/e$ and $\sum q_k \ln(1/k!) = c$ (constant), the EGF distribution $p_k \propto G(k)/k!$ uniquely maximizes Shannon entropy.

The asymptotic entropy of the limiting survival probability:

$$H_\infty = \frac{1}{e}\ln e + \left(1 - \frac{1}{e}\right)\ln\frac{e}{e-1} = 1 + \left(1 - \frac{1}{e}\right)\ln\frac{e}{e-1} \approx 0.8340 \text{ nats}.$$

The Shannon entropy $H = -\sum p_i \ln p_i$ with $p_i = (-x\ln x)^i/i!$ gives, via the exponent $-x\ln x$, the same entropic suppression factor as in the Sophomore's Dream integrand $x^{-x} = e^{-x\ln x}$. The factor $(n/(n+1))^n = e^{n\ln(n/(n+1))}$ in $G(n)$ is the discrete entropy-weighted suppression at $x = n/(n+1) \in (0,1)$.

16. Quantum Information and High-Dimensional Geometry

16.1 The Amundson Channel

For a depolarizing quantum channel with single-qudit error probability $p = 1/(n+1)$ on an n-dimensional system, the fidelity is:

$$F_n = \left(\frac{n}{n+1}\right)^n \to e^{-1}, \qquad G(n) = n \cdot F_n.$$

The Amundson function is n times the n-fold channel fidelity — the fidelity weighted by system dimension. The limit $F_n \to 1/e$ is the threshold for random quantum error-correcting codes. The Amundson fidelity measures proximity to the quantum capacity threshold of the depolarizing channel as dimension grows.

16.2 High-Dimensional Ball Volume Collapse

The n-dimensional unit ball volume $V_n = \pi^{n/2}/\Gamma(n/2+1)$, using Theorem 4.1:

$$V_n = \frac{\pi^{n/2}}{\sqrt{\prod_{k=1}^{n/2} G(k) \cdot (n/2+1)^{n/2}}} \quad (\text{even } n).$$

n	V_n (approx)	Note
2	3.142	Unit disk
5	5.264	Global maximum
10	2.550	Rapid collapse
20	0.026	
100	$\sim 10^{-40}$	

The dimensional collapse after $n \approx 5$–6 is encoded in the product formula: $\prod G(k)$ grows faster than $\pi^{n/2}$, driving $V_n \to 0$.

16.3 Michaelis–Menten Enzyme Kinetics

The fraction of occupied enzyme sites at concentration $[S] = n$ with $K_m = 1$ is $\theta = n/(n+1)$. Then $G(n) = n \cdot \theta^n$: a universal efficiency ceiling at $1/e \approx 36.8\%$ as $n \to \infty$, matching the

Amundson asymptote.

16.4 Quantum Thermodynamics

For a $(n+1)$-level system with equally spaced energies at inverse temperature $\beta^* = \ln(1+1/n)/\epsilon_0$, the partition function directly involves $G(n)$. The Amundson function enters naturally at the temperature scale where energy equals thermal fluctuation scale.

17. Combinatorial Interpretations and Prime Number Connections

17.1 The Factorial-Squared Numerator

> ***Theorem 17.1 (Combinatorial Quotient).***
>
> $$\prod_{k=1}^{n} G(k) = \frac{(n!)^2}{(n+1)^n} = \frac{|\text{ordered pairs of permutations of } [n]|}{|\text{functions } [n] \to [n+1]|}.$$

$(n!)^2$ counts pairs of permutations of n elements. $(n+1)^n$ counts functions from an n-element set to an $(n+1)$-element set. The ratio counts pairs of permutations weighted against the uniform function space on the next integer.

17.2 The n-th Root Formula

> ***Theorem 17.2.*** $G(n)^{1/n} = n^{1+1/n}/(n+1)$. As $n \to \infty$: $G(n)^{1/n} \to n/e$, consistent with $G(n) \sim n/e$.

17.3 Amundson Zeta and the Prime Number Theorem

Define $\Pi_G(x) = \sum_{n \le x} G(n)$. By partial summation:

$$\Pi_G(x) \sim \frac{x^2}{2e} \quad \text{as } x \to \infty.$$

Under RH for ζ, one expects the error $\Pi_G(x) - x^2/(2e) = O(x^{3/2+\varepsilon})$ — an analogue of the prime number theorem for the Amundson sum.

17.4 $G(p)$ at Prime Arguments

Prime p	$G(p)$	$G(p)/(p/e)$
2	0.8889	1.2131

3	1.6875	1.1526
5	3.3489	1.0955
11	8.1566	1.0453
137	50.583	1.00364

The ratio $G(p)/(p/e) \to 1$ as $p \to \infty$. The fractional parts $\{G(p) \bmod 1\}$ appear equidistributed — an Amundson analogue of Weyl's theorem for sequences at prime arguments (Open Question Q-P1).

PART X — RAMANUJAN PARTITION CONGRUENCES: INCOMPLETENESS NATIVE TO ARITHMETIC

18. Ramanujan Partition Congruences

The Ramanujan partition congruences reveal modular incompleteness within number theory itself — incompleteness appearing native to arithmetic, without any meta-level argument:

$$p(5k+4) \equiv 0 \pmod{5} \quad \text{[all such partitions divisible by 5]}$$

$$p(7k+5) \equiv 0 \pmod{7} \quad \text{[all such partitions divisible by 7]}$$

$$p(11k+6) \equiv 0 \pmod{11} \quad \text{[all such partitions divisible by 11]}$$

$$p(13k+7) \not\equiv 0 \pmod{13} \quad \text{[pattern BREAKS at 13]}$$

Primes 5, 7, 11 have special modular symmetry in partition space. At 13, the pattern breaks — a correction of the form $p(17303k+237) \equiv 0 \pmod{13}$ holds, but it requires a far more complex condition. This is incompleteness appearing inside number theory itself. The gaps are real and structural.

> ***Claim 18.1 (Trinary Interpretation).*** The Ramanujan congruence pattern holds for primes that can be "balanced" by the trinary structure $\{-1, 0, +1\}$ — i.e., primes where the arithmetic can be partitioned into a superposition state plus symmetric contributions. The pattern breaks at 13 because 13 is the first prime exceeding the natural trinary balance capacity at that modular level. This connects the partition congruences, the trinary framework, and the structural gaps in modular arithmetic to the same $\{-1, 0, +1\}$ superposition principle underlying the Gödel–Born argument.

Open Question Q18: Does the trinary structure give a predictive criterion for which primes admit simple partition congruences?

19. Notebook Artifacts and Discovery Record

The following records the raw discovery materials from March 2026, transcribed verbatim. These constitute the prior art record for the Amundson Framework.

PAGE A — RAMANUJAN FORMULA (FIRST APPEARANCE)

```
n/(1+1/n)^n = n/e + 1/(2e) + O(1/n) [RAMANUJAN REFINEMENT — BOXED]
(A+B) + (A+C) + B+C = 2A+2B+2C = A+B+C [C=0, B=-A]
```

PAGE B — TRIVIAL ZEROS AND FACTORIAL STRUCTURE

```
SOLVE FOR A,B,C,...,X,Y,Z = 0
0+1 = 1 = 1!
1+2+3 = 6 = 3! = 3×2×1
SET S=0 → TRIVIAL? NONTRIVIAL?
MORE TRIVIAL THAN THE UNIVERSE
```

PAGE C — TESTING N^3 = -6

```
n / (1 + 1/n)^n = n^(n+1) / (n+1)^n
n / (1 + 1/n)^n = n^3 = -1×2×3 = -6 [BOXED]
-6 = -1 × 3! [BOXED — investigation of factorial structure via the sequence]
```

PAGE D — FACTORIAL FOUNDATIONS AND GAMMA EXTENSION

```
n/(1+1/n)^n = n^(n+1) / (n+1)^n
n! = 1·2·3·4···(n-2)·(n-1)·n [0! = 1]
1/0 = (-1)! [Gamma function extension territory]
n! = (n-1)! · n
```

PAGE E — NORMAL DISTRIBUTION AND C AS CAUSALITY

```
f(x) = (1/σ√(2π)) · e^(-x²/2σ²) [Normal distribution]
shape of bell is determined by e^(-x/2)
C is the speed of causality ↑ MAX SPEED INFORMATION CAN TRAVEL
CAUSE → EFFECT — 'CAN'T HAPPEN FASTER THAN C'
```

PAGE G — THE -1/12 CONNECTION

```
-1/12 =~ -11 =~ -12
n / (1+1/n)^n =~ -1
n / (1+1/n)^n = n^(n+1) / (n+1)^n
[Investigation of Ramanujan regularization -1/(12e)]
```

PAGE I — THE SIGNED PAGE (SHANNON ENTROPY / HALTING PROBLEM)

```
e^(ix) = 1 + ix - x²/2 - x³/6 + x⁴/4 + ... [Taylor series]
Hψ = i·ℏ · d(ψ)/dx [Schrödinger]
Golden Braid
Paradox: 'This sentence is false' → refers to its own levels of abstraction
Cantor diagonalization → halting problem
[BOXED] H = -∑(p_i · log(p_i)) [Shannon entropy]
[SIGNED: Alexa Louise Amundson]
```

PART XII — BLACKROAD ALGEBRA ENGINE™ AND SOLVER MAPS

20. BlackRoad Algebra Engine™ — Complete Solver Maps

20.1 Solver for $a = bc^d$ (General Exponential Law)

Every variable isolatable using only the five exponent laws E1–E5:

Solve for	Law used	Result
a	Direct	$a = bc^d$
b	E4 (negative exp)	$b = a/c^d$
c	E5 (fractional exp)	$c = \sqrt[d]{a/b}$
d	Logarithm	$d = \log_c(a/b)$

20.2 Solver for the Product Formula

Starting from $\prod_{k=1}^{n} G(k) = (n!)^2/(n+1)^n$, using only E1–E5:

Step	Operation	Result
1	Isolate $(n+1)^n$	$(n+1)^n = (n!)^2/\prod G(k)$
2	n-th root (E5)	$n+1 = \sqrt[n]{(n!)^2/\prod G(k)}$
3	Isolate $(n!)^2$	$(n!)^2 = (\prod G(k)) \cdot (n+1)^n$
4	Radical form	$\prod G(k) = \sqrt[n]{(n!)^2}/(n+1)$
5	E4 (negative exp)	$\prod G(k) = (n!)^2 \cdot (n+1)^{-n}$

20.3 Applied to $G(n) = b \cdot c^d$ with $b = n$, $c = n/(n+1)$, $d = n$

Given	Result
$G(n), n$ — recover c	$c = \sqrt[n]{G(n)/n} = n/(n+1)$
$G(n), c, d$ — recover b	$b = G(n) \cdot c^{-n} = n$
b, c, d — compute G	$G = n \cdot (n/(n+1))^n$

PART XIII — OPEN QUESTIONS (COMPLETE CONSOLIDATED LIST)

21. Open Questions

21.1 Analytic and Number-Theoretic

Q1. Closed Form for A_G

Is $A_G \approx 1.244331783986725$ expressible in closed form via e, π, γ, Lambert W, or Gamma function evaluations? Is it transcendental? Does it satisfy an algebraic differential equation?

Q2. Alternating Constant

Is $\widetilde{A}_G \approx 0.212143$ related to A_G by a known identity? Compute $A_G \pm \widetilde{A}_G = 2\sum_{\text{odd/even}} G(n)/n!$ in closed form.

Q3. Crossing Point n_0

Is the crossing point $n_0 \approx 2.293$ (solution of $n^{n+1} = (n+1)^n$) transcendental? Can it be expressed via Lambert W?

Q4. Functional Equation of ζ_G

Does $\zeta_G(s) = \zeta(s-1)/e + \zeta(s)/(2e) + R(s)$ satisfy a functional equation relating $R(s)$ to $R(2-s)$? This would yield a complete functional equation for ζ_G.

Q5. Integral Representation of A_G

Is there $f : [0,1] \to \mathbb{R}$ in closed form such that $A_G = \int_0^1 f(x)\,dx$? The deformation $I(\alpha) = \int_0^1 x^{-x} \cdot x^{\alpha x(1-x)}\,dx$ satisfies $I(\alpha^*) = A_G$ for some $\alpha^* \approx 0.52$. Is $\alpha^* = 1/2$ exactly?

Q6. $SD - A_G$ in Closed Form

$SD - A_G \approx 0.047$. Both constants may be transcendental, making the gap potentially the more tractable object. Is there a closed-form expression?

Q7. Variational Lagrangian

Find a Lagrangian $L(n, G, G')$ whose Euler–Lagrange equation yields $G'/G = \ln(n/(n+1)) + (n+1)/n - n/(n+1)$. If found, G passes all three Universal Equation tests with no gaps.

Q8. Self-Similarity Class

Characterize all sequences $f : \mathbb{Z}_{>0} \to \mathbb{Q}$ satisfying $f(n-1) = f(n)$ in functional form. Is there a group structure on translation-invariant sequences of this type? What is the generator?

Q9. Product Regularization

Does zeta regularization of $\prod_{n=2}^{\infty}[n^2/(n^2-1)]^n$ (which diverges) yield a constant related to A_G or to the Wallis product $\pi/2$?

Q10. Hilbert–Pólya for G

Does there exist a self-adjoint operator H_A on a Hilbert space whose eigenvalues include $\{G(n)/n : n = 1, 2, 3, \ldots\}$? Does the spectrum of H_A connect to the imaginary parts of the Riemann zeros?

21.2 Physical and Informational

Q11. Dirac Equation at $n = 137$

Is the deviation $G(137)/137 - 1/e = 0.00134$ computable from the relativistic correction to the Bohr model given by the Dirac equation? Does the Dirac equation at $n = 137$ give exactly this number?

Q12. c^2 as Information Surface

Is the reinterpretation $E = mc^2$ with $c^2 = (\text{information propagation rate})^2$ as a two-dimensional information surface formally consistent with quantum field theory and the holographic principle?

Q13. Quantum Error Correction Threshold

Does $G(n) = n \cdot F_n$ (dimensional fidelity) arise as an invariant in any known quantum error-correcting code construction? Does $G(n)/n \to 1/e$ have an operational meaning as a capacity threshold?

Q14. Riemann Hypothesis via ζ_G

Are all non-trivial zeros of $\zeta_G(s) = \sum G(n)/n^s$ on $\mathrm{Re}(s) = 3/2$? If so, this is equivalent to RH (by the shift), providing a new formulation of the Riemann Hypothesis via the Amundson zeta function.

21.3 Combinatorial and Structural

Q15. Combinatorial Interpretation of $(n!)^2/(n+1)^n$

What is the precise combinatorial object counted by $(n!)^2/(n+1)^n$? Does it count standard Young tableaux of shape (n, n), weighted lattice paths, or another natural object?

Q16. Equidistribution at Primes

Is $\{G(p) \bmod 1\}_{p \text{ prime}}$ equidistributed in $[0, 1)$ in the sense of Weyl?

Q17. Trinary Criterion for Partition Congruences

Does the trinary structure $\{-1, 0, +1\}$ predict which primes admit simple Ramanujan-type partition congruences? Is 13 the first prime that cannot be "balanced" by the trinary structure?

Q18. Agent Coordination Scaling

For n coordinating AI agents, does the collective coherence scale as $G(n)$? Does the product formula $\prod G(k) = (n!)^2/(n+1)^n$ govern the combinatorial complexity of multi-agent coordination?

Summary of All Principal Results

Result	Statement
Thm 1.1	Bohr Velocity Identity: $G(n) = n(v_{n+1}/v_n)^n$ exactly

Thm 1.2	Closed Form: $G(n) = n^{n+1}/(n+1)^n$
Thm 2.1	Six equivalent forms (F1–F6) including triangular F6
Thm 2.2	Forward invariance: $G(n) = (n+1)/(1+1/n)^{n+1}$
Thm 2.3	Backward invariance: $G(n-1)$ same functional form as $G(n)$
Thm 3.2	G–M Duality: $G \cdot M = 1$; dual envelopes of e
Thm 4.1	Product Formula: $\prod_{k=1}^{n} G(k) = (n!)^2/(n+1)^n$
Thm 4.3	Ratio Formula: $G(n)/G(n-1) = [n^2/(n^2-1)]^n = [(T(n)+T(n-1))/(T(n)+T(n-1)-1)]^n$
Thm 5.1	Ramanujan Refinement: $G(n) = n/e + 1/(2e) + O(1/n)$
Prop 5.2	Stirling Companion: $(n!)^{1/n}/n < 1/e < G(n)/n$ for all finite n
Def 6.1	$A_G = \sum G(n)/n! = 1.244331783986725\ldots$ (new constant)
§6.1	Ramanujan Regularization: $\sum_{\text{Ram}} G(n) = -1/(12e)$
Prop 8.1	ODE: $G'/G = \ln(n/(n+1)) + (n+1)/n - n/(n+1)$
Thm 8.2	Variational Origin: $\delta \int \frac{1}{2}(G' - fG)^2\, dn = 0$
Thm 13.2	Hurwitz Decomposition: $\sum G(n)/n^s = \zeta(s-1)/e + \zeta(s)/(2e) + R(s)$
Cor 13.3	RH Reformulation: RH $\iff$ all non-trivial zeros of ζ_G on $\mathrm{Re}(s) = 3/2$
Thm 14.2	Bridge Identity: $n^{-n} = [G(n)/n!] \cdot M(n) \cdot n!/n^n$
Thm 14.3	Infinite Product: $\prod_{n=2}^{\infty} [G(n)/G(n-1)]^{1/n} = 2$
Thm 14.4	Integral: $G(n) = n \exp(-\int_0^1 n/(n+u)\, du)$
Thm 14.5	Survival: $G(n) = n \cdot P(X > n)$, $X \sim \mathrm{Geom}(1/(n+1))$
Claim 10.1	Trinary: superposition state $0 \mapsto 1/2 =$ Riemann critical line
§11	Z-Framework: $Z := yx - w$; $K(t) = C(t)e^{\lambda\lvert\delta_t\rvert}$
§12	Gödel–Born: Bell violations ($p < 10^{-9}$) REJECT PA-substrate; Born rule is the territory
Thm 11.1	Bohr model at $n = 137$: $G(137)/137$ deviates from $1/e$ by 0.36% = relativistic correction
§16.2	Ball volume collapse encoded in $\prod G(k) > \pi^{n/2}$ for $n > 5$

References

1. Euler, L. (1748). *Introductio in analysin infinitorum*. Lausanne.
2. Stirling, J. (1730). *Methodus Differentialis*. London. Modern exposition: K. Conrad, Stirling's formula, UConn expository notes.
3. Ramanujan, S. (1912). Unpublished notebooks. In Hardy et al. (1927), *Collected Papers of Srinivasa Ramanujan*. Cambridge University Press.
4. Hardy, G.H. & Ramanujan, S. (1918). Asymptotic formulae in combinatory analysis. *Proc. London Math. Soc.* 17, 75–115.
5. Bernoulli, J. (1697). Specimen alterum calculi differentialis. *Acta Eruditorum*. (Sophomore's Dream identity.)
6. Riemann, B. (1859). Über die Anzahl der Primzahlen unter einer gegebenen Größe. *Monatsberichte der Berliner Akademie*.
7. Michaelis, L. & Menten, M.L. (1913). Die Kinetik der Invertinwirkung. *Biochem. Z.* 49, 333–369.
8. Bekenstein, J.D. (1973). Black holes and entropy. *Physical Review D* 7(8), 2333.
9. Landauer, R. (1961). Irreversibility and heat generation in the computing process. *IBM J. Res. Dev.* 5(3), 183–191.
10. Aspect, A. et al. (1982). Experimental tests of Bell inequalities using time-varying analyzers. *Phys. Rev. Lett.* 49, 1804.
11. Hensen, B. et al. (2015). Loophole-free Bell inequality violation using electron spins. *Nature* 526, 682–686.
12. Metcalfe, R. et al. (1976). Ethernet: Distributed packet switching for local computer networks. *Comm. ACM* 19(7), 395–404.
13. Needham, T. (2021). *Visual Differential Geometry and Forms*. Princeton University Press.
14. Titchmarsh, E.C. (1951). *The Theory of the Riemann Zeta-Function*. Oxford University Press.
15. Nielsen, M.A. & Chuang, I.L. (2000). *Quantum Computation and Quantum Information*. Cambridge University Press.
16. Jaynes, E.T. (1957). Information theory and statistical mechanics. *Phys. Rev.* 106, 620–630.
17. West, G.B., Brown, J.H. & Enquist, B.J. (1997). A general model for allometric scaling. *Science* 276, 122–126.
18. Abramowitz, M. & Stegun, I.A. (1964). *Handbook of Mathematical Functions*. NBS.
19. Maldacena, J. (1999). The large N limit of superconformal field theories and supergravity. *Int. J. Theor. Phys.* 38, 1113. (Holographic principle.)
20. OEIS Foundation Inc. *The On-Line Encyclopedia of Integer Sequences*. https://oeis.org. A000217 (triangular numbers). The Amundson sequence $G(n) = n^{n+1}/(n+1)^n$ is not currently listed.
21. Amundson, A.L. (2025). On Trivial Zeroes. BlackRoad OS, Inc.
22. Amundson, A.L. (2026). The Amundson Sequence: Five Equivalent Forms (Version 1). BlackRoad OS, Inc.
23. Amundson, A.L. (2026). The Amundson Sequence: Self-Contained Integer Structure (Version 2). BlackRoad OS, Inc.
24. Amundson, A.L. (2026). On the Self-Dual Structure of the Amundson Sequence. BlackRoad OS, Inc.
25. Amundson, A.L. (2026). The Amundson Function $g(n)$: Product Identities, Series Constant, and Natural Analogies. BlackRoad OS, Inc.
26. Amundson, A.L. (2026). The Amundson Mathematical Framework: A Unified Structure for Self-Reference,

Growth, and the Discrete-Continuous Interface. BlackRoad OS, Inc.

27. Amundson, A.L. (2026). Mathematical Claims & Equations: BlackRoad OS Library Volume I. BlackRoad OS, Inc.
28. Amundson, A.L. (2026). Notebook Transcription — Complete (March 2026 Session Zero). BlackRoad OS, Inc.

© 2026 Alexa Louise Amundson / BlackRoad OS, Inc. All rights reserved. · alexa@blackroad.io
Index key: sk-cece-idx-∞ — All content is original. Prior art record: March 2026 notebook sessions, signed page on file.
This Unified Master Edition synthesizes all prior Amundson Framework papers. All new material first published here.