

Canadian Consensus Meeting on Cyclosporine Monitoring: Report of the Consensus Panel

Leslie M. Shaw,¹ Randall W. Yatscoff,^{2,3} Larry D. Bowers,⁴ David J. Freeman,⁵ John R. Jeffery,³ Paul A. Keown,⁶ Iain J. McGilveray,⁷ Thomas G. Rosano,⁸ and Pui-Yuen Wong⁹

The Bureau of Drug Research, Health Protection Branch of the Government of Canada, requested of the transplant and clinical chemistry professional groups that a consensus meeting be held, to develop a "state-of-the-art" consensus paper on cyclosporine (CsA) monitoring. The often confusing scientific literature on CsA monitoring, created largely as a result of the use of different nonspecific and specific methods for measurement of CsA, the use of different sample matrices in which the drug is measured, and the varied criteria used for defining renal toxicity or rejection had led to this request. In response, the Canadian Society of Clinical Chemists in conjunction with the Canadian Transplant Society planned and organized the Canadian Consensus Meeting, which was held at Minaki, Ontario, May 11 through 13, 1990. Twenty-one scientists representing the clinical, methodological, and pharmacological aspects of CsA presented to 72 attendees the latest research data on mechanisms of the immunosuppressive and nephrotoxic activity of CsA; immunosuppressive and nephrotoxic activities of the major human metabolites of CsA; distribution of CsA between blood plasma, blood cells, and tissues; specific methods for measuring CsA; the impact on clinical outcome of adjusting CsA dosage on the basis of results of monitoring the parent drug concentration in whole blood; CsA pharmacokinetics; and methods for detecting rejection. Based on a detailed consideration of all data pertinent to the therapeutic drug monitoring of CsA, a set of recommendations was developed, as requested by the Canadian Health Protection Branch, by a consensus panel consisting of the nine individuals who co-authored this paper. At the conclusion of the meeting these recommendations and the rationale for them were presented for critical review to all who attended the meeting.

During the conference there was considerable debate regarding the following: (a) the appropriateness of one sample matrix over another (whole blood, plasma, serum,

or plasma ultrafiltrate) for measuring CsA in transplant patients, and (b) the relative merits of the area-under-the-curve method compared with trough concentrations for regulation of CsA dosing. Although all of the transplant centers represented at the meeting use CsA monitoring to adjust dosage, several investigators in the transplant field still believe that adjusting CsA dosage in response to changes in clinical indices provides good long-term clinical outcomes in transplant patients. Future studies will be required to resolve this issue.

It is the hope of the consensus panel that the following recommendations and the rationale for them will provide the type of practical information that could foster an improvement in the center-to-center consistency in the practice of cyclosporine therapeutic drug monitoring. This report is therefore regarded as a complement to the Hawk's Cay consensus report (1), which described the areas of controversy but did not make specific recommendations on the time of sample collection, the method for measurement of CsA, or the sample matrix in which to measure the drug.

Recommendations and Guidelines

- 1) The recommended time for collection of the sample in which to measure CsA is 11–12 h after the last CsA dose for patients receiving two doses per day and 23–24 h after the last CsA dose for patients receiving one dose per day.
- 2) It is recommended that the time of food intake in relation to CsA dosing be consistent, because effects of food on CsA absorption can be considerable.
- 3) Information as to the amount and time of administration of the last dose should be submitted together with the request for CsA determination.
- 4) In most clinical situations during the immediate post-transplant period, the maximum recommended frequency of monitoring is once every 24 h.
- 5) Until one matrix is shown to be clinically superior to another, analytical reasons, including the pressing need for center-to-center consistency, should be the basis for selection of sample matrix. This is the basis for selection of whole blood as the recommended sample. EDTA is the recommended anticoagulant. It is important to assure artifact-free collection of the blood specimen. Therefore, for patients receiving the drug through an in-dwelling catheter, venipuncture from the contralateral arm is required.
- 6) In vitro, CsA has long-term stability over a wide range of temperatures. If required, samples can be shipped at ambient temperatures.
- 7) Use of a validated specific method for measuring the parent drug is recommended.
- 8) Performance characteristics for CsA measurements should meet the criteria for acceptable clinical laboratory practice.
- 9) No data currently support the monitoring of CsA

¹ Department of Pathology and Laboratory Medicine, The University of Pennsylvania Medical Center, Philadelphia, PA 19104.

Departments of ² Clinical Chemistry and ³ Medicine, Health Sciences Centre, University of Manitoba, Winnipeg, Manitoba, Canada.

⁴ Department of Laboratory Medicine and Pathology, University of Minnesota, Minneapolis, MN.

⁵ Roberts Research Institute, University of Western Ontario, London, Ontario, Canada.

⁶ Departments of Pathology and of Medicine, Vancouver General Hospital, University of British Columbia, Vancouver, BC, Canada.

⁷ Health Protection Branch, Ottawa, Ontario, Canada.

⁸ Department of Laboratory Medicine, Albany Medical Center, Albany, NY.

⁹ Department of Clinical Biochemistry, University of Toronto, Toronto, Ontario, Canada.

Received May 29, 1990; accepted July 30, 1990.

metabolites as a guide to dosage adjustment in transplant patients. Should one or more metabolites subsequently be shown to be clinically important, it is recommended that they be measured with a specific method.

10) Target or therapeutic ranges have been empirically derived. Their definition is subject to all of the difficulties resulting from multiple confounding factors. Tables are provided listing representative ranges and factors such as the method used for measurement of CsA that may influence the derivation of each target range and, where available, the criteria used for defining rejection and nephrotoxicity.

11) It is recommended that laboratories provide same-day (before the next dose) turnaround assay for acute transplant patients.

12) To maintain the quality of the method for measuring CsA, participation in an external quality-improvement program is essential in addition to an effective internal quality-assurance program. It is recommended that a pure, well-characterized CsA preparation be developed to make possible the effective standardization of CsA measurements from laboratory to laboratory.

Time of Sample Collection

Most transplant centers throughout the world specify that the time for collection of the specimen in which CsA is to be measured be just before administration of the next dose (2). As is the case for the majority of therapeutic drugs, this selection of the trough time is based on its reproducibility; it occurs after absorption of the drug is complete, which places it in a more reproducible part of the drug concentration-vs-time curve; CsA toxicity correlates better with trough concentrations than with either peak or peak minus trough increments (3). Several investigators have proposed other sample times: 5 or 6 h after the last dose (1) or multiple timed samples collected throughout the dosage interval for determination of the area under the CsA concentration in blood-vs-time curve (AUC), and the average steady-state concentration (4, 5). On the basis of this information, standard pharmacokinetic formulas are used to determine what dosages are subsequently required to achieve target steady-state concentrations of CsA. Future studies are needed to test the value of this approach by directly comparing it with the standard trough-concentration approach in relation to achieving target concentrations in blood and the desired clinical outcome. The major practical limitations of AUC monitoring are the need for extended hospitalization or clinic visits, the frequency of blood sampling required each time the patient's CsA status is to be evaluated, and the cost. Despite these important practical difficulties this proposal has the potential advantage of providing a pharmacokinetically more rigorous approach to determining the patient's subsequent dosage regimen (4-6). In situations where it is difficult to attain the desired trough concentrations, pharmacokinetic profiling has been helpful in defining the nature of the problem (e.g., poor absorption or abnormal clearance of CsA) and in establishing an effective dosage regimen (2).

Sample Stability

CsA in EDTA-anticoagulated blood, without any additives or preservatives, is stable at either room temperature or at 2-8 °C for up to seven days. Samples can then be shipped at ambient temperatures to central processing locations. For prolonged storage, samples should be kept at

-20 °C. In the presence of preservatives, samples are stable for up to four months at room temperature (7).

Sample Matrix

To date, there are neither pharmacological nor clinical studies in which CsA concentrations, measured specifically in different sample matrices (whole blood, plasma, serum, or plasma ultrafiltrate) prepared from individual patients, were compared to determine which of these correlates best with pharmacological effects and clinical events such as rejection or nephrotoxicity. It is difficult to draw meaningful conclusions about the comparative clinical value of different sample matrices from the literature because of confounding factors, e.g., the use of different methods for CsA determination, different dosing intervals, the temporal relationship between clinical events and monitoring CsA concentration, the concomitant use of other immunosuppressive agents, the lack of "gold standards" for use in determining the degree of immunosuppression and nephrotoxicity, and the presence of other risk factors that may predispose a transplant patient to nephrotoxicity (1, 2, 8, 9).

Until appropriate comparative studies are performed, the primary factors upon which a rational choice of sample matrix can be based are analytical factors and the need for center-to-center consistency of results. The analytical reasons for recommending whole blood as the sample matrix are the greater signal-to-noise ratio made possible by the higher (twofold on average) concentration in whole blood as compared with plasma prepared at 37 °C, and the requirement that plasma be separated from blood cells under stringent conditions of time and temperature. The use of plasma prepared under different temperature and (or) time conditions for CsA determination would require multiple target ranges for therapeutic drug monitoring. The use of whole-blood CsA concentrations makes it easier to compare results from center to center. Furthermore, for transplant patients who may need to have their CsA measured in different laboratories, measurement of concentrations in whole blood will give the most comparable results, especially when validated liquid-chromatographic or specific monoclonal-antibody immunoassays are used for the measurement (2, 8, 10).

Method for Measurement

The specific measurement of CsA in which only parent drug and not the metabolites are quantified can be accomplished by two procedures: "high-pressure" liquid chromatography (HPLC) and immunoassays involving a selective monoclonal antibody (11-16).

The former method, when well-validated, has been considered the reference method for the specific measurement of the drug (1, 2, 8). Several reversed-phase HPLC procedures for CsA determination have been described (8, 17-24). The major differences among them are the sample preparation procedure, the sample volume, the HPLC column, and the mobile phase. The absolute analytical recovery of the drug should be >80%, with the recovery relative to an internal standard being in the range of 90-110%.

Performance Characteristics

The error for the method (either HPLC or selective immunoassay) at 50 µg/L should be no greater than 10 µg/L and at a concentration of 300 µg/L should be no greater than 30 µg/L. This requires that the standard

deviation of the analytical method be less than or equal to 5 and 15 $\mu\text{g/L}$, at those respective concentrations. The rationale for this is that the concentration of CsA measured in blood has been decreasing as clinical protocols have been developed. In addition, several centers have shown that concentrations that decrease with time after the transplant are efficacious (25). A recent report indicates that for one center the median CsA concentration observed with a specific assay was 135 $\mu\text{g/L}$, with 33% of the samples having concentrations less than 100 $\mu\text{g/L}$ (26). Medical decisions regarding therapy are being made at the 100 $\mu\text{g/L}$ level.

Results by the analytical method (either HPLC or immunoassay) should be compared with those by a validated HPLC procedure as the reference method (x -axis) for at least 200 specimens of blood from persons with all transplant types (kidney, liver, pancreas, heart, heart-lung, bone marrow). Standard statistical evaluation (see reference 27, for example) of the linear-regression data should be used, including, but not limited to, the slope and standard deviation of the slope, the y -intercept and standard deviation of the intercept, and the standard deviation of the estimate ($s_{y/x}$, standard error of the regression, standard error of the residuals). An appropriate statistical test (t -test) should be performed on the slope and intercept to determine that they are not statistically different from 1.00 or 0.0, respectively, at the 95% confidence interval. We recommend that the slope differ by no more than 10% from the line of identity (slope = 1.00; acceptable 0.90–1.10), the intercept by no more than 15 $\mu\text{g/L}$, or the $s_{y/x}$ by no more than 15 $\mu\text{g/L}$.

The analytical method should be validated for accuracy. The validation of the assay depends on the availability of pure CsA standard material. Standards are available from the Sandoz Co. and from the United States Pharmacopoeia. The latter material has been checked for purity and assayed in several reference laboratories and, thus, is preferred. Precautions should be taken when drying and desiccating the reference material before standards are prepared.

The consensus meeting panel strongly advocates development of a secondary reference material in a whole-blood matrix, assayed by reference laboratories, to assist in improving accuracy of CsA testing. A potentially promising reference procedure for confirming the CsA concentration in this reference material would be liquid chromatography/mass spectroscopy, with deuterated CsA as an internal standard.

The analytical recovery of CsA and the internal standard should be determined in the assay—absolute recovery, not recovery relative to the internal standard. It has been shown that recoveries of CsA and the common internal standards cyclosporin D and dihydrocyclosporin C are not the same when hexane washes are used in the analysis (28, 29).

A set of "challenge samples" containing both metabolite and parent drug, and preferably in a whole-blood matrix, should be analyzed. The concentration of CsA in the challenge sample should be determined by reference laboratories. These could be part of an external quality-assurance program, such as the Canadian Quality Assessment Program. In validating the assay, at least 10 such challenge samples should be run. Thereafter, the consensus panel recommends the analysis of at least one set of challenge samples per year.

Cyclosporine Metabolites

More than 20 metabolites of CsA have been identified (30–39). Several studies have been published during the past few years in which the immunosuppressive (32–34, 37, 40–45) and toxic activities (46–49) of these metabolites have been investigated. Although further investigation is still required, some conclusions can be made from the data currently available.

In vitro experiments indicate that metabolites are considerably less immunosuppressive and toxic than is CsA. The immunosuppressive and toxic activities of the metabolites appear not to be associated (47). In vivo studies in animals have been hampered by the lack of sufficient quantities of metabolites and of a suitable model (43, 50). Preliminary results, obtained with rats, suggest that the metabolites are less immunosuppressive than CsA, which corroborates the in vitro findings. Clinical studies have indicated that the routine monitoring of metabolites is not warranted for the majority of clinical situations (51–57). If future data indicate a clinical requirement for metabolite measurement, then a specific measurement for each metabolite of significance would be warranted.

Therapeutic Range

Historically, the therapeutic range for CsA has been difficult to establish, due in large part to the variation in matrix and selectivity of the assays as well as the clinical criteria used for rejection and toxicity (2, 30, 58–70). Taking into account two of the recommendations proposed here—namely, the use of selective assays with standards and samples in a whole-blood matrix for routine monitoring—should facilitate the establishment of a consensus therapeutic range. Table 1 lists therapeutic ranges for solid-organ transplantation from several Canadian and United States centers where whole blood and selective assays are used. Where available, the criteria for rejection and CsA toxicity are also listed (71–75). It is important that each transplant center, before adopting a reference range established elsewhere, ensure that the methods of analysis and the criteria for diagnosis of clinical events are the same as those originally used to establish this range. The institution should then confirm the adequacy of this reference range for use in their own transplant population.

Quality Assurance

Several quality-assessment programs have described their experience to date regarding center-to-center reproducibility when a given method is used for measuring CsA, the recovery of cyclosporine added to drug-free or pooled specimens of whole blood from the transplant patients, and the detection of CsA in samples of pooled drug-free whole blood (7). These programs have been helpful in providing feedback to participating laboratories about the consistency of their performance and about how their performance compares with that of other laboratories. In the future it will be important for these programs to provide suggestions as to the possible causes of a particular problem and how to improve performance. One further recommendation is the development of a well-characterized CsA preparation in whole blood to provide all laboratories a common source of a single material against which to compare CsA standards in routine use.

We thank Kenneth Copeland for helping to gather data for this paper and Marie Simoens for secretarial assistance.

Table 1. Comparison of Therapeutic Ranges among Various Transplant Centers Using a Whole-Blood Matrix

Center	Analytical method	Dosing interval	Sample time	Other immuno-suppressants	Therapeutic range, $\mu\text{g/L}$	Transplant	Diagnosis of rejection and (or) toxicity
Health Sciences Ctr. Winnipeg, MB	Selective RIA	BID	Trough	Aza, Pred	200–300 <1 m 100–200 >1 m	K	Biopsy, >15% \uparrow Cr
Hosp. of the Univ. of Pennsylvania Philadelphia, PA	HPLC	QD	Trough	Aza, Pred	100–250 <3 m 80–125 >3 m 250–350 <3 m 100–150 >3 m	K K L L, H	Biopsy, >25% \uparrow Cr
Mayo Clinic Rochester, MN	HPLC HPLC	QD BID	Trough Trough	Pred Pred	250–350 150–250 <2 m 80–200 >2 m	L K	—
Ottawa General Hosp. Ottawa, ON	Selective RIA	BID	Trough	Aza, Pred	>200 post-tx <200 >1 w	K	Biopsy, ultrasound, \uparrow Cr
St. Joseph's Hosp. Hamilton, ON	Selective RIA	BID	Trough	Aza, Pred	150–250 <3 m 100–150 >3 m	K	Biopsy, ultrasound, >10% \uparrow Cr
Toronto General Hosp. Toronto, ON	Selective RIA	BID	Trough	Aza, Pred	100–300 100–400	K L	Biopsy, \uparrow Cr
University Hospital London, ON	Selective RIA	BID	Trough	Pred	250–450 <3 m 100 >3 m 400 <3 m 200–300 3–12 m 100–200 >1 y 300–600 <3 m 200–350 3–12 m 100–200 >1 y	K H L	Biopsy, 10–20% \uparrow Cr
Univ. of Cincinnati Cincinnati, OH	HPLC	BID	Trough	Aza, Pred	150–250 <6 m 75–150 >6 m 150–300 >6 m 75–150 >3 m	K K L H	Biopsy, >10% \uparrow Cr
Univ. of Minnesota Minneapolis, MN	HPLC	BID	Trough	Aza, Pred	150–200 <1 m 100–150 1–12 m 50–100 >12 m 200–250 200–300	K K K L H	Biopsy, >25% \uparrow Cr
Univ. of Pittsburgh Pittsburgh, PA	HPLC	BID	Trough	Aza, Pred	150–250	K	—
Univ. of Texas Medical Center Houston, TX	Selective RIA	QD	AUC	Pred	370 target	K	—
Vancouver General Hosp. Vancouver, BC	Selective RIA	BID	Trough	Aza, Pred	300–400 <1 w 250–400 <1 m 200–300 <3 m 100–200 <6 m 100–150 <2 y 300–600 <1 w 200–450 <1 m 200–350 <3 m 200–300 <6 m 150–250 <1 y 300–450 <1 w 300–400 <1 m 200–300 <3 m 100–200 <2 y	K L H	Biopsy, >10% \uparrow Cr >10% \uparrow Cr >10% \uparrow Cr
Victoria General Hosp. Halifax, NS	Selective RIA	BID	Trough	Aza, Pred	250–350 <2 w 200–250 <3 m 150 target >3 m	K	Biopsy, \uparrow Cr

K: kidney transplant recipients; L: liver transplant recipients; H: heart transplant recipients; Aza: azathioprine; Pred: prednisone; Cr: creatinine; QD: once daily; BID: twice daily; AUC: area under the curve; m: month; w: week; y: year; tx, transplantation.

References

- Kahan BD, Shaw LM, Holt D, Grevel J, Johnston A. Consensus document: Hawk's Cay Meeting on therapeutic drug monitoring of cyclosporine. *Clin Chem* 1990;36:1510-6.
- Shaw LM, Bowers LD, Demers L, et al. Critical issues in cyclosporine monitoring: report of the task force on cyclosporine monitoring [Special Report]. *Clin Chem* 1987;33:1269-88.
- Kahan BD, Van Buren CT, Nan Lin S, et al. Immunopharmacological monitoring of cyclosporin A-treated recipients of cadaveric kidney allografts. *Transplantation* 1982;34:36-45.
- Grevel J, Welsh MS, Kahan BD. Pharmacokinetic monitoring of cyclosporine therapy. *Transplant Proc* 1990;22:in press.
- Awni WM, Heim-Duthoy K, Kasiske BL. Monitoring of cyclosporine by serial post-transplant pharmacokinetic studies in renal transplant patients. *Transplant Proc* 1990;22:1343-4.
- Kahan BD, Grevel J. The benefit of pharmacokinetic monitoring of cyclosporine therapy. *Clin Biochem*, in press.
- Wong PY, Mee AV, Glenn J, Keown PA. Quality assessment of cyclosporine monitoring—Canadian validation. *Transplant Proc* 1990;22:1216-7.
- Vine W, Bowers LD. Cyclosporine: structure, pharmacokinetics and therapeutic drug monitoring [Review]. *Crit Rev Clin Lab Sci* 1987;25:275-311.
- Kahan BD, Mickey R, Flechner SM, et al. Multivariate analysis of risk factors impacting on immediate and eventual cadaver allograft survival in CsA-treated recipients. *Transplantation* 1987;43:65-70.
- Annesley TM, Giacherio DA, Feldkamp CS. Cyclosporine distribution in blood: concomitant effect of hematocrit, concentration of metabolites and temperature. *J Clin Immunoassay* 1987;10:226-31.
- Shaw LM. Cyclosporine monitoring [Editorial]. *Clin Chem* 1989;35:5-6.
- Ball PE, Munzer H, Keller HP, et al. Specific ^3H radioimmunoassay with a monoclonal antibody for monitoring cyclosporine in blood. *Clin Chem* 1988;34:256-60.
- Copeland KR, Yatscoff RW. Use of a monoclonal antibody for the therapeutic monitoring of cyclosporine in plasma and whole blood. *Ther Drug Monit* 1988;10:453-8.
- Sgoutas DE, Hammarstrom M. Comparison of specific radioimmunoassays for cyclosporine. *Transplantation* 1989;47:668-70.
- Schroeder TJ, Brunson ME, Pesce AJ, et al. A comparison of the clinical utility of the radioimmunoassay, high-performance liquid chromatography and Tdx cyclosporine assays in outpatient renal transplant recipients. *Transplantation* 1989;47:262-6.
- Tredger JM, Steward CM, Williams R. Cyclosporine blood levels and evaluation of radioimmunoassay with selective monoclonal or polyclonal antibodies and high-performance liquid chromatography in liver transplant recipients. *Transplantation* 1988;46:681-6.
- Furlanut M, Plebani M, Burlina A. Cyclosporin monitoring: methodological considerations. *J Liq Chromatogr* 1989;12:1759-89.
- Sawchuk RJ, Cartier LL. Liquid-chromatographic determination of cyclosporin A in blood and plasma. *Clin Chem* 1981;27:1366-71.
- Niederberger W, Schaub P, Beveridge T. High-performance liquid chromatographic determination of cyclosporin A in human plasma and urine. *J Chromatogr Biomed Appl* 1980;182:454-8.
- Carruthers SG, Freeman DJ, Koegler JC, et al. Simplified liquid chromatographic analysis for cyclosporin A and comparison with radioimmunoassay. *Clin Chem* 1983;29:180-3.
- Kahn GC, Shaw LM, Kane MD. Routine monitoring of cyclosporine in whole blood and in kidney tissue using high performance liquid chromatography. *J Anal Toxicol* 1986;10:28-34.
- Moyer TP, Johnson P, Faynor SM, Sterioff S. Cyclosporine: a review of drug monitoring problems and presentation of a simple, accurate liquid chromatographic procedure that solves these problems. *Clin Biochem* 1986;19:3-9.
- Kabra PM, Wall JH. Improved liquid chromatographic analysis of cyclosporine in whole blood with solid phase (Bond-Elut) extraction. *J Liq Chromatogr* 1987;10:477-90.
- Bowers LD, Singh J. A gradient HPLC method for quantitation of cyclosporine and its metabolites in blood and bile. *J Liq Chromatogr* 1987;10:411-20.
- Klintmalm G, Sawe J, Ringden O, et al. Cyclosporine plasma levels in renal transplant patients. Association with renal toxicity and allograft rejection. *Transplantation* 1985;39:132-7.
- Vine W, Olson KN, Bowers LD. Therapeutic drug monitoring of cyclosporine by high performance liquid chromatography in high volume. *Transplant Proc* 1988;20(Suppl 2):354-6.
- Carey RN, Garber CC. Evaluation of methods. In: Kaplan LA, Pesce AJ, eds. *Clinical chemistry: theory, analysis and correlations*, 2nd ed. St. Louis: Mosby, 1989;290-310.
- Giesbrecht E, Soldin S, Wong PY. A rapid, reliable high-performance liquid chromatographic micromethod for the measurement of cyclosporine in whole blood. *Ther Drug Monit* 1989;11:332-6.
- Bowers LD. Cyclosporine analysis by HPLC: precision, accuracy and minimum detectable quantity. *Transplant Proc* 1990;22:1150-4.
- Shaw LM. Advances in cyclosporine pharmacology, measurement and therapeutic monitoring [Review]. *Clin Chem* 1989;35:1299-308.
- Maurer G, Loosli HR, Schreier E, et al. Disposition of cyclosporine in several animal species and man. *Drug Metab Dispos* 1984;12:120-6.
- Copeland KR, Yatscoff RW, McKenna RM. Immunosuppressive activity of cyclosporine metabolites compared and characterized by mass spectroscopy and nuclear magnetic resonance. *Clin Chem* 1990;36:225-9.
- Abecassis M, Wong PY, Mason N, et al. Cyclosporin A metabolites suppress T-cell proliferation by concanavalin A and in a mixed lymphocyte reaction. *Can J Surg* 1988;31:145-50.
- Wallenmacq PE, Lhoest G, Latinne D, et al. Isolation, characterization and in vitro activity in human cyclosporin A metabolites. *Transplant Proc* 1989;21:906-10.
- Wang PC, Hartman NR, Venkataramanan R, et al. Isolation of 10 cyclosporine metabolites from human bile. *Drug Metab Dispos* 1989;17:292-6.
- Meier GP, Park SB, Yee GC. Isolation and identification of a novel human metabolite of cyclosporin A, dihydro-CsA-M-17. *Drug Metab Dispos* 1990;18:68-71.
- Rosel T, LeGrue SJ, Liehr J, et al. Isolation and in vitro characterization of a novel immunosuppressive cyclosporine metabolite. *Transplantation* 1989;47:149-56.
- Henricsson S. A sulfate conjugate of cyclosporin. *Pharmacol Toxicol* 1990;66:53-5.
- Bowers LD, Covey T, Henion JD. Tandem quadrupole mass spectral characterization of a cyclic undecapeptide, cyclosporine, and selected analogs. *Biomed Environ Mass Spectrom* 1990;in press.
- Rosano TG, Freed BM, Cerilli J, et al. Immunosuppressive metabolites of cyclosporine in the blood of renal allograft recipients. *Transplantation* 1986;42:262-7.
- Freed BM, Rosano TG, Lempert N. In vitro immunosuppressive properties of cyclosporine metabolites. *Transplantation* 1987;43:123-7.
- Schlitt HJ, Christians V, Wonigeit K, et al. Immunosuppressive activity of cyclosporine metabolites in vitro. *Transplant Proc* 1987;19:248-51.
- Ryffel B, Foxwell BMJ, Mihatsch MJ, et al. Biological significance of cyclosporine metabolites. *Transplant Proc* 1988;20(Suppl 2):575-84.
- Zeevi A, Venkataramanan R, Burckart G, et al. Sensitivity of activated human lymphocytes to cyclosporine and its metabolites. *Hum Immunol* 1988;21:143-53.
- Heidecke CD, Nicolaus C, Schafer B, et al. Immunosuppressive activities of cyclosporine metabolites M-17 and M-21 within a bioassay based on inhibition of interleukin-2 production. *Transplant Proc* 1989;21:839.
- Cole E, Chung F, Wong PY, et al. Toxic effects on renal cells in culture—a comparison of cyclosporin A and its metabolites. *Transplant Proc* 1989;21:943-5.
- Copeland KR, Yatscoff RW. Immunosuppressive activity and toxicity of cyclosporine metabolites characterized by mass spectroscopy and nuclear magnetic resonance. *Transplant Proc* 1990;22:1146-9.
- Boosterli UA, Bouis P, Brouillard J-F, et al. In vivo toxicity assessment of cyclosporin A and its analogs in a primary rat hepatocyte culture model. *Toxicol Appl Pharmacol* 1988;96:212-21.

49. Bowers LD. Studies of cyclosporine and metabolite toxicity in renal and hepatocyte culture systems. *Transplant Proc* 1990;27(Suppl 3):1135-6.
50. Donatsch P, Rickenbacher U, Ryffel B, Brouillard J-F. Cyclosporine metabolites—their potential to cause adverse reactions in the rat. *Transplant Proc* 1990;27(Suppl 3):1137-40.
51. Rosano TG, Pell MA, Freed BM, et al. Cyclosporine and metabolites in blood from renal allograft recipients with nephrotoxicity, rejection or good renal function. Comparative high-performance liquid chromatography and monoclonal radioimmunoassay studies. *Transplant Proc* 1988;20(Suppl 2):330-8.
52. Lensmeyer GL, Wiebe DA, Carlson IH. Identification and analysis of nine metabolites of cyclosporine in whole blood by liquid chromatography. 2: Comparison of patients' results. *Clin Chem* 1987;33:1851-5.
53. Yee GC, Gmur DJ, Meier P. Measurement of cyclosporine metabolite concentrations with a new column-switching high-performance liquid chromatographic assay. *Transplant Proc* 1988;20(Suppl 2):585-90.
54. Wang CP, Burckart GJ, Ptachcinski RJ. Cyclosporine metabolite concentrations in the blood of liver, heart, kidney and bone marrow transplant patients. *Transplant Proc* 1988;20(Suppl 2):591-6.
55. Awni WM, Kasiske B, Heim-Duthoy K. Changes in the pharmacokinetics of cyclosporine and three of its metabolites in renal transplant patients early in the post-transplant period. *Transplant Proc* 1988;20(Suppl 2):623-4.
56. Haven MC, Sobeski LM, Ear RA, et al. Cyclosporine concentrations in blood after liver transplantation: correlation of immunoassay results with clinical events. *Clin Chem* 1989;35:564-8.
57. Wonigeit K, Kohlhaw K, Winkler M, Schaefer O, Pichlmayer R. Cyclosporine monitoring in liver allograft recipients. Two distinct patterns of blood level derangements associated with nephrotoxicity. *Transplant Proc* 1990;27(Suppl 3):1305-11.
58. Irschik E, Tilg H, Niederwieser D, Gastl G, Huber C, Margreiter R. Cyclosporin blood levels do correlate with clinical complications. *Lancet* 1984;ii:692-3.
59. Henry ML, Bowers VD, Fanning WJ, et al. Cyclosporine levels are not helpful. *Transplant Proc* 1988;20(Suppl 2):419-21.
60. Sommer BG, Sing DE, Henry ML, et al. Serum cyclosporine kinetic profile. Failure to correlate with nephrotoxicity or rejection episodes following sequential immunotherapy for renal transplantation. *Transplantation* 1988;45:86-90.
61. Kahan BD, Gibbons S, Jarowenko M, et al. The value of serial serum trough cyclosporine levels in human renal transplantation. *Transplant Proc* 1986;16:1195-9.
62. Keown PA, Stiller CR, Laupacis AL, et al. The effects and side effects of cyclosporine: relationship to drug pharmacokinetics. *Transplant Proc* 1982;14:659-61.
63. Holt DW, Marsden JT, Johnston A, et al. Blood cyclosporine concentrations and renal allograft dysfunction. *Br Med J* 1986;293:1057-9.
64. Moyer TP, Post GR, Sterioff S, Anderson CF. Cyclosporine nephrotoxicity is minimized by adjusting dosage on the basis of drug concentration in blood. *Mayo Clin Proc* 1988;63:241-7.
65. Macris MP, Ford EG, Van Buren CT, Frazier OH. Predictors of severe renal dysfunction after heart transplantation and intravenous cyclosporine therapy. *J Heart Transplant* 1989;8:444-8.
66. Trull A, Hue K, Tan K, et al. Cross-correlation of cyclosporin concentrations and biochemical measures of kidney and liver function in heart and heart-lung transplant recipients. *Clin Chem* 1990;36:1474-8.
67. Krom RAF, Wiesner RH, Rettke SR, et al. The first 100 liver transplantations at the Mayo Clinic. *Mayo Clin Proc* 1989;64:84-94.
68. Shaw LM, Audet PR, Grossman RA, Fields L, Lensmeyer GL, Dafoe DC. Adjustments of cyclosporine dosage in renal transplant patients based on concentration measured specifically in whole blood: clinical outcome, results, and diagnostic utility. *Transplant Proc* 1990;22:1267-73.
69. Grevel J, Napoli KL, Gibbons S, Kahan BD. Area-under-the-curve monitoring of cyclosporine therapy: performance of different assay methods and their target concentrations. *Ther Drug Monit* 1990;12:8-15.
70. Copeland KR, Yatscoff RW, Jeffery J, Rush D. Longitudinal monitoring of cyclosporine using a radioimmunoassay specific for the parent drug. *Transplantation* 1987;47:899-902.
71. Sibley R, Rynasiewicz J, Ferguson RM, et al. Morphology of cyclosporine nephrotoxicity and acute rejection in patients immunosuppressed with cyclosporine and prednisone. *Surgery* 1983;94:225-31.
72. Taube D, Neiled G, Hobby P, Holt D, Welsh K, Cameron JS. A comparison of the clinical, histological, cytologic, and biochemical features of renal transplant rejection, cyclosporin A nephrotoxicity, and stable renal function. *Transplant Proc* 1985;17:179-88.
73. Solez K, McGraw DJ, Beschoner WE, et al. Reflections on use of the renal biopsy as the gold standard in distinguishing transplant rejection from cyclosporine nephrotoxicity. *Transplant Proc* 1985;17(Suppl 1):123-33.
74. Myers BD, Ross J, Newton L, et al. Cyclosporine-associated chronic nephropathy. *N Engl J Med* 1984;311:699-705.
75. Mihatsch MJ, Thiel G, Ryffel B. Morphological diagnosis of cyclosporine nephrotoxicity. *Semin Diagn Pathol* 1988;5:104-21.

The Proceedings of the Canadian Consensus Meeting on Cyclosporine Monitoring will be published in their entirety in the February 1991 issue of *Clinical Biochemistry*.