ECE 351 - 52

Signals and Systems 1 Lab 9

Submitted By:
Owen Blair

Contents

0.1	H(jw) Magnitude Calculation	2
	0.1.1 Original $H(j\omega)$ Function	2
	0.1.2 Hand Calculation	2
0.2	H(jw) Phase Calculation	2

Owen Blair

0.1 H(jw) Magnitude Calculation

0.1.1 Original $H(j\omega)$ Function

$$\frac{\frac{j\omega}{RC}}{(j\omega)^2 + \frac{j\omega}{RC} + \frac{1}{LC}}$$

0.1.2 Hand Calculation

$$|H(j\omega)| = \frac{\sqrt{\left(\frac{\omega}{RC}\right)^2}}{\sqrt{\omega^2 + \left(\frac{\omega}{RC}\right)^2 + \left(\frac{1}{LC}\right)^2}}$$
$$|H(j\omega)| = \frac{\left(\frac{\omega}{RC}\right)}{\sqrt{\omega^2 + \left(\frac{\omega}{RC}\right)^2 + \left(\frac{1}{LC}\right)^2}}$$

0.2 H(jw) Phase Calculation

$$\angle H(j\omega) = \angle numerator(j\omega) - \angle denominator(j\omega)$$

$$\angle H(j\omega) = tan^{-1}(\frac{\omega}{RC}) - tan^{-1}(\frac{\frac{\omega}{RC}}{\omega^2 + \frac{1}{LC}})$$